



# LATTICE PLATONIC SOLIDS AND THEIR EHRHART POLYNOMIAL

E. J. IONASCU

ABSTRACT. First, we calculate the Ehrhart polynomial associated with an arbitrary cube with integer coordinates for its vertices. Then, we use this result to derive relationships between the Ehrhart polynomials for regular lattice tetrahedra and those for regular lattice octahedra. These relations allow one to reduce the calculation of these polynomials to only one coefficient.

## 1. INTRODUCTION

In the 1960's, Eugène Ehrhart ([14], [15]) proved that given a  $d$ -dimensional compact simplicial complex in  $\mathbb{R}^n$  ( $1 \leq d \leq n$ ), denoted here generically by  $\mathcal{P}$ , whose vertices are in the lattice  $\mathbb{Z}^n$ , there exists a polynomial  $L(\mathcal{P}, t) \in \mathbb{Q}[t]$  of degree  $d$ , associated with  $\mathcal{P}$  and satisfying

$$(1) \quad L(\mathcal{P}, t) = \text{the cardinality of } \{t\mathcal{P}\} \cap \mathbb{Z}^n, \quad t \in \mathbb{N}.$$

It is known that

$$L(\mathcal{P}, t) = \text{Vol}(\mathcal{P})t^n + \frac{1}{2} \text{Vol}(\partial\mathcal{P})t^{n-1} + \dots + \chi(\mathcal{P}),$$

---

Received July 11, 2012; revised September 20, 2012.

2010 *Mathematics Subject Classification*. Primary 52C07, 05A15, 68R05.

*Key words and phrases*. Ehrhart polynomial; linear Diophantine equations; lattice points; cubes; tetrahedra; octahedra.




Go back

Full Screen

Close

Quit



where  $\text{Vol}(\mathcal{P})$  is the usual volume of  $\mathcal{P}$ ,  $\text{Vol}(\partial\mathcal{P})$  is the surface area of  $\mathcal{P}$  normalized with respect to the sublattice on each face of  $\mathcal{P}$  and  $\chi(\mathcal{P})$  is the Euler characteristic of  $\mathcal{P}$ . In general, the other coefficients are less understandable, but significant progress has been done (see [5], [27] and [28]).

In [13], Eugène Ehrhart classified the regular convex polyhedra in  $\mathbb{Z}^3$ . It turns out that only cubes, regular tetrahedra and regular octahedra can be embedded in the usual integer lattice. We arrived at the same result in [23] using a construction of these polyhedra from equilateral triangles. This led us to the following simple description of all cubes in  $\mathbb{Z}^3$ . If we take an odd positive integer, say  $d$ , and a primitive solution of the Diophantine equation  $a^2 + b^2 + c^2 = 3d^2$  ( $\gcd(a, b, c) = 1$ ), then there are equilateral triangles in any plane having equation  $ax + by + cz = f$ , which can be parameterized in terms of two integers  $m$  and  $n$  (see [18], [19] and [22]). The side-lengths of such a triangle are equal to  $d\sqrt{2(m^2 + mn + n^2)}$ . In order to rise in space from such a triangle to form a regular tetrahedron, we need to satisfy the necessary and sufficient condition

$$(2) \quad m^2 + mn + n^2 = k^2 \text{ for some odd } k \in \mathbb{N}.$$

If (2) is satisfied, there are two possibilities. If  $k$  is a multiple of 3, then we can complete the triangle in both sides of the plane to a regular tetrahedron in  $\mathbb{Z}^3$ , and if  $k$  is not divisible by 3, then we can complete the triangle in exactly one side to form a regular tetrahedron in  $\mathbb{Z}^3$  (see Figure 1). Every such regular tetrahedron can then be completed to a cube in  $\mathbb{Z}^3$  with side-lengths equal to  $dk$ . Every regular octahedron in  $\mathbb{Z}^3$  is the dual of the double of a cube in  $\mathbb{Z}^3$ . We will make these constructions very specific in the last section.

It is natural to ask the question that we think Ehrhart himself asked: “What is the form that the polynomial in (1) takes for these regular lattice polyhedra?”. The purpose of this paper is to answer this question for cubes (in a very simple way), and give some partial answers for regular tetrahedra and octahedra.



Go back

Full Screen

Close

Quit




Figure 1. Expand an equilateral triangle.

For completeness and due credit, we include Ehrhart's idea in [16] to characterize all cubes in  $\mathbb{Z}^3$ . This is based on a theorem of Olinde Rodrigues: *The set of 3-by-3 orthogonal matrices can be given by four real parameters  $a, b, c, d$ , not simultaneously zero, as follows*

$$(3) \quad \frac{\pm 1}{a^2 + b^2 + c^2 + d^2} \begin{bmatrix} a^2 + b^2 - c^2 - d^2 & 2(bc + da) & 2(bd - ca) \\ 2(bc - da) & a^2 - b^2 + c^2 - d^2 & 2(cd + ba) \\ 2(bd + ca) & 2(cd - ba) & a^2 - b^2 - c^2 + d^2 \end{bmatrix}.$$

It is clear that every cube in  $\mathbb{Z}^3$  can be translated in such way that a vertex becomes the origin and the three vectors defined by the three sides starting from the origin give an orthogonal




Go back

Full Screen

Close

Quit



basis for  $\mathbb{R}^3$ . Hence, we can construct a 3-by-3 orthogonal matrix from these vectors which has rational entries. Conversely, we can construct a cube in  $\mathbb{Z}^3$  from such an orthogonal matrix which has rational entries. In what follows we will do this association so that the vectors (points) are determined by the rows. The construction here is to take four integers  $a, b, c$  and  $d$  in (3), simplify by whatever is possible and then get rid of the denominators to obtain the three vectors with integer coordinates that determine the cube. This construction is similar to the classical parametrization of the Heronian triangles.

Our approach to the classification allows us to start in terms of the side lengths. However, Ehrhart's construction is useful answering other questions about these objects. For instance, we can see that there are such cubes of any side length (other than the trivial ones, multiples of the unit cube) since every natural number can be written as a sum of four perfect squares. It turns out that there are only odd number side lengths for irreducible cubes, i.e., a cube which is not an integer multiple of a smaller cube in  $\mathbb{Z}^3$ .

Let us begin with some of the smallest irreducible cubes. We introduce them here by orthogonal matrices with rational entries and define up to the usual symmetries of the space (equivalent classes relative to the 48-order subgroup of all orthogonal matrices with entries 0 or  $\pm 1$ , denoted by  $\mathcal{S}_o$ ). As we mentioned before, this will make a difference, the cubes are essentially determined by the rows. Obviously, the Ehrhart polynomials are identical for all cubes in the same equivalence class (left or right).

We will denote the Ehrhart polynomial for an irreducible cube  $C_\ell$  of side-length  $\ell = 2k - 1$ ,  $k \in \mathbb{N}$ , by  $L(C_\ell, t)$ . From the general theory we have

$$(4) \quad L(C_\ell, t) = \ell^3 t^3 + \lambda_1 t^2 + \lambda_2 t + 1, \quad t \in \mathbb{N},$$

where  $\lambda_1$  is half the sum of the areas of the faces of the cube  $C_\ell$ , each face being normalized by the area of a fundamental domain of the sublattice contained in that face. The coefficient  $\lambda_2$  is in




Go back

Full Screen

Close

Quit



general a problem (see, for example [6]), but in this case it takes a simple form as we will show in Section 3.

For the unit cube  $C_1 = I$  (the identity matrix), obviously,  $L(C_1, t) = (t+1)^3$ . There is only one cube (right or left equivalence classes modulo  $\mathcal{S}_o$ ) for each  $\ell = 2k - 1$  for  $k = 1, 2, 3, 4, 5$  and 6:

$$C_1 = I \qquad C_3 := \frac{1}{3} \begin{bmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{bmatrix},$$

$$C_5 := \frac{1}{5} \begin{bmatrix} 4 & 3 & 0 \\ 3 & -4 & 0 \\ 0 & 0 & 5 \end{bmatrix}, \qquad C_7 := \frac{1}{7} \begin{bmatrix} -2 & 6 & 3 \\ 3 & -2 & 6 \\ 6 & 3 & -2 \end{bmatrix},$$

$$C_9 := \frac{1}{9} \begin{bmatrix} 7 & 4 & -4 \\ 4 & 1 & 8 \\ -4 & 8 & 1 \end{bmatrix}, \qquad C_{11} := \frac{1}{11} \begin{bmatrix} 2 & 9 & 6 \\ 9 & 2 & -6 \\ 6 & -6 & 7 \end{bmatrix}.$$

For  $k = 7$  we have  $C_{13} := \frac{1}{13} \begin{bmatrix} -3 & 12 & 4 \\ 4 & -3 & 12 \\ 12 & 4 & -3 \end{bmatrix}$ , and an extra orthogonal matrix:

$$\hat{C}_{13} := \frac{1}{13} \begin{bmatrix} 5 & 12 & 0 \\ 12 & -5 & 0 \\ 0 & 0 & 13 \end{bmatrix}.$$



Go back

Full Screen

Close

Quit



One peculiar thing about the Ehrhart polynomials associated with these cubes so far is that there is an unexpected factor in their factorization:

$$\begin{aligned}
 L(C_3, t) &= (3t + 1)(9t^2 + 1), & L(C_5, t) &= (5t + 1)(25t^2 + 2t + 1), \\
 L(C_7, t) &= (7t + 1)(49t^2 - 4t + 1), & L(C_9, t) &= (9t + 1)(81t^2 - 6t + 1), \\
 L(C_{11}, t) &= (11t + 1)(121t^2 - 8t + 1), & L(C_{13}, t) &= (13t + 1)(169t^2 - 10t + 1),
 \end{aligned}$$

and

$$L(\hat{C}_{13}, t) = (13t + 1)(169t^2 + 2t + 1).$$

This suggests that

$$(5) \quad L(C_\ell, t) = (\ell t + 1)(\ell^2 t^2 + \alpha t + 1), \quad t \in \mathbb{N}, \text{ and some } \alpha \in \mathbb{Z}.$$

We can easily prove that this is indeed the case for cubes of a special form like  $C_5$  and  $\hat{C}_{13}$  above. Let us consider a primitive Pythagorean triple  $(a, b, c)$  with  $a^2 + b^2 = c^2$ . In the  $xy$ -plane, we construct the square with vertices  $O(0, 0, 0)$ ,  $A(a, b, 0)$ ,  $B(a - b, a + b, 0)$ , and  $C(-b, a, 0)$  (Figure 2). We then translate this face along the vector  $\vec{c}k$  to form a cube of side-lengths equal to  $c$ . Let us denote this cube by  $C_{a,b,c}$ . It is easy to argue that  $(a, b, c)$  is primitive because we have no lattice points on the sides of  $OABC$ , other than its vertices. The coefficient  $\lambda_1$  in (4) is equal to  $\frac{1}{2}(c^2 + c^2 + 4c)$  because two of the faces have to be normalized by 1 and four of the faces have to be normalized by  $c(\frac{c}{c}) = c$ . By Pick's theorem, applied to  $OABC$ , we have

$$\begin{aligned}
 c^2 &= \frac{\#\{\text{points on the sides}\}}{2} + \#\{\text{interior points of OABC}\} - 1 \\
 &= \#\{\text{interior points of OABC}\} + 1.
 \end{aligned}$$



Go back

Full Screen

Close

Quit




Figure 2. One face of the cube.

Hence the number of lattice points in the interior of  $OABC$  is  $c^2 - 1$ . Therefore the number of lattice points in  $C_{a,b,c}$  is  $(c + 1)(c^2 + 3) = c^3 + c^2 + 3c + 3$ . The polynomial

$$L(C_{a,b,c}, t) = c^3 t^3 + (c^2 + 2c)t^2 + (c + 2)t + 1 = (ct + 1)(c^2 t^2 + 2t + 1)$$



Go back

Full Screen

Close

Quit



satisfies exactly the condition  $L(C_{a,b,c}, 1) = (c + 1)(c^2 + 3)$ . So we have shown that (5) is true for infinitely many cubes  $C_\ell$ .

**Proposition 1.1.** *Given a primitive Pythagorean triple,  $a^2 + b^2 = c^2$ , the cubes in the class of*

$$C_{a,b,c} := \frac{1}{c} \begin{bmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & c \end{bmatrix} \text{ have the same Ehrhart polynomial given by}$$

$$L(C_c, t) = (ct + 1)(c^2t^2 + 2t + 1), \quad t \in \mathbb{N}.$$

This proposition follows easily from the general theory since the polytope is a product of a square and a segment. The general formula is proved in Section 3. Section 2 is basically dealing with the second coefficient in (4). In Section 4, we look at the Ehrhart polynomial for regular tetrahedra and regular octahedra with lattice vertices.

## 2. THE COEFFICIENT $\lambda_1$

Let us prove the following well known lemma (see the acknowledgement note) which we include here for completeness.

**Lemma 2.1.** *For  $n \in \mathbb{N}$ ,  $n \geq 2$ , let  $a_1, a_2, \dots, a_n$  be  $n$  integers such that  $\gcd(a_1, a_2, \dots, a_n) = 1$ . Then the determinant of the lattice  $\mathcal{L}$  of points  $(x_1, x_2, \dots, x_n) \in \mathbb{Z}^n$  in the hyperplane  $a_1x_1 + \dots + a_nx_n = 0$  is given by  $\sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$ .*

*Proof.* We define  $p = a_1^2 + a_2^2 + \dots + a_n^2$  and consider the sublattice  $L$  of points  $(x_1, x_2, \dots, x_n) \in \mathbb{Z}^n$  such that  $a_1x_1 + \dots + a_nx_n = 0 \pmod{p}$ . Since  $\gcd(a_1, a_2, \dots, a_n) = 1$ , the index of  $L$  in  $\mathbb{Z}^n$  is  $p$  and hence the determinant of  $L$  is  $p$ . On the other hand, a basis for  $L$  can be obtained by appending a basis for the lattice  $\mathcal{L}$  by the vector  $(a_1, a_2, \dots, a_n)$  whose length is  $\sqrt{p}$  and which is perpendicular to all other basis vectors. Therefore, the determinant of the lattice  $\mathcal{L}$  is  $\frac{p}{\sqrt{p}} = \sqrt{p}$ .  $\square$




Go back

Full Screen

Close

Quit





Let us now assume that we have an arbitrary cube in  $\mathbb{Z}^3$ ,

$$(6) \quad C_\ell = \frac{1}{\ell} \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_2 & c_3 \end{bmatrix},$$

with  $a_i$ ,  $b_i$  and  $c_i$  integers such that  $a_i a_j + b_i b_j + c_i c_j = \delta_{i,j} \ell^2$  for all  $i, j$  in  $\{1, 2, 3\}$ . We define  $d_i := \gcd(a_i, b_i, c_i)$ . It is clear that  $d_i$  are divisors of  $\ell$ . Let us also introduce the numbers  $d'_i = \frac{\ell}{d_i}$ ,  $i = 1, 2, 3$ . Then, we have the following expression for the first coefficient in (4).

**Theorem 2.2.** *The coefficient  $\lambda_1$  is given by*

$$(7) \quad \lambda_1 = \ell(d_1 + d_2 + d_3) \text{ where } d_i := \gcd(a_i, b_i, c_i), \quad i \in \{1, 2, 3\}.$$

*Proof.* We use Lemma 2.1 for each of the faces of the cube. Opposite faces will have the same contribution. Say we take a face containing the points  $(a_1, b_1, c_1)$  and  $(a_2, b_2, c_2)$ . The irreducible normal vector to this face is clearly  $\frac{1}{d_3}(a_3, b_3, c_3)$ . The area of a fundamental domain here is given by  $\sqrt{\frac{1}{d_3^2}(a_3^2 + b_3^2 + c_3^2)} = D_3$ . By the general theory  $\lambda_1 = \frac{1}{2}(2\frac{\ell^2}{d_1} + 2\frac{\ell^2}{d_2} + 2\frac{\ell^2}{d_3}) = \ell(d_1 + d_2 + d_3)$ .  $\square$

Naturally, at this point, the question is whether or not it is possible to have all of the  $d_i$ 's bigger than one. It turns out that this is possible and as before, in our line of similar investigations, the first  $\ell$  is  $\ell = 1105 = 5(13)(17)$

$$(8) \quad C_{1105} = \frac{1}{1105} \begin{bmatrix} -65 & 156 & 1092 \\ 420 & 1015 & -120 \\ 1020 & -408 & 119 \end{bmatrix}.$$



Go back

Full Screen

Close

Quit



**Corollary 2.3.** For a matrix  $C_\ell$  as in (6) such that the  $C_\ell^{-1}$  is in the same equivalence class modulo  $\mathcal{S}_o$ , we have

$$(9) \quad d_1 + d_2 + d_3 = \gcd(a_1, a_2, a_3) + \gcd(b_1, b_2, b_3) + \gcd(c_1, c_2, c_3).$$

*Proof.* The Ehrhart polynomial must be the same for the corresponding cubes in the same equivalence class.  $\square$

We believe that this corollary applies to all  $\ell < 1105$ , and of course to a lot of other cases, but we do not have a proof of this fact. A counterexample to the hypothesis of this corollary is given by the matrix given in (8). In this case,  $d_1 + d_2 + d_3 = 35$  and  $\gcd(a_1, a_2, a_3) + \gcd(b_1, b_2, b_3) + \gcd(c_1, c_2, c_3) = 7$ .

### 3. THE COEFFICIENT $\lambda_2$

The main idea in calculating the coefficient  $a_2$  is to take advantage of the fact that every cube defined by (6) can be used to form a fundamental domain (wandering set)  $W$  for the space, under integer translations along the vectors  $\vec{\alpha} = (a_1, b_1, c_1)$ ,  $\vec{\beta} = (a_2, b_2, c_2)$  and  $\vec{\gamma} = (a_3, b_3, c_3)$ , i.e.,

$$\mathbb{R}^3 = \bigcup_{i,j,k \in \mathbb{Z}} (W + i\vec{\alpha} + j\vec{\beta} + k\vec{\gamma}),$$

where  $\overset{\circ}{\bigcup}$  means a union of mutually disjoint sets.

The fundamental domain  $W$  that we will consider here, associated with a generic cube as in Figure 3, is the set of all points formed by the interior points of the cube to which we add the points of the faces OAEC, OADB and OBFC except the (closed) edges AD, DB, BF, FC, CE, and EA. It is easy to see that such a set is indeed a wandering set. We were informed that this notion



Go back

Full Screen

Close

Quit





**Theorem 3.1** (Ehrhart-Macdonald Reciprocity Law). *Given a compact simplicial lattice complex  $P$  (as before) of dimension  $n$ , then*

$$L(\overset{\circ}{P}, t) = (-1)^n L(P, -t), \quad t \in \mathbb{N},$$

where  $\overset{\circ}{P}$  denotes the interior of  $P$ , as usual.

With the notation from the previous section, we have the following result.

**Theorem 3.2.** *The coefficient  $\lambda_2$  is equal to  $d_1 + d_2 + d_3$ .*

*Proof.* Let us denote by  $k$  the number of lattice points in  $W$ . For  $n \in \mathbb{N}$ , the number of lattice points in

$$\bigcup_{i,j,k \in \{1,2,\dots,n\}} (W + i\vec{\alpha} + j\vec{\beta} + k\vec{\gamma}),$$

is equal to  $n^3k$ . On the other hand, this number is equal to  $L(C_\ell, n) + K$ , where  $K$  is the number of lattice points on three big faces of  $nC$ . It is easy to see that  $K$  is  $O(n^2)$ , and so  $k = \lim_{n \rightarrow \infty} \frac{1}{n^3}(L(C_\ell, n) + O(n^2)) = \ell^3$ .

Hence, according to Theorem 3.1, the number of lattice points in the interior of  $C_\ell$  is  $\ell^3 - \lambda_1 + \lambda_2 - 1$ . So the number of lattice points on the boundary of  $C_\ell$  is  $2\lambda_1 + 2$ . Let us denote by  $\sigma$  the number of lattice points on the interior of the sides  $\overline{OA}$ ,  $\overline{OB}$  and  $\overline{OC}$ . Then we have

$$2\lambda_1 + 2 = 2[k - (\ell^3 - \lambda_1 + \lambda_2 - 1)] + 2\sigma + 6 \Rightarrow \lambda_2 = \sigma + 3.$$

Since  $\sigma = (d_1 - 1) + (d_2 - 1) + (d_3 - 1)$ , the claim follows. □

Putting these facts together we obtain the following theorem.



Go back

Full Screen

Close

Quit



**Theorem 3.3.** Given a cube  $C_\ell$  constructed from a matrix as in (6), its Ehrhart polynomial is given by

$$(20) \quad L(C_\ell, t) = (\ell t + 1)[\ell^2 t^2 + (d_1 + d_2 + d_3 - \ell)t + 1], \quad t \in \mathbb{N}.$$

There are some natural questions at this point. One of them is: “What is the maximum number of lattice points that can be contained in a lattice cube of side lengths  $\ell$ ?” We have the following corollary to the above theorem.

**Corollary 3.4.** Given a cube  $C_\ell$  constructed from a matrix as in (6), the maximum of lattice points inside or on the boundary of this cube cannot be more than  $(\ell + 1)^3$ . This value is attained for the cube  $\ell C_1$ .

*Proof.* Since  $d_i$  is a divisor of  $\ell$ , we have  $d_i \leq \ell$ , so the corollary follows from (10). □

What is the maximum of lattice points contained in an irreducible cubes of sides  $\ell$ ? This is a more complicated problem which depends heavily on  $\ell$  and relates to the number of irreducible cubes (their Ehrhart polynomials, in fact) of sides  $\ell$ .

#### 4. REGULAR TETRAHEDRA AND REGULAR OCTAHEDRA

We remind the reader that a cube in space (Figure 4) is determined by an orthogonal matrix as in (6) by taking its vertices  $O$  (the origin),  $A, B, C, D, E, F$  and  $G$  whose position vectors are  $\overrightarrow{OA} = \vec{\alpha} = (a_1, b_1, c_1)$ ,  $\overrightarrow{OB} = \vec{\beta} = (a_2, b_2, c_2)$ ,  $\overrightarrow{OC} = \vec{\gamma} = (a_3, b_3, c_3)$ ,  $\overrightarrow{OD} = \vec{\alpha} + \vec{\beta}$ ,  $\overrightarrow{OE} = \vec{\beta} + \vec{\gamma}$ ,  $\overrightarrow{OF} = \vec{\gamma} + \vec{\alpha}$  and  $\overrightarrow{OG} = \vec{\alpha} + \vec{\beta} + \vec{\gamma}$ .

In [23], we rediscovered Ehrhart’s characterization ([13]) of all regular polyhedra which can be imbedded in  $\mathbb{Z}^3$ . Only the cubes, the tetrahedra and octahedra exist in  $\mathbb{Z}^3$  and there are infinitely many in each class. We have constructed all of these equilateral triangles. In general, once a




Go back

Full Screen

Close

Quit



tetrahedron is constructed, this can be always completed to a cube. Vice versa, for a cube given by (6), there are two regular tetrahedra as shown in Figure 4, which are in the same equivalence class, modulo the orthogonal matrices with entries  $\pm 1$ , denoted earlier by  $\mathcal{S}_0$ . Regular octahedra can be obtained by doubling the coordinates of the cube  $C_\ell$  and then taking the centers of each face. This procedure is exhaustive. An octahedron in the same class can be obtained by simply taking the vertices whose position vectors are  $\pm\vec{\alpha}$ ,  $\pm\vec{\beta}$  and  $\pm\vec{\gamma}$ . We will use the notations  $T_\ell$  and  $O_\ell$  for the tetrahedra and octahedra constructed this way from  $C_\ell$ . Since we are interested in irreducible  $T_\ell$  and  $O_\ell$ , we may assume that  $\ell$  is odd. The  $T_\ell$  and  $O_\ell$  have side-lengths equal to  $\ell\sqrt{2}$ . From the general Ehrhart theory (see [3]), we have

$$(11) \quad L(T_\ell, t) = \frac{\ell^3}{3}t^3 + \mu_1 t^2 + \mu_2 t + 1, \quad L(O_\ell, t) = \frac{4\ell^3}{3}t^3 + \nu_1 t^2 + \nu_2 t + 1.$$

Let us first look at some of the examples of the smallest side-lengths.

$$T_1 := \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}, \quad O_1 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ with}$$

$$L(T_1, t) = \frac{t^3}{3} + t^2 + \frac{5t}{3} + 1 \quad \text{and} \quad L(O_1, t) = \frac{4}{3}t^3 + 2t^2 + \frac{8t}{3} + 1.$$

For the next side-lengths,

$$T_3 := \begin{bmatrix} 1 & 1 & 4 \\ 1 & 4 & 1 \\ 4 & 1 & 1 \end{bmatrix}, \quad O_3 := \begin{bmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{bmatrix} \text{ with}$$

$$L(T_3, t) = 9t^3 + \frac{9}{2}t^2 + \frac{13t}{2} + 1 \quad \text{and} \quad L(O_3, t) = 36t^3 + 9t^2 - t + 1.$$



Go back

Full Screen

Close

Quit




Figure 4. A cube determines two tetrahedra.

These polynomials were computed with the help of a computer.

#### 4.1. The coefficients $\mu_1$ and $\nu_1$

From the general theory we know that these coefficients can be computed in terms of the areas of faces and normalized by the area of the fundamental domains of the sub-lattice of  $\mathbb{Z}^3$  corresponding




Go back

Full Screen

Close

Quit



to that face. Since the number of faces for  $O_\ell$  is twice as big as  $T_\ell$  and basically the parallel faces are in the same equivalence class, we have  $\nu_1 = 2\mu_1$ .

Let us introduce the divisors

$$(12) \quad \begin{aligned} D_1 &= \gcd(a_1 + a_2 + a_3, b_1 + b_2 + b_3, c_1 + c_2 + c_3), \\ D_2 &= \gcd(a_1 + a_2 - a_3, b_1 + b_2 - b_3, c_1 + c_2 - c_3), \\ D_3 &= \gcd(a_1 - a_2 + a_3, b_1 - b_2 + b_3, c_1 - c_2 + c_3) \quad \text{and} \\ D_4 &= \gcd(-a_1 + a_2 + a_3, -b_1 + b_2 + b_3, -c_1 + c_2 + c_3). \end{aligned}$$

Let us observe that the vectors  $\vec{\alpha} + \vec{\beta} + \vec{\gamma}$ ,  $\vec{\alpha} + \vec{\beta} - \vec{\gamma}$ ,  $\vec{\alpha} - \vec{\beta} + \vec{\gamma}$ ,  $-\vec{\alpha} + \vec{\beta} + \vec{\gamma}$  are vectors normal to the faces of the  $T_\ell$ . By Lemma 2.1, we see that the area of each fundamental domain corresponding to a face of  $T_\ell$  is given by one of the numbers  $\frac{\ell\sqrt{3}}{D_i}$ .

**Proposition 4.1.** *The coefficients  $\mu_1$  and  $\nu_1$  in (11) are given by*

$$(13) \quad \mu_1 = \frac{\nu_1}{2} = \frac{\ell(D_1 + D_2 + D_3 + D_4)}{4}.$$

This explains the coefficients of  $t^2$  in the next examples which were obtained by brute force counting using Maple:

$$T_5 := \begin{bmatrix} 7 & -1 & 0 \\ 4 & 3 & 5 \\ 3 & -4 & 5 \end{bmatrix}, \quad O_5 := \begin{bmatrix} 4 & 3 & 0 \\ 3 & -4 & 0 \\ 0 & 0 & 5 \end{bmatrix} \quad \text{with}$$
$$L(T_5, t) = \frac{125}{3}t^3 + 5t^2 + \frac{1}{3}t + 1 \quad \text{and} \quad L(O_5, t) = \frac{500}{3}t^3 + 10t^2 + \frac{16}{3}t + 1.$$




Go back

Full Screen

Close

Quit





## 4.2. The coefficients $\mu_2$ and $\nu_2$

Let us observe that the cube in Figure 4 can be decomposed into four triangular pyramids OABC, DABG, FCGB and EGCA, which can be translated and some reflected into the origin to form half of  $O_\ell$  and the regular tetrahedron  $T_\ell$ . We remind the reader of a notation we used in the proof of Theorem 3.2 where we denoted by  $\sigma$  the number of lattice points on the interior of the edges  $\overline{OA}$ ,  $\overline{OB}$  and  $\overline{OC}$ . We showed that  $\sigma = d_1 + d_2 + d_3 - 3$ .

Let us balance the number  $M$  of the lattice interior points of  $C_\ell$  using the above decomposition. According to Theorem 3.3, and Theorem 3.1 we have

$$\begin{aligned} M &= -L(C_\ell, -1) = (\ell - 1)(\ell^2 - d_1 - d_2 - d_3 + \ell + 1) \\ &= \ell^3 - (d_1 + d_2 + d_3)(\ell - 1) - 1. \end{aligned}$$

Part of the lattice points counted in  $M$  are in the regular tetrahedron which are counted by  $L(T_\ell, 1) = \frac{\ell^3}{3} + \mu_1 + \mu_2 + 1$ , from which we need to subtract the number of lattice points on the interior of its sides, which we will denote by  $\tau$  and subtract 4 for its vertices. The rest of the points counted in  $M$  is in the interior of the four pyramids. If we multiply this number by two and add the number of lattice points in the interior of the cube faces of the cube less  $\tau$ , we get the number of interior points of  $O_\ell$  minus  $2\sigma + 1$ . The number of lattice interior points of the cube faces is equal to  $2\lambda_1 + 2 - 4\sigma - 8$ . In other words, we have

$$\begin{aligned} &2\left(M - \frac{\ell^3}{3} - \mu_1 - \mu_2 - 1 + \tau + 4\right) + 2\lambda_1 + 2 - 4\sigma - 8 - \tau \\ &= \frac{4\ell^3}{3} - \nu_1 + \nu_2 - 1 - (2\sigma + 1). \end{aligned}$$

Taking into account that  $\nu_1 = 2\mu_1$  and  $\lambda_1 = (d_1 + d_2 + d_3)\ell = (\sigma + 3)\ell$ , this can be simplified to

$$(14) \quad \nu_2 + 2\mu_2 = 6 + \tau.$$




Go back

Full Screen

Close

Quit



We close this section concluding what we have shown.

**Theorem 4.1.** *For a regular tetrahedron  $T_\ell$  and a regular octahedron  $O_\ell$  constructed as before from an orthogonal matrix with rational coefficients as in (6), the coefficients  $\mu_2$  and  $\nu_2$  in (11) satisfy*

$$(15) \quad \nu_2 + 2\mu_2 = (d_1 + d_2 + d_3 + d_4 + d_5 + d_6),$$

where  $d_1, d_2, d_3$  are defined as before and  $d_4 = \gcd(a_1 - a_2, b_1 - b_2, c_1 - c_2)$ ,  $d_5 = \gcd(a_1 - a_3, b_1 - b_3, c_1 - c_3)$  and  $d_6 = \gcd(a_3 - a_2, b_3 - b_2, c_3 - c_2)$ .

We have tried to find another relation that will help us find the two coefficients but it seems there is not an easy way to avoid, what are called in [3], the building blocks of the lattice-point enumeration, the Dedekind sums. These numbers require a little more computational power and we are wonder if a shortcut doesn't really exist. One would expect that the answer to our questions for such regular objects is encoded in the coordinates of their vertices in a relatively simple way. We leave this problem to the interest of a reader.

**Acknowledgment.** The author wishes to express his sincere appreciation for the anonymous referee's constructive report which informed us about the general formulation of Lemma 2.1 and its proof.

1. Athanasiadis C. A., *Ehrhart polynomials, simplicial polytopes, magic squares and a conjecture of Stanley*, J. reine angew. Math. **583** (2005), 163–174.
2. Barvinok A., *Computing the Ehrhart quasi-polynomial of a rational simplex*, Mathematics of Computation, **75** (2006), 1449–1466.




Go back

Full Screen

Close

Quit



3. Beck M. and Robins S., *Computing the Continuous Discretely: Integer-Point Enumeration in Polyhedra*, Undergraduate Texts in Mathematics, Springer-Verlag, New York, 2007; also available at <http://math.sfsu.edu/beck/ccd.html>
4. Beck M., Diaz R. and Robins S., *The Frobenius problem, rational polytopes, and Fourier-Dedekind sums*, Journal of Number Theory **96** (2002), 1–21.
5. Berline N. and Vergne M., *Local Euler-Maclaurin formula for polytopes*, Mosc. Math. J. **7(3)** (2007), 355–386.
6. Diaz R. and Robins S., *The Ehrhart polynomial of a lattice polytope*, Ann. of Math. **145** (1997) 503–518.
7. Braun B. J., *Ehrhart Theory for Lattice Polytopes*, Ph. D. Thesis, 2007.
8. ———, *An Ehrhart Series Formula For Reflexive Polytopes*, Electronic Journal of Combinatorics, **13(1)** (2006), N 15.
9. Braun B. J. and Develin M., *Ehrhart Polynomial Roots and Stanley’s Non-negativity Theorem*, Integer Points in Polyhedra–Geometry, Number Theory, Representation Theory, Algebra, Optimization, Statistics, Contemporary Mathematics **452** (2008), 67–78.
10. Brion M. and Vergne M., *Lattice Points in Simple Polytopes*, Journal of the American Mathematical Society **10(2)** (1997), 371–392.
11. Chandler R. and Ionascu E. J., *A characterization of all equilateral triangles in  $\mathbb{Z}^3$* , Integers, Art. **8** (2008), A19.
12. Chen B., *Weight Functions, Double Reciprocity Laws, and Formulas for Lattice Polyhedra*, Proceedings of the National Academy of Sciences of the United States of America **95(16)** (1998), 9093–9098.
13. Ehrhart E., *Sur les polygones et les polyèdres réguliers entiers*, Enseignement Math. **5(2)** (1959), 81–85.
14. ———, *Sur les polyèdres rationnels homothétiques à  $n$  dimensions*, C. R. Acad. Sci. Paris 254 (1962), 616–618.
15. ———, *Sur un problème de géométrie diophantienne linéaire. I. Polyèdres et réseaux*, J. reine angew. Math. **226** (1967), 1–29.
16. ———, *Solution to problem 6179 [1977, 744]*, Amer. Math. Monthly 1980, 826–827.
17. Schoenberg I. J., *Regular Simplices and Quadratic Forms*, J. London Math. Soc. **12** (1937), 48–55.
18. Ionascu E. J., *A parametrization of equilateral triangles having integer coordinates*, Journal of Integer Sequences, **10** (2007), 09.6.7.
19. ———, *Counting all equilateral triangles in  $\{0, 1, 2, \dots, n\}^3$* , Acta Math. Univ. Comenianae, **LXXVII(1)**, (2008), 129–140.
20. ———, *A characterization of regular tetrahedra in  $\mathbb{Z}^3$* , J. Number Theory, **129** (2009), 1066–1074.




Go back

Full Screen

Close

Quit



21. ———, *Regular tetrahedra with integer coordinates of their vertices*, Acta Math. Univ. Comenianae, **LXXX(2)** (2011) 161–170.
22. Ionascu E. J. and Chandler R., *A characterization of all equilateral triangles in  $\mathbb{Z}^3$* , Integers, **8** (2008), A19.
23. Ionascu E. J. and Markov A., *Platonic solids in  $\mathbb{Z}^3$* , J. Number Theory 131(1) (2011), 138–145.
24. Ionascu E. J. and Obando R., *Cubes in  $\{0, 1, 2, \dots, n\}^3$* , to appear in “Integers”, arXiv:1003.4569
25. Ionascu E. J., *Ehrhart’s polynomial for equilateral triangles in  $\mathbb{Z}^3$* , to appear in Australas. J. Combinatorics, (2012)
26. Liu F., *Contributions to the theory of Ehrhart polynomials*, Ph. D. Thesis 2006.
27. Pommersheim J., *Toric varieties, lattice points, and Dedekind sums*, Math. Ann. **295** (1993), 1–24.
28. Pommersheim J. and Thomas H., *Cycles representing the Todd class of a toric variety*, J. Amer. Math. Soc. **17(4)** (2004), 983–994 (electronic).
29. Sam S., *A bijective proof for a theorem of Ehrhart*, Amer. Math. Monthly **116(8)** (2009), 688–701.
30. Sloane Neil J. A., *The On-Line Encyclopedia of Integer Sequences*, 2005,  
<http://www.research.att.com/~njas/sequences/>.
31. Stapledon A., *Equivariant Ehrhart theory*, Adv. Math. **226(4)** (2011), 3622–3654.

E. J. Ionascu, Department of Mathematics Columbus State University 4225 University Avenue Columbus, GA 31907, e-mail: [ionascu@columbusstate.edu](mailto:ionascu@columbusstate.edu), [math@ejionascu.ro](mailto:math@ejionascu.ro)



Go back

Full Screen

Close

Quit