

# On Gaussian Polynomials and Content Ideal

Chahrazade Bakkari    Najib Mahdou

*Department of Mathematics, Faculty of Science and Technology of Fez  
Box 2202, University S. M. Ben Abdellah Fez, Morocco*

**Abstract.** Loper and Roitman proved that every integral domain satisfies the property “each Gaussian polynomial has locally principal content”. In this paper, we study this property in a ring with zero-divisors, and give then a class of such rings which does not satisfying this property and another class of rings with zero-divisors satisfying this property.

Keywords: Gaussian polynomial, content ideal, locally principal, trivial ring extension, direct product

## 1. Introduction

Throughout this paper all rings are assumed to be commutative with identity elements and all modules are unital.

Let  $R$  be a commutative ring. We say that an ideal is regular if it contains a regular element, i.e., a non-zerodivisor element. A ring  $R$  is called locally has a property  $(P)$  if each localisation of  $R$  by a maximal ideal of  $R$  has the property  $(P)$ .

The content  $C(f)$  of a polynomial  $f \in R[X]$  is the ideal of  $R$  generated by all coefficients of  $f$ . One of its properties is that  $C(\cdot)$  is semi-multiplicative, that is  $C(fg) \subseteq C(f)C(g)$ , and a polynomial  $f \in R[X]$  is said to be Gaussian over  $R$  if  $C(fg) = C(f)C(g)$ , for every polynomial  $g \in R[X]$ . A polynomial  $f \in R[X]$  is Gaussian provided  $C(f)$  is locally principal by [7, Remark 1.1]. Our guiding question is the converse of this property, that is “each regular Gaussian polynomial has locally principal content”. Notice for convenience that the conjecture has a local character since the Gaussian condition is a local property (i.e., a polynomial

is Gaussian over a ring  $R$  if and only if its image is Gaussian over  $R_M$  for each maximal ideal  $M$  of  $R$ ), [12, Lemma 5].

Significant progress has been made on this conjecture. Glaz and Vasconcelos proved it for normal Noetherian domains [6]. Then Heinzer and Huneke established this conjecture over locally approximately Gorenstein rings and over locally Noetherian domains [7, Theorem 1.5 and Corollary 3.4]. Recently, Loper and Roitman established the conjecture for (locally) domains [11, Theorem 4], and then Lucas extended their result to arbitrary rings by restricting to polynomials with regular content [12, Theorem 3.6]. Finally, in [1], by using pullbacks, the authors construct a new class of rings that are not locally domains, nor locally Noetherians, and satisfy this conjecture. Let us note that Heinzer and Huneke, in [7, Remark 1.6], give an example showing that the conjecture is false in general.

Let  $A$  be a ring,  $E$  be an  $A$ -module and  $R := A \rtimes E$  be the set of pairs  $(a, e)$  with pairwise addition and multiplication given by  $(a, e)(b, f) = (ab, af + be)$ .  $R$  is called the trivial ring extension of  $A$  by  $E$ . Note that a prime (respectively, maximal) ideal of  $R$  has the form  $P \rtimes E$  (respectively,  $M \rtimes E$ ) where  $P$  (respectively,  $M$ ) is a prime (respectively, maximal) ideal of  $A$  (by [9, Theorem 25.1]).

Considerable work has been concerned with trivial ring extensions. Part of it has been summarized in Glaz's book [4] and Huckaba's book (where  $R$  is called the idealization of  $E$  by  $A$ ) [9].

The goal of this work is to exhibit a class of rings (with zero-divisors) that does not satisfy the property "each Gaussian polynomial has locally principal content" and a second class of rings with zero-divisors satisfying this property. For this purpose, we study the transfer of this property to trivial ring extension and direct product.

## 2. Main results

This section develops a result of the transfer of the property "each Gaussian polynomial has locally principal content" for a particular context of trivial ring extensions. And so, we will construct a class of rings (with zero-divisors) that does not satisfy the property "each Gaussian polynomial has locally principal content".

**Theorem 2.1.** *Let  $(A, M)$  be a local ring which is not a field such that  $M^2 = 0$  and let  $R := A \rtimes E$  be the trivial ring extension of  $A$  by  $E$ , where  $E$  is a free  $A$ -module. Then,  $R$  does not satisfy the property "each Gaussian polynomial has locally principal content" in the following cases:*

- (1)  $\text{rank}_A(E) = 1$  and  $M$  is not principal,
- (2)  $\text{rank}_A(E) \geq 2$ .

Before proving Theorem 2.1, we establish the following Lemma.

**Lemma 2.2.** *Let  $(A, M)$  be a local ring which is not a field such that  $M^2 = 0$  and let  $R := A \rtimes E$  be the trivial ring extension of  $A$  by  $E$ , where  $E$  is an  $A$ -module. Assume that there exists  $a, b \in E - \{0\}$  such that  $Ma = Mb = 0$  and the  $A$ -module generated by  $\{a, b\}$  is not principal. Then,  $R$  does not satisfy the property “each Gaussian polynomial has locally principal content”.*

*Proof.* Under the hypothesis of Lemma 2.2, remark that  $R$  is local with maximal ideal  $M \rtimes E$  by [9, Theorem 25.1] since  $(A, M)$  is local. Let  $f = (0, a) + (0, b)X \in R[X]$ . Our aim is to show that  $f$  is Gaussian and  $C(f)$  is not (locally) principal. We claim that  $f$  is Gaussian. Indeed, let  $g \in R[X]$ . We may assume that  $g \in (M \rtimes E)[X]$  (since if  $g \notin (M \rtimes E)[X]$ , then  $C(g) = R$  and so  $C(fg) = C(f)C(g)$ ). Hence,  $C(f)C(g) = [R(0, a) + R(0, b)]C(g) \subseteq [R(0, a) + R(0, b)][M \rtimes E] = 0$  since  $aM = bM = 0$  and so  $C(fg) = C(f)C(g) = 0$  which means that  $f$  is Gaussian.

We claim that  $C(f)$  is not principal. Deny. There exists  $(c, d) \in R$  such that  $C(f) := R(0, a) + R(0, b) (= 0 \rtimes (Aa + Ab)) = R(c, d)$  since  $R$  is local. Hence,  $c = 0$  and so  $Aa + Ab = Ad$ , a contradiction since the  $A$ -module generated by  $\{a, b\}$  is not principal. Therefore,  $C(f)$  is not locally principal and this completes the proof of Lemma 2.2.

*Proof of Theorem 2.1.* Let  $(A, M)$  be a local ring which is not a field such that  $M^2 = 0$  and let  $R := A \rtimes E$  be the trivial ring extension of  $A$  by  $E$ , where  $E$  is a free  $A$ -module.

1) Assume that  $\text{rank}_A(E) = 1$  and  $M$  is not principal. We may assume that  $E = A$ . Hence, there exists  $m_1, m_2 \in M - \{0\}$  such that  $m_2 \notin Am_1$ . Set  $f = (m_1, 0) + (0, m_2)X \in R[X]$ . Our aim is to show that  $f$  is Gaussian and  $C(f)$  is not (locally) principal.

We claim that  $f$  is a Gaussian polynomial. Let  $g = \sum_{i=0}^n (n_i, a_i)X^i \in R[X]$ .

If  $n_i \notin M$  for some  $i = 0, \dots, n$ , then  $n_i$  is invertible in  $A$  and so  $(n_i, a_i)$  is invertible in  $R$ . Hence,  $C(g) = R$  and so  $g$  is Gaussian; thus  $C(fg) = C(f)C(g)$ .

Assume that  $n_i \in M$  for each  $i = 0, \dots, n$ . We have  $fg = \sum_{i=0}^n (m_1, 0)(n_i, a_i)X^i +$

$$\sum_{i=0}^n (0, m_2)(n_i, a_i)X^{i+1} = \sum_{i=0}^n (0, m_1a_i)X^i$$

since  $m_1, m_2, n_i \in M$  and  $M^2 = 0$ . Hence

$$C(fg) = \sum_{i=0}^n R(0, m_1a_i).$$

On the other hand,  $C(f)C(g)$

$$= [R(m_1, 0) + R(0, m_2)][\sum_{i=0}^n R(n_i, a_i)] = \sum_{i=0}^n R(m_1, 0)(n_i, a_i) = \sum_{i=0}^n R(0, m_1a_i)$$

since  $m_1, m_2, n_i \in M$  and  $M^2 = 0$ . Hence,  $C(fg) = C(f)C(g)$  and so  $f$  is Gaussian.

We claim that  $C(f)$  is not principal (since  $R$  is local). Deny. Then  $C(f)(= R(m_1, 0) + R(0, m_2)) = R(n, e)$  for some  $n \in M$  and  $e \in A$  and so  $(m_1, 0) = (a, b)(n, e)$  for some  $(a, b) \in R$ .

If  $a \in M$ , then  $m_1 = an = 0$  since  $n \in M$ , a contradiction.

If  $a \notin M$ , then  $a$  is invertible in  $A$  and so  $(a, b)$  is invertible in  $R$ . Hence,  $R(m_1, 0) = R(n, e)(= R(m_1, 0) + R(0, m_2))$  and so  $(0, m_2) \in R(m_1, 0)$ . Therefore,  $(0, m_2) = (a, b)(0, m_1) = (0, am_1)$  and so  $m_2 \in Am_1$ , a contradiction. Therefore,  $C(f)$  is not (locally) principal.

2) Now, assume that  $\text{rank}_A(E) \geq 2$ . Let  $m \in M - \{0\}$ ,  $(e_i)_{i \in I}$  be a basis of the free  $A$ -module  $E$ ,  $a_i = me_i \in E$  for  $i = 1, 2$ . We have  $Ma_i = 0$  for each  $i = 1, 2$  since  $M^2 = 0$ . We claim that the  $A$ -module generated by  $(a_i)_{i=1,2}$  is not principal.

Assume that the  $A$ -module generated by  $(a_i)_{i=1,2}$  is principal, that is  $Aa_1 + Aa_2 = Ame_1 + Ame_2 = Af$  for some  $f \in E$ . Hence,  $f = b_1me_1 + b_2me_2$  for some  $b_i \in A$ , where  $i = 1, 2$ . But,  $a_1 \in Af$  implies that  $a_1(= me_1) = cf = cb_1me_1 + cb_2me_2$  for some  $c \in A$ . Thus,  $m = cb_1m$  and  $cb_2m = 0$  since  $(e_i)_{i \in I}$  is a basis of the free  $A$ -module  $E$ . Therefore,  $(1 - cb_1)m = 0$  and so  $1 - cb_1 \in M$  since  $(A, M)$  is a local ring and  $m \neq 0$ . Hence,  $cb_1 \notin M$  and so  $cb_1$  is invertible; in particular,  $c$  is invertible.

Hence, the equation  $cb_2m = 0$  implies that  $b_2m = 0$  (as  $c$  is invertible) and so  $b_2 \in M$  (since  $(A, M)$  is a local ring and  $m \neq 0$ ). Hence,  $f = b_1me_1 + b_2me_2 = b_1me_1$  as  $b_2, m \in M$  and  $M^2 = 0$ . But  $a_2(= me_2) \in Aa_1 + Aa_2 = Af$  implies that  $me_2 = df = db_1me_1$  for some  $d \in A$ ; so  $m = 0$  since  $(e_i)_{i \in I}$  is a basis of the free  $A$ -module  $E$ , a contradiction as  $m \neq 0$ .

Therefore, the  $A$ -module generated by  $(a_i)_{i=1,2}$  is not principal and Lemma 2.2 completes the proof of Theorem 2.1.

**Remark 2.3.** The hypothesis “ $M$  is not principal” in Theorem 2.1(1) is necessary. Indeed, let  $K$  be a field considered as a local ring with maximal ideal  $M = 0$ . Hence,  $R := K \times K$  is a local ring with unique proper ideal  $R(0, 1)(= 0 \times K)$ . So,  $R$  satisfies the property “each Gaussian polynomial has locally principal content”.

The goal of the following result is to construct a second class of rings that does not satisfy the property “each Gaussian polynomial has locally principal content”.

**Proposition 2.4.** *Let  $(A, M)$  be a local ring which is not a Bézout ring such that  $M^2 = 0$ . Then  $A$  does not satisfy the property “each Gaussian polynomial has locally principal content”.*

*Proof.* It suffices to show that there exists a polynomial  $f \in A[X]$  such that  $C(f)$  is not principal since  $A$  is local and Gaussian. For that, let's consider  $a, b \in A$  such that the ideal generated by  $\{a, b\}$  is not principal (since  $A$  is not a Bézout ring). Set  $f = a + bX \in A[X]$ . Therefore,  $C(f) := aA + bA$  is not principal and this completes the proof of Proposition 2.4.

The next result gives a second wide class of rings that does not satisfy the property “each Gaussian polynomial has locally principal content”.

**Corollary 2.5.** *Let  $(A, M)$  be a local ring which is not a field such that  $M^2 = 0$  and let  $R := A \rtimes E$ , where  $E$  is a nonzero  $A$ -module such that  $ME = 0$ . Then  $R$  does not satisfy the property “each Gaussian polynomial has locally principal content”.*

*Proof.* Remark that the ring  $R$  is local with maximal ideal  $M \rtimes E$  which satisfies  $(M \rtimes E)^2 = 0$  since  $M^2 = 0$  and  $ME = 0$ . Hence, each polynomial in  $R$  is Gaussian. It remains to show that  $R$  is not Bézout by Proposition 2.4 (since  $(M \rtimes E)^2 = 0$ ).

Let  $a \in M - \{0\}$  and  $e \in E - \{0\}$ . We claim that the ideal  $I$  generated by  $\{(a, 0), (0, e)\}$  is not principal. Deny. Assume that  $I := R(a, 0) + R(0, e) = R(b, h)$ , where  $(b, h) \in M \rtimes E$ . We claim that  $b \neq 0$ .

Indeed, if  $b = 0$ , then  $(a, 0) \in I = R(0, h)$  which implies that  $a = 0$ , a contradiction. Hence  $b \neq 0$ .

But  $(0, e) \in I = R(b, h)$ . Hence,  $(0, e) = (c, l)(b, h) = (cb, ch)$  for some  $(c, l) \in R$  (since  $b \in M$ ). Then,  $cb = 0$  and  $c \in M$  (since  $c \notin M$  implies that  $c$  is invertible and  $b = 0$ , a contradiction). Therefore,  $e = ch = 0$ , a contradiction. Then  $I$  is not a principal ideal of  $R$  and so  $R$  is not a Bézout ring and this completes the proof of Corollary 2.5.

Now, we will construct a wide class of rings satisfying the property “each Gaussian polynomial has locally principal content”. For this, we study the transfer of this property to finite direct products.

**Theorem 2.6.** *Let  $(R_i)_{i=1, \dots, n}$  be a family of rings. Then  $\prod_{i=1}^n R_i$  satisfies the property “each Gaussian polynomial has locally principal content” if and only if so does  $R_i$  for each  $i = 1, \dots, n$ .*

Before proving Theorem 2.6, we establish the following Lemma.

**Lemma 2.7.** *Let  $R$  be a ring and,  $h : R \rightarrow h(R)$  be a ring homomorphism, and  $f$  be a Gaussian polynomial in  $R[X]$ . Then the homomorphic image of  $f$  is a Gaussian polynomial in  $h(R)[X]$ .*

*Proof.* Let  $h : R \rightarrow h(R)$  be a ring homomorphism,  $f$  be a Gaussian polynomial, and let  $g = \sum_{i=0}^n a_i X^i \in R[X]$ . Let's remark first that,

$$\begin{aligned} C_{h(R)}(h(g)) &= \sum_{i=0}^n h(R)h(a_i) \\ &= \sum_{i=0}^n h(Ra_i) \\ &= h\left(\sum_{i=0}^n Ra_i\right) \\ &= h(C_R(g)). \end{aligned}$$

Hence, we have (since  $f$  is Gaussian):

$$\begin{aligned}
 C_{h(R)}(h(f)h(g)) &= C_{h(R)}(h(fg)) \\
 &= h(C_R(fg)) \\
 &= h(C_R(f)C_R(g)) \\
 &= h(C_R(f))h(C_R(g)) \\
 &= C_{h(R)}(h(f))C_{h(R)}(h(g)).
 \end{aligned}$$

As desired.

*Proof of Theorem 2.6.* We will prove the result for  $i = 1, 2$ , and the theorem will be established by induction on  $n$ . Let  $f = (f_1, f_2)$  be a Gaussian polynomial in  $(R_1 \times R_2)[X]$ , and  $M$  be a maximal ideal of  $R_1 \times R_2$ . Then  $M = m_1 \times R_2$  or  $M = R_1 \times m_2$  where  $m_i \in \text{Max}(R_i)$  for  $i = 1, 2$ .

We may assume that  $M = m_1 \times R_2$  (the case  $M = R_1 \times m_2$  is similar). We wish prove that  $C_{R_1 \times R_2}(f)_M$  is principal. But  $(R_1 \times R_2)_M$  is naturally isomorphic to  $(R_1)_{m_1}$  and  $C_{R_1 \times R_2}(f)_M$  is isomorphic to  $C(f_1)_{m_1}$ . Therefore,  $C_{R_1 \times R_2}(f)_M$  is principal since  $f_1$  is supposed Gaussian by Lemma 2.7 and so  $R_1 \times R_2$  satisfies the property “each Gaussian polynomial has locally principal content”.

Conversely, assume that the polynomial  $f_1$  is Gaussian in  $R_1[X]$  (it is the same for  $f_2$ ). We easily check that  $f := (f_1, 0)$  is Gaussian in  $(R_1 \times R_2)[X]$ . Let  $m_1 \in \text{Max}(R_1)$ . Therefore,  $(C_{R_1 \times R_2}(f))_{m_1 \times R_2}$  is principal since  $R_1 \times R_2$  satisfies the property “each Gaussian polynomial has locally principal content”. Hence,  $(C_{R_1}(f_1))_{m_1}$  is principal (since  $(R_1 \times R_2)_M$  is naturally isomorphic to  $(R_1)_{m_1}$  and  $C_{R_1 \times R_2}(f)_M$  is isomorphic to  $C(f_1)_{m_1}$ ), which means that  $C_{R_1}(f_1)$  is locally principal and this completes the proof of Theorem 2.6.

Now, we are able to construct a wide class of rings satisfying the property “each Gaussian polynomial has locally principal content”.

**Corollary 2.8.** *Let  $(R_i)_{i=1, \dots, n}$  be a family of domains. Then  $\prod_{i=1}^n R_i$  satisfies the property “each Gaussian polynomial has locally principal content”.*

*Proof.* By Theorem 2.5 and since every domain satisfies the property “each Gaussian polynomial has locally principal content” (by [11, Theorem 4]).

**Acknowledgements.** The authors would like to express their sincere thanks for the referee for his/her helpful suggestions and comments, which have greatly improved this paper.

## References

- [1] Bakkari, C.; Mahdou, N.: *Gaussian polynomials and content ideal in pull-backs*. Comm. Algebra **34**(8) (2006), 2727–2732. [Zbl 1101.13010](https://doi.org/10.1080/009278706005713010)

- [2] Bazzoni, S.; Glaz, S.: *Prüfer rings in “Multiplicative Ideal Theory in Commutative Algebra”*. In: J. W. Brewer (ed.) et al, Springer, New York 2006, 263–277. [Zbl 1115.13024](#)
- [3] Corso, A.; Heinzer, W.; Huneke, C.: *A generalized Dedekind-Mertens lemma and its converse*. Trans. Am. Math. Soc. **350** (1998), 5095–5109. [Zbl 0903.13001](#)
- [4] Glaz, S.: *Commutative coherent rings*. Lect. Notes Math. **1371**, Springer-Verlag, Berlin 1989. [Zbl 0745.13004](#)
- [5] Glaz, S.: *The weak dimensions of Gaussian rings*. Proc. Am. Math. Soc. **133**(9) (2005), 2507–2513. [Zbl 1077.13009](#)
- [6] Glaz, S.; Vasconcelos, W.: *The content of Gaussian polynomials*. J. Algebra **202** (1998), 1–9. [Zbl 0923.13007](#)
- [7] Heinzer, W.; Huneke, C.: *Gaussian polynomials and content ideals*. Proc. Am. Math. Soc. **125** (1997), 739–745. [Zbl 0860.13005](#)
- [8] Heinzer, W.; Huneke, C.: *The Dedekind-Mertens lemma and the content of polynomials*. Proc. Am. Math. Soc. **126** (1998), 1305–1309. [Zbl 0887.13005](#)
- [9] Huckaba, J. A.: *Commutative rings with zero divisors*. Marcel Dekker, New York 1988. [Zbl 0637.13001](#)
- [10] Kabbaj, S.; Mahdou, N.: *Trivial extensions defined by coherent-like conditions*. Commun. Algebra **32**(10) (2004), 3937–3953. [Zbl 1068.13002](#)
- [11] Loper, K. A.; Roitman, M.: *The content of a Gaussian polynomial is invertible*. Proc. Am. Math. Soc. **133**(5) (2004), 1267–1271. [Zbl 1137.13301](#)
- [12] Lucas, T. G.: *Gaussian polynomials and invertibility*. Proc. Am. Math. Soc. **133**(7) (2005), 1881–1886. [Zbl pre02150822](#)
- [13] Rotman, J. J.: *An introduction to homological algebra*. Academic Press, New York 1979. [Zbl 0441.18018](#)
- [14] Rush, D. E.: *The Dedekind-Mertens lemma and the contents of polynomials*. Proc. Am. Math. Soc. **128** (2000), 2879–2884. [Zbl 0974.13007](#)
- [15] Tsang, H.: *Gauss’s Lemma*. Ph.D. thesis, University of Chicago, Chicago 1965.

Received May 14, 2007