



Existence and rapid convergence results for nonlinear Caputo nabla fractional difference equations

Xiang Liu¹, Baoguo Jia¹, Lynn Erbe² and Allan Peterson ²

¹School of Mathematics, Sun Yat-Sen University, Guangzhou, 510275, China

²Department of Mathematics, University of Nebraska-Lincoln, Lincoln, NE 68588-0130, U.S.A.

Received 26 January 2017, appeared 24 June 2017

Communicated by Paul Eloe

Abstract. This paper is concerned with finding properties of solutions to initial value problems for nonlinear Caputo nabla fractional difference equations. We obtain existence and rapid convergence results for such equations by use of Schauder's fixed point theorem and the generalized quasi-linearization method, respectively. A numerical example is given to illustrate one of our rapid convergence results.


Keywords: Caputo nabla fractional difference equation, Schauder's fixed point theorem, generalized quasi-linearization, existence, rapid convergence.

2010 Mathematics Subject Classification: 39A12, 39A70.

1 Introduction

It is well known that there is a large quantity of research on integer-order difference equations. Since the study was begun very early, much classical content has been established, and we refer specifically to the monographs [2,17]. However, the study of fractional difference equations is quite recent. The basic theory of linear and nonlinear fractional difference equations can be found in [13–16]. Note that the theory of nonlinear fractional difference equations is not complete and the convergence of approximate solutions is one of the most studied problems. This has an important affect on the development of the qualitative theory.

Generalized quasi-linearization is an efficient method for constructing approximate solutions of nonlinear problems. This method originated in dynamic programming theory and was initially applied by Bellman and Kalaba [8]. A systematic development of the method to ordinary differential equations was provided by Lakshmikantham and Vatsala [18], and there are some generalized results of the method to various types of differential equations and we refer to the monographs [19,20], for functional differential equations [3,11], for impulsive equations [4,7], for partial differential equations [5,9,23], for differential equations on time scales [22], for fractional differential equations [10,21,26], for other types [12,24,25], and the references cited therein. For nonlinear fractional difference equations see the paper [6].

 Corresponding author. Email: apeterson1@math.unl.edu

In this paper, we attempt to extend the applications of generalized quasi-linearization with certain conditions on the forcing function, and study the rate of convergence of the approximate solutions for the nonlinear Caputo nabla fractional difference equation. In order to do this, we first prove the existence of solutions for such equations, and then using an appropriate iterative scheme, we obtain two monotone sequences which converge uniformly and rapidly to the solution of the problem. Finally, we provide a numerical example to illustrate the application of the obtained results.

2 Preliminary definitions

For the convenience of readers, we will list some relevant results here. We use the notation $\mathbb{N}_a := \{a, a + 1, a + 2, \dots\}$, where a is a real number. For the function $f : \mathbb{N}_a \rightarrow \mathbb{R}$, the backward difference or nabla operator is defined as $\nabla f(t) = f(t) - f(t - 1)$ for $t \in \mathbb{N}_{a+1}$ and the higher order differences are defined recursively by $\nabla^n f(t) = \nabla(\nabla^{n-1} f(t))$ for $t \in \mathbb{N}_{a+n}$, $n \in \mathbb{N}$. In addition, we take ∇^0 as the identity operator. We define the definite nabla integral of $f : \mathbb{N}_a \rightarrow \mathbb{R}$ by

$$\int_a^b f(s) \nabla s = \begin{cases} \sum_{s=a+1}^b f(s), & a < b, \\ 0, & a = b, \\ -\sum_{s=b+1}^a f(s), & a > b, \end{cases} \quad (2.1)$$

where $b \in \mathbb{N}_a$.

Definition 2.1 (See [15, Definition 3.4]). The (generalized) rising function is defined by

$$t^{\bar{r}} = \frac{\Gamma(t+r)}{\Gamma(t)} \quad (2.2)$$

for those values of t and r for which the right-hand side of (2.2) is defined. Also, we use the convention that if t is a nonpositive integer, but $t+r$ is not a nonpositive integer, then $t^{\bar{r}} = 0$. We then define the ν -th order Taylor monomials based at a (see [15, Definition 3.56] by

$$H_\nu(t, a) = \frac{(t-a)^{\bar{\nu}}}{\Gamma(\nu+1)},$$

for $\nu \neq -1, -2, \dots, t \in \mathbb{N}_a$.

For some important formulas for these Taylor fractional monomials see [15, Theorem 3.57 and Theorem 3.93].

Definition 2.2 (Nabla fractional sum [15, Definition 3.58]). Let $f : \mathbb{N}_a \rightarrow \mathbb{R}$ be given and assume $\nu > 0$. Then

$$\nabla_a^{-\nu} f(t) = \int_a^t H_{\nu-1}(t, \rho(s)) f(s) \nabla s, \quad t \in \mathbb{N}_a, \quad (2.3)$$

where $\rho(t) := t - 1$ and by convention $\nabla_a^{-\nu} f(a) = 0$.

Definition 2.3 (Nabla fractional difference [15, Definition 3.61]). Let $f : \mathbb{N}_a \rightarrow \mathbb{R}$, $\nu > 0$ be given, and let $N := \lceil \nu \rceil$, where $\lceil \cdot \rceil$ is the ceiling function. Then we define the ν -order nabla fractional difference operator $\nabla_a^\nu f(t)$ by

$$\nabla_a^\nu f(t) = \nabla^N \nabla_a^{-(N-\nu)} f(t), \quad t \in \mathbb{N}_{a+N}. \quad (2.4)$$

Definition 2.4 (Caputo nabla fractional difference [15, Definition 3.117]). Let $f : \mathbb{N}_a \rightarrow \mathbb{R}$, $\nu > 0$ be given, and let $N := \lceil \nu \rceil$. Then we define the ν -order Caputo nabla fractional difference operator $\nabla_{a^*}^\nu f(t)$ by

$$\nabla_{a^*}^\nu f(t) = \nabla_a^{-(N-\nu)} [\nabla^N f(t)], \quad t \in \mathbb{N}_{a+N}. \quad (2.5)$$

Now it follows from this definition that $\nabla_{a^*}^\nu c = 0$ for $\nu > 0$ with any constant c .

Lemma 2.5 (See [15, Definition 3.61 and Theorem 3.62]). Assume $f : \mathbb{N}_a \rightarrow \mathbb{R}$, $\nu > 0$, $\nu \notin \mathbb{N}_1$, and choose $N \in \mathbb{N}_1$ such that $N - 1 < \nu < N$. Then

$$\nabla_a^\nu f(t) = \int_a^t H_{-\nu-1}(t, \rho(s)) f(s) \nabla s, \quad t \in \mathbb{N}_a, \quad (2.6)$$

where $\rho(t) := t - 1$ and by convention $\nabla_a^\nu f(a) = 0$.

Lemma 2.6 (See [1]). Assume $f : \mathbb{N}_a \rightarrow \mathbb{R}$, for any $\nu > 0$, we have

$$\nabla_{a^*}^\nu f(t) = \nabla_a^\nu f(t) - \sum_{k=0}^{N-1} H_{-\nu+k}(t, a) \nabla^k f(a). \quad (2.7)$$

In particular, when $0 < \nu < 1$, we have

$$\nabla_{a^*}^\nu f(t) = \nabla_a^\nu f(t) - H_{-\nu}(t, a) f(a). \quad (2.8)$$

3 Existence and comparison results

Consider the following initial value problem (IVP) for a nonlinear Caputo nabla fractional difference equation

$$\begin{cases} \nabla_{a^*}^\nu x(t) = f(t, x(t)), & t \in \mathbb{N}_{a+1}^b, \\ x(a) = x_0, \end{cases} \quad (3.1)$$

where $f : \mathbb{N}_{a+1}^b \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous with respect to x , $x : \mathbb{N}_a \rightarrow \mathbb{R}$, and $0 < \nu < 1$.

In this paper, we define the norm of x on \mathbb{N}_a^b by $\|x\| = \max_{s \in \mathbb{N}_a^b} |x(s)|$. Throughout this paper, we use the notation $\frac{\partial^k f(t, x)}{\partial^k x} = f^{(k)}(t, x)$ ($k = 0, 1, 2, \dots$). We define the following set for convenience:

$$\Omega = \{(t, x) : \alpha_0(t) \leq x(t) \leq \beta_0(t), t \in \mathbb{N}_{a+1}^b\}.$$

where $\alpha_0(t)$ and $\beta_0(t)$ are defined on \mathbb{N}_a^b with $\alpha_0(t) \leq \beta_0(t)$ on \mathbb{N}_a^b .

Definition 3.1. The function $\alpha_0(t)$, $t \in \mathbb{N}_a^b$, is said to be a lower (an upper) solution of the IVP (3.1), if

$$\begin{cases} \nabla_{a^*}^\nu \alpha_0(t) \leq (\geq) f(t, \alpha_0(t)), & t \in \mathbb{N}_{a+1}^b, \\ \alpha_0(a) \leq (\geq) x_0. \end{cases} \quad (3.2)$$

Lemma 3.2. *The function $x(t)$ is a solution of the IVP (3.1) if and only if the function $x(t)$ has the following representation*

$$x(t) = x_0 + \sum_{s=a+1}^t H_{\nu-1}(t, \rho(s)) f(s, x(s)). \quad (3.3)$$

Proof. Applying the operator $\nabla_a^{-\nu}$ on both sides of the first equality of the IVP (3.1), we have

$$\nabla_a^{-\nu} [\nabla_{a^*}^{\nu} x(t)] = \nabla_a^{-\nu} f(t, x(t)),$$

which can be written as

$$\nabla_a^{-\nu} [\nabla_a^{-(1-\nu)} \nabla x(t)] = \nabla_a^{-\nu} f(t, x(t)).$$

That is,

$$\nabla^{-1} \nabla x(t) = \nabla_a^{-\nu} f(t, x(t)).$$

Then, we have

$$x(t) = x_0 + \sum_{s=a+1}^t H_{\nu-1}(t, \rho(s)) f(s, x(s)).$$

Conversely, assume that x has the representation (3.3). By means of (2.3), we obtain that (3.3) is equivalent to

$$x(t) = x_0 + \nabla_a^{-\nu} f(t, x(t)). \quad (3.4)$$

Applying $\nabla_{a^*}^{\nu}$ to both sides of (3.4), we get

$$\nabla_{a^*}^{\nu} x(t) = \nabla_{a^*}^{\nu} x_0 + \nabla_{a^*}^{\nu} [\nabla_a^{-\nu} f(t, x(t))].$$

Using (2.8), we obtain

$$\nabla_{a^*}^{\nu} x(t) = \nabla_{a^*}^{\nu} x_0 + \left(\nabla_a^{\nu} \nabla_a^{-\nu} f(t, x(t)) - H_{-\nu}(t, a) \nabla_a^{-\nu} f(a, x(a)) \right).$$

Thus, we have

$$\nabla_{a^*}^{\nu} x(t) = f(t, x(t)).$$

The proof is complete. □

Now we present an existence result for the IVP (3.1), which we will use in our main results. Since the proof is a standard application of Schauder's fixed point theorem we will omit the proof of this lemma.

Lemma 3.3. *Assume that*

(H_{3.1}) *the function $f : \bar{R} \rightarrow \mathbb{R}$ is continuous with respect to x , $|f(t, x)| \leq Q$ on \bar{R} , $D = H_{\nu}(b, a)$, and $D \leq \frac{M}{Q}$, where*

$$\bar{R} = \{(t, x) : t \in \mathbb{N}_{a+1}^b, \|x - x_0\| \leq M\}.$$

Then the IVP (3.1) has a solution.

Lemma 3.4. *Assume that*

(H_{3.2}) *the function $f : \Omega \rightarrow \mathbb{R}$ is continuous in its second variable.*

(H_{3.3}) *the functions $\alpha_0, \beta_0 : \mathbb{N}_a^b \rightarrow \mathbb{R}$ are lower and upper solutions respectively of the IVP (3.1) such that $\alpha_0(t) \leq \beta_0(t)$ on \mathbb{N}_a^b ;*

Then there exists a solution $x(t)$ of the IVP (3.1) satisfying $\alpha_0(t) \leq x(t) \leq \beta_0(t)$ on \mathbb{N}_a^b .

Proof. Let $P : \mathbb{N}_{a+1}^b \times \mathbb{R} \rightarrow \mathbb{R}$ be defined by $P(t, x) = \max\{\alpha_0(t), \min\{x, \beta_0(t)\}\}$. Then $f(t, P(t, x))$ defines an extension of f to $\mathbb{N}_{a+1}^b \times \mathbb{R}$, which is bounded and continuous with respect its second variable on \mathbb{N}_{a+1}^b . Therefore, by Lemma 3.3, $\nabla_{a^*}^\nu x(t) = f(t, P(t, x))$, $x(a) = x_0$ has a solution on \mathbb{N}_a^b .

To complete the proof, we need to show that $\alpha_0(t) \leq x(t) \leq \beta_0(t)$ on \mathbb{N}_a^b . We now show that $x(t) \leq \beta_0(t)$ on \mathbb{N}_a^b . Clearly, $x(a) \leq \beta_0(a)$, we now only need to show $x(t) \leq \beta_0(t)$ on \mathbb{N}_{a+1}^b . If it is not true, there exists a point $c \in \mathbb{N}_{a+1}^b$ such that $x(t) - \beta_0(t)$ has a positive maximum, that is,

$$x(c) - \beta_0(c) = \max\{x(t) - \beta_0(t) : t \in \mathbb{N}_{a+1}^b\} > 0,$$

and

$$x(t) - \beta_0(t) \leq x(c) - \beta_0(c) \quad \text{on } \mathbb{N}_{a+1}^c.$$

First, we will show that $\nabla_{a^*}^\nu x(c) > \nabla_{a^*}^\nu \beta_0(c)$, that is,

$$\nabla_a^\nu x(c) - H_{-\nu}(c, a)x(a) > \nabla_a^\nu \beta_0(c) - H_{-\nu}(c, a)\beta_0(a).$$

Since $x(a) \leq \beta_0(a)$ and

$$H_{-\nu}(c, a) = \frac{(c-a)^{-\bar{\nu}}}{\Gamma(1-\nu)} = \frac{\Gamma(c-a-\nu)}{\Gamma(c-a)\Gamma(1-\nu)} > 0, \quad c \in \mathbb{N}_{a+1}^b,$$

we have

$$-H_{-\nu}(c, a)x(a) \geq -H_{-\nu}(c, a)\beta_0(a).$$

Next, we show $\nabla_a^\nu x(c) > \nabla_a^\nu \beta_0(c)$. In view of the fact that $\frac{(1)^{-\bar{\nu}-1}}{\Gamma(-\nu)} = \frac{\Gamma(-\nu)}{\Gamma(-\nu)\Gamma(1)} = 1$ and $H_{-\nu}(c, a) > 0$, it follows that

$$\begin{aligned} \nabla_a^\nu x(c) - \nabla_a^\nu \beta_0(c) &= \frac{1}{\Gamma(-\nu)} \sum_{s=a+1}^c (c-\rho(s))^{-\bar{\nu}-1} (x(s) - \beta_0(s)) \\ &= \left[\frac{1}{\Gamma(-\nu)} \sum_{s=a+1}^{c-1} (c-\rho(s))^{-\bar{\nu}-1} (x(s) - \beta_0(s)) \right] + (x(c) - \beta_0(c)) \\ &\geq \left[1 + \frac{1}{\Gamma(-\nu)} \sum_{s=a+1}^{c-1} (c-\rho(s))^{-\bar{\nu}-1} \right] (x(c) - \beta_0(c)) \\ &= \left[\sum_{s=a+1}^c H_{-\nu-1}(c, \rho(s)) \right] (x(c) - \beta_0(c)) \\ &= H_{-\nu}(c, a)(x(c) - \beta_0(c)) > 0. \end{aligned}$$

Then, we have $\nabla_a^\nu x(c) > \nabla_a^\nu \beta_0(c)$. Thus, we conclude that $\nabla_{a^*}^\nu x(c) > \nabla_{a^*}^\nu \beta_0(c)$.

On the other hand, due to $x(c) - \beta_0(c) > 0$, so $P(c, x(c)) = \beta_0(c)$. Hence

$$\nabla_{a^*}^\nu \beta_0(c) \geq f(c, P(c, x(c))) = \nabla_{a^*}^\nu x(c),$$

which is a contradiction. Thus, we obtain $x(t) \leq \beta_0(t)$ on \mathbb{N}_a^b . Similarly, we can show that $\alpha_0(t) \leq x(t)$ on \mathbb{N}_a^b . Therefore, it follows that $x(t)$ is actually a solution of IVP (3.1). The proof is complete. \square

For the convenience of readers, we give a result for the linear Caputo nabla fractional difference equation.

Consider the Caputo nabla fractional difference inequality

$$\nabla_a^\nu x(t) - Cx(t) \leq 0, \quad x(a) \leq 0, \quad t \in \mathbb{N}_{a+1}^b. \quad (3.5)$$

Lemma 3.5. *Assume that*

$(H_{3.4})$ *the positive constant C satisfies $CH_\nu(b, a) < 1$.*

Then $x(t) \leq 0$ on \mathbb{N}_a^b .

Proof. Setting $x(t) = \nabla_a^{-\nu} y(t)$, according to the Definition 2.2, we have

$$\begin{aligned} x(t) = \nabla_a^{-\nu} y(t) &= \int_a^t H_{\nu-1}(t, \rho(s)) y(s) \nabla s \\ &= \sum_{s=a+1}^t \frac{(t - \rho(s))^{\overline{\nu-1}}}{\Gamma(\nu)} y(s) \\ &= \sum_{s=a+1}^t \frac{\Gamma(t - s + \nu)}{\Gamma(\nu) \Gamma(t - s + 1)} y(s). \end{aligned} \quad (3.6)$$

We get from (3.6) that $y(t) \leq 0$ implies $x(t) \leq 0$ on \mathbb{N}_{a+1}^b , so we only need to prove $y(t) \leq 0$ on \mathbb{N}_{a+1}^b . If this is false, there exists a $c \in \mathbb{N}_{a+1}^b$ such that $y(c) = \max\{y(t) : t \in \mathbb{N}_{a+1}^b\} > 0$, and $y(t) \leq y(c)$ on \mathbb{N}_{a+1}^c . It follows from (2.8) that (3.5) is equivalent to

$$\nabla_a^\nu x(t) - H_{-\nu}(t, a)x(a) - Cx(t) \leq 0. \quad (3.7)$$

Letting $x(t) = \nabla_a^{-\nu} y(t)$ in (3.7) yields

$$\nabla_a^\nu \nabla_a^{-\nu} y(t) - C \nabla_a^{-\nu} y(t) \leq 0,$$

which can be written as

$$y(t) \leq \frac{C}{\Gamma(\nu)} \sum_{s=a+1}^t (t - \rho(s))^{\overline{\nu-1}} y(s).$$

Hence, we have

$$\begin{aligned} y(c) &\leq \frac{C}{\Gamma(\nu)} \sum_{s=a+1}^c (c - \rho(s))^{\overline{\nu-1}} y(s) \\ &\leq \frac{C}{\Gamma(\nu)} \left(\sum_{s=a+1}^c (c - \rho(s))^{\overline{\nu-1}} \right) y(c) \\ &= Cy(c)H_\nu(c, a), \end{aligned}$$

that is,

$$(1 - CH_\nu(c, a))y(c) \leq 0.$$

Since $y(c) > 0$, so we have $(1 - CH_\nu(c, a)) \leq 0$.

On the other hand, from the condition $(H_{3.4})$ and the increasing property of the function $H_\nu(t, a)$, we have $(1 - CH_\nu(c, a)) > 0$, which is a contradiction. Then, we have $y(t) \leq 0$ on \mathbb{N}_{a+1}^b . Hence, we conclude that $x(t) \leq 0$ on \mathbb{N}_a^b . The proof is complete. \square

4 Rapid convergence

In this section, we consider the IVP (3.1) with $f(t, x) = f_1(t, x) + f_2(t, x)$. We show that the convergence of the sequences of successive approximations is of order m where m is $2k + 1$ or $2k$ ($k \geq 1$) by applying the method of generalized quasi-linearization for nonlinear Caputo nabla fractional difference equations.

Theorem 4.1. *Assume that the conditions $(H_{3,3})$ – $(H_{3,4})$ hold, and*

(A_{4.1}) the functions $f_1, f_2 : \Omega \rightarrow \mathbb{R}$ are such that $f_1^{(i)}(t, x), f_2^{(i)}(t, x)$ ($i = 0, 1, \dots, 2k$) exist, are continuous in the second variable, and for $C_1 > 0, C_2 > 0, C = C_1 + C_2$,

$$f_1^{(1)}(t, x) \leq C_1, f_2^{(1)}(t, x) \leq C_2 \quad \text{on } \Omega;$$

(A_{4.2}) there exist $M_1, M_2 > 0$ such that for $x_1 \geq x_2, y_1 \geq y_2$ the functions $f_1^{(2k)}(t, x), f_2^{(2k)}(t, x)$ satisfy the following conditions:

$$\begin{aligned} 0 &\leq f_1^{(2k)}(t, x_1) - f_1^{(2k)}(t, x_2) \leq M_1(x_1 - x_2) \quad \text{on } \Omega, \\ 0 &\geq f_2^{(2k)}(t, y_1) - f_2^{(2k)}(t, y_2) \geq -M_2(y_1 - y_2) \quad \text{on } \Omega. \end{aligned}$$

Then there exist two monotone sequences $\{\alpha_n(t)\}, \{\beta_n(t)\}, n \geq 0$ which converge uniformly and monotonically to a solution of the IVP (3.1) and the convergence is of order $2k + 1$.

Proof. From the condition (A_{4.2}), and the Taylor expansion with Lagrange remainder, we obtain

$$f_1(t, x_1) \geq \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, x_2)}{i!} (x_1 - x_2)^i, \quad (4.1)$$

$$f_2(t, x_1) \geq \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, x_2)}{i!} (x_1 - x_2)^i + \frac{f_2^{(2k)}(t, x_1)}{(2k)!} (x_1 - x_2)^{2k} \quad (4.2)$$

for $(t, x_1), (t, x_2) \in \Omega, x_2 \leq x_1$. Similarly, we have

$$f_1(t, x_1) \leq \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, x_2)}{i!} (x_1 - x_2)^i, \quad (4.3)$$

$$f_2(t, x_1) \leq \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, x_2)}{i!} (x_1 - x_2)^i + \frac{f_2^{(2k)}(t, x_1)}{(2k)!} (x_1 - x_2)^{2k} \quad (4.4)$$

for $(t, x_1), (t, x_2) \in \Omega, x_1 \leq x_2$.

Consider the following nonlinear Caputo nabla fractional difference equations:

$$\begin{cases} \nabla_{a^*}^{\nu} y(t) = \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \alpha)}{i!} (y - \alpha)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \alpha)}{i!} (y - \alpha)^i + \frac{f_2^{(2k)}(t, \beta)}{(2k)!} (y - \alpha)^{2k} \\ \quad \equiv F(t, \alpha, \beta; y), \quad t \in \mathbb{N}_{a+1}^b, \\ y(a) = x_0, \end{cases} \quad (4.5)$$

and

$$\begin{cases} \nabla_{a^*}^v z(t) = \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \beta)}{i!} (z - \beta)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \beta)}{i!} (z - \beta)^i + \frac{f_2^{(2k)}(t, \alpha)}{(2k)!} (z - \beta)^{2k} \\ \equiv G(t, \alpha, \beta; z), \quad t \in \mathbb{N}_{a+1}^b, \\ z(a) = x_0. \end{cases} \quad (4.6)$$

We develop the sequences $\{\alpha_n(t)\}$, $\{\beta_n(t)\}$ using the above IVPs (4.5), (4.6), respectively. Letting $\alpha = \alpha_0$, $\beta = \beta_0$ in IVPs (4.5), (4.6). We first prove that $\alpha_0(t)$ and $\beta_0(t)$ are lower and upper solutions of the IVP (4.5) respectively. In fact, from the condition $(H_{3.3})$, we have

$$\begin{aligned} \nabla_{a^*}^v \alpha_0(t) &\leq f_1(t, \alpha_0) + f_2(t, \alpha_0) = F(t, \alpha_0, \beta_0; \alpha_0), \quad t \in \mathbb{N}_{a+1}^b, \\ \alpha_0(a) &\leq x_0, \end{aligned}$$

and by using the inequalities (4.1), (4.2), it follows that

$$\begin{aligned} \nabla_{a^*}^v \beta_0(t) &\geq \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \alpha_0)}{i!} (\beta_0 - \alpha_0)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \alpha_0)}{i!} (\beta_0 - \alpha_0)^i + \frac{f_2^{(2k)}(t, \beta_0)}{(2k)!} (\beta_0 - \alpha_0)^{2k} \\ &= F(t, \alpha_0, \beta_0; \beta_0), \quad t \in \mathbb{N}_{a+1}^b, \\ \beta_0(a) &\geq x_0, \end{aligned}$$

which imply that $\alpha_0(t)$ and $\beta_0(t)$ are lower and upper solutions of the IVP (4.5), respectively. Furthermore, we can see that $F(t, \alpha_0, \beta_0; y)$ is continuous with respect to y . Thus, by Lemma 3.4, there exists a solution $\alpha_1(t)$ of the IVP (4.5) such that $\alpha_0(t) \leq \alpha_1(t) \leq \beta_0(t)$ on \mathbb{N}_a^b .

Similarly, applying the fact that $\alpha_1(t)$ is a solution of the IVP (4.5), the inequalities (4.1)–(4.4), and the conditions $(H_{3.3})$, $(A_{4.2})$, we obtain

$$\begin{aligned} \nabla_{a^*}^v \alpha_1(t) &\stackrel{(A_{4.2})}{\leq} \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \alpha_0)}{i!} (\alpha_1 - \alpha_0)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \alpha_0)}{i!} (\alpha_1 - \alpha_0)^i + \frac{f_2^{(2k)}(t, \alpha_1)}{(2k)!} (\alpha_1 - \alpha_0)^{2k} \\ &\stackrel{(4.1), (4.2)}{\leq} f_1(t, \alpha_1) + f_2(t, \alpha_1) \\ &\stackrel{(4.3), (4.4)}{\leq} \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \beta_0)}{i!} (\alpha_1 - \beta_0)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \beta_0)}{i!} (\alpha_1 - \beta_0)^i + \frac{f_2^{(2k)}(t, \alpha_1)}{(2k)!} (\alpha_1 - \beta_0)^{2k} \\ &\stackrel{(A_{4.2})}{\leq} \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \beta_0)}{i!} (\alpha_1 - \beta_0)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \beta_0)}{i!} (\alpha_1 - \beta_0)^i + \frac{f_2^{(2k)}(t, \alpha_0)}{(2k)!} (\alpha_1 - \beta_0)^{2k} \\ &= G(t, \alpha_0, \beta_0; \alpha_1), \quad t \in \mathbb{N}_{a+1}^b, \\ \alpha_1(a) &= x_0, \end{aligned}$$

and

$$\begin{aligned} \nabla_{a^*}^v \beta_0(t) &\geq f_1(t, \beta_0) + f_2(t, \beta_0) = G(t, \alpha_0, \beta_0; \beta_0), \quad t \in \mathbb{N}_{a+1}^b, \\ \beta_0(a) &\geq x_0, \end{aligned}$$

which show that $\alpha_1(t)$ and $\beta_0(t)$ are lower and upper solutions of the IVP (4.6), respectively. Furthermore, we can find that $G(t, \alpha_0, \beta_0; z)$ is continuous with respect to z . Hence, in view of Lemma 3.4, we see that there exists a solution $\beta_1(t)$ of the IVP (4.6) such that $\alpha_1(t) \leq \beta_1(t) \leq \beta_0(t)$ on \mathbb{N}_a^b .

Next, we must show that $\alpha_1(t)$ and $\beta_1(t)$ are lower and upper solutions respectively of the IVP (3.1). For this purpose, using the conclusion that $\alpha_1(t)$ is a solution of the IVP (4.5), the condition (A_{4.2}), and the inequalities (4.1), (4.2), we have

$$\begin{aligned}\nabla_{a^*}^{\nu} \alpha_1(t) &\leq \sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \alpha_0)}{i!} (\alpha_1 - \alpha_0)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \alpha_0)}{i!} (\alpha_1 - \alpha_0)^i + \frac{f_2^{(2k)}(t, \alpha_1)}{(2k)!} (\alpha_1 - \alpha_0)^{2k} \\ &\leq f_1(t, \alpha_1) + f_2(t, \alpha_1), \quad t \in \mathbb{N}_{a+1}^b, \\ \alpha_1(a) &= x_0,\end{aligned}$$

which proves that $\alpha_1(t)$ is a lower solution of the IVP (3.1) on \mathbb{N}_a^b . Similar arguments show that

$$\begin{aligned}\nabla_{a^*}^{\nu} \beta_1(t) &\geq f_1(t, \beta_1) + f_2(t, \beta_1), \quad t \in \mathbb{N}_{a+1}^b, \\ \beta_1(a) &= x_0,\end{aligned}$$

which shows that $\beta_1(t)$ is an upper solution of the IVP (3.1) on \mathbb{N}_a^b . Therefore, we obtain

$$\alpha_0(t) \leq \alpha_1(t) \leq \beta_1(t) \leq \beta_0(t) \quad \text{on } \mathbb{N}_a^b.$$

By induction, we have

$$\alpha_0(t) \leq \alpha_1(t) \leq \cdots \leq \alpha_n(t) \leq \beta_n(t) \leq \cdots \leq \beta_1(t) \leq \beta_0(t) \quad \text{on } \mathbb{N}_a^b.$$

In addition, using the fact that $\alpha_n(t), \beta_n(t)$ are lower and upper solutions of the IVP (3.1) with $\alpha_n(t) \leq \beta_n(t)$, and the conditions of Lemma 3.4 are satisfied, we can conclude that there exists a solution $x_n(t)$ of the IVP (3.1) such that $\alpha_n(t) \leq x_n(t) \leq \beta_n(t)$ on \mathbb{N}_a^b . From this we can obtain that

$$\alpha_0(t) \leq \alpha_1(t) \leq \cdots \leq \alpha_n(t) \leq x_n(t) \leq \beta_n(t) \leq \cdots \leq \beta_1(t) \leq \beta_0(t) \quad \text{on } \mathbb{N}_a^b.$$

For any fixed $t \in \mathbb{N}_a^b$, the monotone sequences $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$ are uniformly bounded by $\{\alpha_0(t)\}$ and $\{\beta_0(t)\}$. Hence, we have that both monotone sequences $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$ are convergent on \mathbb{N}_a^b , that is, there exist functions $\rho, r : \mathbb{N}_a^b \rightarrow \mathbb{R}$ such that

$$\lim_{n \rightarrow \infty} \alpha_n(t) = \rho(t) \leq r(t) = \lim_{n \rightarrow \infty} \beta_n(t).$$

Taking $\alpha = \alpha_n, \beta = \beta_n$ in IVPs (4.5), (4.6), then we can show easily that $\rho(t), r(t)$ are solutions of the IVP (3.1). Next, we show that $\rho(t) \geq r(t)$. Let $p(t) := r(t) - \rho(t)$ so that $r(a) - \rho(a) = 0$. Using the condition (A_{4.1}), and the mean value theorem, we have

$$\begin{aligned}\nabla_{a^*}^{\nu} p(t) &= f_1(t, r) + f_2(t, r) - f_1(t, \rho) - f_2(t, \rho) \\ &= f_1^{(1)}(t, \xi_1)(r - \rho) + f_2^{(1)}(t, \xi_2)(r - \rho) \\ &\leq C_1 p + C_2 p \\ &= C p,\end{aligned}$$

where $\rho(t) \leq \xi_1(t), \xi_2(t) \leq r(t)$. By Lemma 3.5, we conclude that $r(t) \leq \rho(t)$ on \mathbb{N}_a^b . This proves that $\rho(t) = r(t)$. By the squeeze theorem, we have $\lim_{n \rightarrow \infty} x_n(t)$ exists, and

$$\lim_{n \rightarrow \infty} \alpha_n(t) = \rho(t) = \lim_{n \rightarrow \infty} x_n(t) = r(t) = \lim_{n \rightarrow \infty} \beta_n(t).$$

Set $\lim_{n \rightarrow \infty} x_n(t) = x(t)$, then we have $\rho(t) = x(t) = r(t)$. Hence $\alpha_n(t)$ and $\beta_n(t)$ converge uniformly and monotonically to a solution of the IVP (3.1).

Finally, we shall show that the convergence of the sequences $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$ to a solution $x(t)$ of the IVP (3.1) is of order $2k + 1$. For this purpose, set

$$\begin{aligned} p_n(t) &= x(t) - \alpha_n(t) \geq 0, \\ q_n(t) &= \beta_n(t) - x(t) \geq 0 \end{aligned}$$

for $t \in \mathbb{N}_a^b$ with $p_n(a) = q_n(a) = 0$.

From the conditions (A_{4.1}), (A_{4.2}), the Taylor's expansion with Lagrange remainder, and the mean value theorem, we obtain

$$\begin{aligned} & \nabla_{a^*}^v p_{n+1}(t) \\ &= f_1(t, x) + f_2(t, x) \\ & \quad - \left[\sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \alpha_n)}{i!} (\alpha_{n+1} - \alpha_n)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \alpha_n)}{i!} (\alpha_{n+1} - \alpha_n)^i + \frac{f_2^{(2k)}(t, \beta_n)}{(2k)!} (\alpha_{n+1} - \alpha_n)^{2k} \right] \\ &= f_1(t, x) + f_2(t, x) \\ & \quad - \left[f_1(t, \alpha_{n+1}) - \frac{f_1^{(2k)}(t, \xi_3)}{(2k)!} (\alpha_{n+1} - \alpha_n)^{2k} + \frac{f_1^{(2k)}(t, \alpha_n)}{(2k)!} (\alpha_{n+1} - \alpha_n)^{2k} + f_2(t, \alpha_{n+1}) \right. \\ & \quad \left. - \frac{f_2^{(2k)}(t, \xi_4)}{(2k)!} (\alpha_{n+1} - \alpha_n)^{2k} + \frac{f_2^{(2k)}(t, \beta_n)}{(2k)!} (\alpha_{n+1} - \alpha_n)^{2k} \right] \\ & \stackrel{(A_{4.2})}{\leq} f_1^{(1)}(t, \eta_1)(x - \alpha_{n+1}) + f_2^{(1)}(t, \eta_2)(x - \alpha_{n+1}) \\ & \quad + \frac{M_1}{(2k)!} (\alpha_{n+1} - \alpha_n)^{2k} (\xi_3 - \alpha_n) + \frac{M_2}{(2k)!} (\alpha_{n+1} - \alpha_n)^{2k} (\beta_n - \xi_4) \\ & \stackrel{(A_{4.1})}{\leq} C_1 p_{n+1} + C_2 p_{n+1} + K_1 p_n^{2k+1} + K_2 p_n^{2k} (p_n + q_n) \\ &= C p_{n+1} + (K_1 + K_2) p_n^{2k+1} + K_2 p_n^{2k} q_n \\ & \leq C p_{n+1} + K p_n^{2k} (p_n + q_n), \end{aligned}$$

where $\alpha_n \leq \xi_3, \xi_4 \leq \alpha_{n+1}$, $\alpha_{n+1} \leq \eta_1, \eta_2 \leq x$, $\frac{M_1}{(2k)!} = K_1$, $\frac{M_2}{(2k)!} = K_2$, $K = K_1 + K_2$. According to Lemma 3.5, we have $p_{n+1}(t) \leq x(t)$ on \mathbb{N}_a^b , where $x(t)$ is the solution of

$$\begin{cases} \nabla_{a^*}^v x(t) = Cx + K p_n^{2k} (p_n + q_n), & t \in \mathbb{N}_{a+1}^b, \\ x(a) = 0. \end{cases} \quad (4.7)$$

Hence, using the expression for $x(t)$ in Lemma 3.2, the solution $x(t)$ of (4.7) is given by

$$\begin{aligned} x(t) &= \sum_{s=a+1}^t H_{v-1}(t, \rho(s)) \left[Cx(s) + K p_n^{2k}(s)(p_n(s) + q_n(s)) \right] \\ & \leq \left[C \max_{s \in \mathbb{N}_{a+1}^t} x(s) + K \max_{s \in \mathbb{N}_{a+1}^t} p_n^{2k}(s)(p_n(s) + q_n(s)) \right] \sum_{s=a+1}^t H_{v-1}(t, \rho(s)) \\ &= \left[C \max_{s \in \mathbb{N}_{a+1}^t} x(s) + K \max_{s \in \mathbb{N}_{a+1}^t} p_n^{2k}(s)(p_n(s) + q_n(s)) \right] \int_a^t H_{v-1}(t, \rho(s)) \nabla s \end{aligned}$$

$$\begin{aligned}
&= \left[C \max_{s \in \mathbb{N}_{a+1}^b} x(s) + K \max_{s \in \mathbb{N}_{a+1}^b} p_n^{2k}(s)(p_n(s) + q_n(s)) \right] H_v(t, a) \\
&\leq H_v(b, a) \left[C \max_{s \in \mathbb{N}_{a+1}^b} x(s) + K \max_{s \in \mathbb{N}_{a+1}^b} p_n^{2k}(s)(p_n(s) + q_n(s)) \right].
\end{aligned}$$

Then, we have

$$\max_{s \in \mathbb{N}_{a+1}^b} x(s) \leq CH_v(b, a) \max_{s \in \mathbb{N}_{a+1}^b} x(s) + KH_v(b, a) \max_{s \in \mathbb{N}_{a+1}^b} p_n^{2k}(s)(p_n(s) + q_n(s)).$$

According to the condition $(H_{3.4})$, the above inequality can be written as

$$\max_{s \in \mathbb{N}_{a+1}^b} x(s) \leq (1 - CH_v(b, a))^{-1} KH_v(b, a) \max_{s \in \mathbb{N}_{a+1}^b} p_n^{2k}(s)(p_n(s) + q_n(s)).$$

Thus, we have

$$\max_{s \in \mathbb{N}_{a+1}^b} p_{n+1}(s) \leq (1 - CH_v(b, a))^{-1} KH_v(b, a) \max_{s \in \mathbb{N}_{a+1}^b} p_n^{2k}(s)(p_n(s) + q_n(s)). \quad (4.8)$$

Therefore, we conclude that the following inequality holds

$$\|p_{n+1}\| \leq L \|p_n\|^{2k} (\|p_n\| + \|q_n\|),$$

where $L = \frac{KH_v(b, a)}{1 - CH_v(b, a)}$ is a positive constant, which is the desired result.

Similarly, utilizing the conditions $(A_{4.1})$, $(A_{4.2})$, the Taylor's expansion with Lagrange remainder, and the mean value theorem, we have

$$\begin{aligned}
&\nabla_{a^*}^v q_{n+1}(t) \\
&= \left[\sum_{i=0}^{2k} \frac{f_1^{(i)}(t, \beta_n)}{i!} (\beta_{n+1} - \beta_n)^i + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \beta_n)}{i!} (\beta_{n+1} - \beta_n)^i + \frac{f_2^{(2k)}(t, \alpha_n)}{(2k)!} (\beta_{n+1} - \beta_n)^{2k} \right] \\
&\quad - f_1(t, x) - f_2(t, x) \\
&= \left[f_1(t, \beta_{n+1}) - \frac{f_1^{(2k)}(t, \xi_5)}{(2k)!} (\beta_{n+1} - \beta_n)^{2k} + \frac{f_1^{(2k)}(t, \beta_n)}{(2k)!} (\beta_{n+1} - \beta_n)^{2k} \right. \\
&\quad \left. + f_2(t, \beta_{n+1}) - \frac{f_2^{(2k)}(t, \xi_6)}{(2k)!} (\beta_{n+1} - \beta_n)^{2k} + \frac{f_2^{(2k)}(t, \alpha_n)}{(2k)!} (\beta_{n+1} - \beta_n)^{2k} \right] \\
&\quad - f_1(t, x) - f_2(t, x) \\
&\stackrel{(A_{4.2})}{\leq} f_1^{(1)}(t, \eta_3)(\beta_{n+1} - x) + f_2^{(1)}(t, \eta_4)(\beta_{n+1} - x) \\
&\quad + \frac{M_1}{(2k)!} (\beta_{n+1} - \beta_n)^{2k} (\beta_n - \xi_5) + \frac{M_2}{(2k)!} (\beta_{n+1} - \beta_n)^{2k} (\xi_6 - \alpha_n) \\
&\stackrel{(A_{4.1})}{\leq} C_1 q_{n+1} + C_2 q_{n+1} + K_1 q_n^{2k+1} + K_2 q_n^{2k} (q_n + p_n) \\
&= C q_{n+1} + (K_1 + K_2) q_n^{2k+1} + K_2 q_n^{2k} p_n \\
&\leq C q_{n+1} + K q_n^{2k} (q_n + p_n),
\end{aligned}$$

where $\beta_{n+1} \leq \xi_5, \xi_6 \leq \beta_n$, $x \leq \eta_3, \eta_4 \leq \beta_{n+1}$. By Lemma 3.5, we obtain $q_{n+1}(t) \leq x(t)$ on \mathbb{N}_a^b , where $x(t)$ is the solution of

$$\begin{cases} \nabla_{a^*}^v x(t) = Cx + Kq_n^{2k}(q_n + p_n), & t \in \mathbb{N}_{a+1}^b, \\ x(a) = 0. \end{cases} \quad (4.9)$$

Similar to the inequality (4.8) for $p_{n+1}(t)$, we can get

$$\max_{s \in \mathbb{N}_{a+1}^b} q_{n+1}(s) \leq (1 - CH_V(b, a))^{-1} KH_V(b, a) \max_{s \in \mathbb{N}_{a+1}^b} q_n^{2k}(s) (q_n(s) + p_n(s)). \quad (4.10)$$

Consequently, we get

$$\|q_{n+1}\| \leq L \|q_n\|^{2k} (\|q_n\| + \|p_n\|).$$

The proof is complete. \square

Theorem 4.2. Assume that the conditions $(H_{3,3})$ – $(H_{3,4})$ hold, and

(A_{4,3}) the functions $f_1, f_2 : \Omega \rightarrow \mathbb{R}$ are such that $f_1^{(i)}(t, x), f_2^{(i)}(t, x)$ ($i = 0, 1, \dots, 2k - 1$) exist, are continuous in its second variable, and for $C_1 > 0, C_2 > 0, C = C_1 + C_2$,

$$f_1^{(1)}(t, x) \leq C_1, \quad f_2^{(1)}(t, x) \leq C_2 \quad \text{on } \Omega.$$

(A_{4,4}) there exist $M_1, M_2 > 0$ such that for $x_1 \geq x_2, y_1 \geq y_2$ the functions $f_1^{(2k-1)}(t, x), f_2^{(2k-1)}(t, x)$ satisfy the following conditions:

$$\begin{aligned} 0 &\leq f_1^{(2k-1)}(t, x_1) - f_1^{(2k-1)}(t, x_2) \leq M_1(x_1 - x_2) && \text{on } \Omega, \\ 0 &\geq f_2^{(2k-1)}(t, y_1) - f_2^{(2k-1)}(t, y_2) \geq -M_2(y_1 - y_2) && \text{on } \Omega. \end{aligned}$$

Then there exist two sequences $\{\alpha_n(t)\}, \{\beta_n(t)\}, n \geq 0$ which converge uniformly and monotonically to a solution of the IVP (3.1) and the convergence is of order $2k$.

Proof. In order to construct monotone sequences $\{\alpha_n(t)\}, \{\beta_n(t)\}, n \geq 0$ which converge uniformly and monotonically to the solution of the IVP (3.1), we need to consider the following nonlinear Caputo nabla fractional difference equations:

$$\left\{ \begin{aligned} \nabla_{a^*}^v \alpha_n(t) &= \sum_{i=0}^{2k-1} \frac{f_1^{(i)}(t, \alpha_{n-1})}{i!} (\alpha_n - \alpha_{n-1})^i + \sum_{i=0}^{2k-2} \frac{f_2^{(i)}(t, \alpha_{n-1})}{i!} (\alpha_n - \alpha_{n-1})^i \\ &\quad + \frac{f_2^{(2k-1)}(t, \beta_{n-1})}{(2k-1)!} (\alpha_n - \alpha_{n-1})^{2k-1} \\ &\equiv F(t, \alpha_{n-1}, \beta_{n-1}; \alpha_n), \quad t \in \mathbb{N}_{a+1}^b, \\ \alpha_n(a) &= x_0, \end{aligned} \right. \quad (4.11)$$

and

$$\left\{ \begin{aligned} \nabla_{a^*}^v \beta_n(t) &= \sum_{i=0}^{2k-2} \frac{f_1^{(i)}(t, \beta_{n-1})}{i!} (\beta_n - \beta_{n-1})^i + \frac{f_1^{(2k-1)}(t, \alpha_{n-1})}{(2k-1)!} (\beta_n - \beta_{n-1})^{2k-1} \\ &\quad + \sum_{i=0}^{2k-1} \frac{f_2^{(i)}(t, \beta_{n-1})}{i!} (\beta_n - \beta_{n-1})^i \\ &\equiv G(t, \alpha_{n-1}, \beta_{n-1}; \beta_n), \quad t \in \mathbb{N}_{a+1}^b, \\ \beta_n(a) &= x_0. \end{aligned} \right. \quad (4.12)$$

Similar to Theorem 4.1, we can show that the convergence is of order $2k$. \square

5 Numerical result

Now, we give a numerical example to illustrate the application of the results established in the previous section.

Example 5.1. Consider the following nonlinear Caputo nabla fractional difference equation

$$\begin{cases} \nabla_{0^*}^{\frac{1}{2}} x(t) = -x^5(t) + x^3(t) + x^2(t) - 5x(t) - 1, & t \in \mathbb{N}_1^5, \\ x(0) = 0. \end{cases} \quad (5.1)$$

Taking $\alpha_0(t) = -1$, $\beta_0(t) = 1$, it is easy to verify that $\alpha_0(t)$, $\beta_0(t)$ are lower and upper solutions of the IVP (5.1), respectively. According to the Definitions 2.2, 2.4, we have $\nabla_{0^*}^{\frac{1}{2}} x(t) = \nabla_0^{-\frac{1}{2}} \nabla x(t) = \sum_{s=1}^t \frac{\Gamma(t-s+\frac{1}{2})}{\Gamma(\frac{1}{2})\Gamma(t-s+1)} \nabla x(s)$. Let $f(t, x)$ denote the right-hand side of (5.1), and split it into two functions as $f(t, x) = f_1(t, x) + f_2(t, x)$, where $f_1(t, x) = x^3(t) + x^2(t) - 5x(t) - 1$, $f_2(t, x) = -x^5(t)$.

Table 5.1: Table of four α -iterates of (5.1).

t	$\alpha_1(t)$	$\alpha_2(t)$	$\alpha_3(t)$	$\alpha_4(t)$	$x(t)$
1	-0.595929	-0.227396	-0.163045	-0.162943	-0.162943
2	-0.611197	-0.248267	-0.176153	-0.175965	-0.175965
3	-0.615457	-0.254921	-0.180447	-0.180254	-0.180254
4	-0.617607	-0.258339	-0.182687	-0.182480	-0.182480
5	-0.618955	-0.260491	-0.184105	-0.183888	-0.183888

Table 5.2: Table of four β -iterates of (5.1).

t	$\beta_4(t)$	$\beta_3(t)$	$\beta_2(t)$	$\beta_1(t)$
1	-0.162943	-0.161992	-0.011633	0.525454
2	-0.175965	-0.174333	-0.000297	0.539207
3	-0.180254	-0.178300	0.003561	0.543062
4	-0.182480	-0.180338	0.005570	0.545007
5	-0.183888	-0.181619	0.006843	0.546226

It is easy to show that

$$\begin{aligned} f_1^{(1)}(t, x) &= 3x^2(t) + 2x(t) - 5 = 3 \left(x(t) + \frac{1}{3} \right)^2 - \frac{16}{3} \leq 0 \quad \text{on } \Omega, \\ f_2^{(1)}(t, x) &= -5x^4(t) \leq 0 \quad \text{on } \Omega, \end{aligned}$$

so we can choose $C = \frac{1}{3}$ such that

$$CH_{\frac{1}{2}}(5, 0) = \frac{1}{3} \cdot \frac{315}{128} = \frac{105}{128} < 1.$$




Figure 5.1: α -iterates (broken lines), β -iterates (unbroken lines).

Furthermore, we can check that for $x_1(t) \geq x_2(t)$, $y_1(t) \geq y_2(t)$,

$$0 \leq f_1^{(2)}(t, x_1) - f_1^{(2)}(t, x_2) = 6(x_1(t) - x_2(t)) \quad \text{on } \Omega,$$

$$0 \geq f_2^{(2)}(t, y_1) - f_2^{(2)}(t, y_2) = -20y_1^3(t) + 20y_2^3(t) \geq -60(y_1(t) - y_2(t)) \quad \text{on } \Omega.$$

Hence, we can apply the iterates of Theorem 4.1. After only four iterates of α and β , we can find the α , β -iterates as given in Tables 5.1, 5.2. Figure 5.1 shows the graphs of some α -iterates (broken lines) and some β -iterates (unbroken lines). Note that the actual graphs of the $\alpha_i(t)$ and the $\beta_i(t)$, $1 \leq i \leq 3$, $1 \leq t \leq 5$, consist just of the points corresponding to the values of these functions at $t = 1, 2, 3, 4, 5$. Note that for all practical purposes the graphs of $\alpha_3(t)$, $\beta_3(t)$ in Figure 5.1 and the solution $x(t)$ are the same to several decimal places.

6 Conclusion

In the above parts, we proved the existence of solutions for nonlinear Caputo nabla fractional difference equations. Based on this fact, we have developed two monotone sequences which converge rapidly to the solution of such equations. In addition, a numerical example is given to show one of the established results.

Acknowledgements

This paper is supported by the National Natural Science Foundation of China (11271380) and Guangdong Province Key Laboratory of Computational Science.

References

- [1] T. ABDELJAWAD, On Riemann and Caputo fractional differences, *Comput. Math. Appl.* **62**(2011), 1602–1611. [MR2824747](#); [url](#)
- [2] R. P. AGARWAL, *Difference equations and inequalities*, Marcel Dekker, New York, NY, USA, 1992. [MR1155840](#)

- [3] B. AHMAD, R. A. KHAN, S. SIVASUNDARAM, Generalized quasi-linearization method for nonlinear functional differential equations, *J. Appl. Math. Stoch. Anal.* **16**(2003), 33–43. [MR1973077](#); [url](#)
- [4] B. AHMAD, A. ALSAEDI, An extended method of quasi-linearization for nonlinear impulsive differential equations with a nonlinear three-point boundary condition, *Electron. J. Qual. Theory Differ. Equ.* **2007**, No. 1, 1–19. [MR2276782](#); [url](#)
- [5] P. AMSTER, P. DE NAPOLI, A quasi-linearization method for elliptic problems with a nonlinear boundary condition, *Nonlinear Anal.* **10**(2007), No. 10, 2255–2263. [MR2311029](#); [url](#)
- [6] F. M. ATICI, F. WU, Existence of solutions for nonlinear fractional difference equations with initial conditions, *Dynam. Systems Appl.* **23**(2014), No. 2–3, 265–276. [MR3241878](#); [url](#)
- [7] D. D. BAINOV, S. G. HRISTOVA, The method of quasi-linearization for the periodic boundary value problem for systems of impulsive differential equations, *Appl. Math. Comput.* **117**(2001), 73–85. [MR1801404](#); [url](#)
- [8] R. BELLMAN, R. KALABA, *Quasi-linearization and nonlinear boundary value problems*, Elsevier, New York, 1965. [MR0178571](#)
- [9] A. BUICĂ, Quasi-linearization method for nonlinear elliptic boundary value problems, *J. Optimiz. Theory. Appl.* **124**(2005), No. 2, 323–338. [MR2130073](#); [url](#)
- [10] J. V. DEVI, F. A. McRAE, Z. DRICI, Generalized quasi-linearization for fractional differential equations, *Comput. Math. Appl.* **59**(2010), 1057–1062. [MR2579470](#); [url](#)
- [11] Z. DRICI, F. A. McRAE, J. V. DEVI, Quasi-linearization for functional differential equations with retardation and anticipation, *Nonlinear Anal.* **70**(2009), 1763–1775. [MR2483597](#); [url](#)
- [12] M. EL-GEBEILY, D. O'REGAN, Existence and generalized quasi-linearization methods for singular nonlinear differential equations, *Dynam. Systems Appl.* **21**(2008), 67–81. [MR2953238](#)
- [13] C. S. GOODRICH, Existence and uniqueness of solutions to a fractional difference equation with nonlocal conditions, *Comput. Math. Appl.* **61**(2009), 191–202. [MR2754129](#); [url](#)
- [14] C. S. GOODRICH, Some new existence results for fractional difference equations, *Int. J. Dyn. Syst. Differ. Equ.* **3**(2014), 145–162. [MR2797046](#); [url](#)
- [15] C. S. GOODRICH, A. C. PETERSON, *Discrete fractional calculus*, Springer, New York, 2015. [MR3445243](#); [url](#)
- [16] J. JONNALAGADDA, Solutions of perturbed linear nabla fractional difference equations, *Differ. Equ. Dyn. Syst.* **22**(2014), No. 3, 281–292. [MR3218901](#); [url](#)
- [17] W. G. KELLEY, A. C. PETERSON, *Difference equations: an introduction with applications*, 2nd ed., Academic Press, New York, 2001. [MR1765695](#)
- [18] V. LAKSHMIKANTHAM, A. S. VATSALA, *Generalized quasi-linearization for nonlinear problems*, Kluwer Academic Publishers, Dordrecht, 1998. [MR1640601](#); [url](#)
- [19] V. LAKSHMIKANTHAM, S. KOKSAL, *Monotone flows and rapid convergence for nonlinear partial differential equations*, Taylor & Francis, London and New York, 2003. [MR2014745](#)

- [20] V. LAKSMIKANTHAM, S. LEELA, ZAHIA DRICI, F. A. McRAE, *Theory of causal differential equations*, World Scientific, Atlantis Press, 2009. [MR2596883](#); [url](#)
- [21] V. A. VIJESH, A short note on the quasi-linearization method for fractional differential equations, *Numer. Funct. Anal. Optim.* **37**(2016), No. 9, 1158–1167. [MR3546127](#); [url](#)
- [22] P. G. WANG, H. X. WU, Y. H. WU, Higher even-order convergence and coupled solutions for second-order boundary value problems on time scales, *Comput. Math. Appl.* **55**(2008), 1693–1705. [MR2405174](#); [url](#)
- [23] P. G. WANG, P. LI, k th Order convergence for a semilinear elliptic boundary value problem in the divergence form, *Appl. Math. Comput.* **217**(2011), 8547–8551. [MR2802266](#); [url](#)
- [24] P. G. WANG, W. GAO, Quasi-linearization of an initial value problem for a set valued integro-differential equation, *Comput. Math. Appl.* **61**(2011), 2111–2115. [MR2785565](#); [url](#)
- [25] P. G. WANG, Y. HOU, Generalized quasi-linearization for the system of fractional differential equations, *J. Funct. Spaces Appl.* **2013**, Art. ID 793263, 5 pp. [MR3053278](#); [url](#)
- [26] A. YAKAR, Initial time difference quasi-linearization for Caputo fractional differential equations, *Adv. Difference Equ.* **2012**, 2012:92, 9 pp. [MR2957672](#); [url](#)