

On a particular second-order nonlinear differential subordination I

Georgia Irina Oros

Abstract

We find conditions on the complex-valued functions A, B, C, D in the unit disc U such that the differential inequality

$$|A(z)z^2p'(z) + B(z)zp'(z) + C(z)p^2(z) + D(z)p(z)| < M$$

implies $|p(z)| < N$, where p is analytic in U , with $p(0) = 0$.

2000 Mathematical Subject Classification: 30C80

1 Introduction and preliminaries

We let $\mathcal{H}[U]$ denote the class of holomorphic functions in the unit disc

$$U = \{z \in \mathbb{C} : |z| < 1\}.$$

For $a \in \mathbb{C}$ and $n \in \mathbb{N}^*$ we let

$$\mathcal{H}[a, n] = \{f \in \mathcal{H}[U], f(z) = a + a_n z^n + a_{n+1} z^{n+1} + \dots, z \in U\}$$

and

$$\mathcal{A}_n = \{f \in \mathcal{H}[U], f(z) = z + a_{n+1}z^{n+1} + a_{n+2}z^{n+2} + \dots, z \in U\}$$

with $\mathcal{A}_1 = \mathcal{A}$.

In [1] chapter IV, the authors have analyzed a second-order linear differential subordination

$$(1) \quad A(z)z^2p'(z) + B(z)zp'(z) + C(z)p(z) + D(z) < h(z),$$

where A, B, C, D and h are complex-valued functions in the unit disc, where $p \in \mathcal{H}[0, n]$. A more general version of (1) is given by:

$$A(z)z^2p'(z) + B(z)zp'(z) + C(z)p(z) + D(z) \in \Omega,$$

where $\Omega \subset \mathbb{C}$.

In [2] we found conditions on the complex-valued functions A, B, C, D in the unit disc U and the positive numbers M and N such that

$$|A(z)zp'(z) + B(z)p^2(z) + C(z)p(z)| < M$$

implies $|p(z)| < N$, where $p \in \mathcal{H}[0, n]$.

In this paper we shall consider the following particular second-order nonlinear differential subordination given by the inequality

$$(2) \quad |A(z)z^2p'(z) + B(z)zp'(z) + C(z)p^2(z) + D(z)p(z)| < M,$$

where $p \in \mathcal{H}[0, n]$.

We find conditions on complex-valued functions A, B, C, D and the positive numbers M and N such that (2) implies

$$|p(z)| < N,$$

where $p \in \mathcal{H}[0, n]$.

In order to prove the new results we shall use the following lemma:

Lemma A. [1, p. 34] *Let $\psi : \mathbb{C}^3 \times U \rightarrow \mathbb{C}$ and $M > 0, N > 0, n$ positive integer, satisfy*

$$(3) \quad |\psi(Ne^{i\theta}, Ke^{i\theta}, L; z)| \geq M$$

whenever

$$\operatorname{Re} [Le^{-i\theta}] \geq (n - 1)K, \quad K \geq nN,$$

$z \in U$ and $\theta \in \mathbb{R}$.

If $p \in \mathcal{H}[0, n]$ and

$$|\psi(p(z), zp'(z), z^2p'(z); z)| < M$$

then $|p(z)| < N$.

2 Main results

Theorem. *Let $M > 0, N > 0$, and let n be a positive integer. Suppose that the functions $A, B, C, D : U \rightarrow \mathbb{C}$ satisfy $A(z) \neq 0$,*

$$(i) \operatorname{Re} \frac{B(z)}{A(z)} \geq -n$$

$$(ii) \operatorname{Re} \frac{B(z) + D(z)}{A(z)} \geq \frac{M + N^2|C(z)|}{N|A(z)|}.$$

If $p \in \mathcal{H}[0, n]$ and

$$|A(z)z^2p'(z) + B(z)zp'(z) + C(z)p^2(z) + D(z)p(z)| < M$$

then

$$|p(z)| < N.$$

Proof. Let $\psi : \mathbb{C}^3 \times U \rightarrow \mathbb{C}$ be defined by

$$(4) \quad \begin{aligned} \psi(p(z), zp'(z), z^2p'(z); z) &= A(z)z^2p'(z) + B(z)zp'(z) + \\ &+ C(z)p^2(z) + D(z)p(z). \end{aligned}$$

From (2) we have

$$(5) \quad |\psi(p(z), zp'(z), z^2p'(z); z)| < M, \text{ for } z \in U.$$

Using (ii) in (4) we have

$$\begin{aligned} |\psi(Ne^{i\theta}, Ke^{i\theta}, L; z)| &= |A(z)L + B(z) + Ke^{i\theta} + C(z)N^2e^{2i\theta} + D(z)Ne^{i\theta}| = \\ &= |A(z)Le^{-i\theta} + B(z)K + C(z)N^2e^{i\theta} + D(z)N| = \\ &= |A(z)| \left| Le^{-i\theta} + \frac{B(z)}{A(z)}K + \frac{C(z)}{A(z)}N^2e^{i\theta} + \frac{D(z)}{A(z)}N \right| \geq \\ &\geq |A(z)| \left[\left| Le^{-i\theta} + K\frac{B(z)}{A(z)} + N\frac{D(z)}{A(z)} \right| - N^2 \left| \frac{C(z)}{A(z)} \right| \right] \geq \\ &\geq |A(z)| \left[\operatorname{Re} Le^{-i\theta} + K\operatorname{Re} \frac{B(z)}{A(z)} + N\operatorname{Re} \frac{D(z)}{A(z)} - N^2 \left| \frac{C(z)}{A(z)} \right| \right] \geq \end{aligned}$$

$$\begin{aligned}
 &\geq |A(z)| \left[(n-1)K + K \operatorname{Re} \frac{B(z)}{A(z)} + N \operatorname{Re} \frac{D(z)}{A(z)} - N^2 \left| \frac{C(z)}{A(z)} \right| \right] \geq \\
 &\geq |A(z)| \left[n(n-1)N + nN \operatorname{Re} \frac{B(z)}{A(z)} + N \operatorname{Re} \frac{D(z)}{A(z)} - N^2 \left| \frac{C(z)}{A(z)} \right| \right] \geq \\
 &\geq |A(z)| \left[nN \operatorname{Re} \frac{B(z)}{A(z)} + N \operatorname{Re} \frac{D(z)}{A(z)} - N^2 \left| \frac{C(z)}{A(z)} \right| \right] \geq \\
 &\geq |A(z)| \left[N \operatorname{Re} \frac{B(z)}{A(z)} + N \operatorname{Re} \frac{D(z)}{A(z)} - N^2 \left| \frac{C(z)}{A(z)} \right| \right] \geq \\
 &\geq |A(z)| \left[N \operatorname{Re} \frac{B(z) + D(z)}{A(z)} - N^2 \left| \frac{C(z)}{A(z)} \right| \right] \geq M.
 \end{aligned}$$

Hence condition (3) holds and by Lemma A we deduce that (5) implies $|p(z)| < N$.

Instead of prescribing the constant N in Theorem, in some cases we can use (ii) to determine an appropriate $N = N(M, n, A, B, C, D)$ so that (2) implies $|p(z)| < N$. This can be accomplished by solving (ii) for N by taking the supremum of the resulting function over U .

Condition (ii) is equivalent to:

$$(6) \quad N^2|C(z)| - N|A(z)| \operatorname{Re} \frac{B(z) + D(z)}{A(z)} + M \leq 0.$$

If we suppose $C(z) \neq 0$, then the inequality (6) holds if

$$(7) \quad |A(z)| \operatorname{Re} \frac{B(z) + D(z)}{A(z)} \geq 2\sqrt{|C(z)|M}$$

If (7) holds, the roots of the trinomial in (6) are

$$N_{1,2} = \frac{|A(z)| \operatorname{Re} \frac{B(z) + D(z)}{A(z)} \pm \sqrt{\left[|A(z)| \operatorname{Re} \frac{B(z) + D(z)}{A(z)} \right]^2 - 4M|C(z)|}}{2|C(z)|}.$$

We let

$$N = \frac{2M}{|A(z)| \operatorname{Re} \frac{B(z) + D(z)}{A(z)} + \sqrt{\left[|A(z)| \operatorname{Re} \frac{B(z) + D(z)}{A(z)} \right]^2 - 4M|C(z)|}}.$$

If this supremum is finite, the Theorem can be rewritten as follows:

Corollary 1. *Let $M > 0$ and let n be a positive integer. Suppose that $p \in \mathcal{H}[0, n]$ and that the functions $A, B, C, D : U \rightarrow \mathbb{C}$, with $A(z) \neq 0$, satisfy*

$$(i) \operatorname{Re} \frac{B(z)}{A(z)} \geq -n$$

$$(ii) N = \sup_{|z| < 1} \frac{2M}{|A(z)| \operatorname{Re} \frac{B(z)+D(z)}{A(z)} + \sqrt{\left[|A(z)| \operatorname{Re} \frac{B(z)+D(z)}{A(z)} \right]^2 - 4M|C(z)|}} < \infty$$

then

$$|A(z)z^2p'(z) + B(z)zp'(z) + C(z)p^2(z) + D(z)p(z)| < M$$

implies

$$|p(z)| < N.$$

Let $n = 1$, $A(z) = 3$, $B(z) = 2 + 4z$, $C(z) = 2$, $D(z) = 10 - 4z$, $M = 8$,

$$N = \frac{4}{3 + \sqrt{5}}.$$

In this case from Corollary 1 we deduce:

Example 1. If $p \in \mathcal{H}[0, 1]$, then

$$|3z^2p'(z) + (2 + 4z)zp'(z) + 2p^2(z) + (10 - 4z)p(z)| < 8$$

implies

$$|p(z)| < \frac{4}{3 + \sqrt{5}}.$$

If $n = 2$, $A(z) = 6$, $B(z) = 8 - 2z$, $C(z) = -4$, $D(z) = 4 + 2z$, $M = 5$,
 $N = \frac{1}{2}$.

Let this case from Corollary 1 we deduce

Example 2. If $p \in \mathcal{H}[0, 2]$ then

$$|6z^2p'(z) + (8 - 2z)zp'(z) - 4p^2(z) + (4 + 2z)p(z)| < 5$$

implies

$$|p(z)| < \frac{1}{2}.$$

If $A(z) = A > 0$ then the Theorem can be rewritten as follows:

Corollary 2. Let $M > 0$, $N > 0$ and let n be a positive integer. Suppose that the functions $B, C, D : U \rightarrow \mathbb{C}$ satisfy

- (i) $\operatorname{Re} B(z) \geq -nA$, $A > 0$,
- (ii) $\operatorname{Re} [B(z) + D(z)] \geq \frac{M + N^2|C(z)|}{N}$.

If $p \in \mathcal{H}[0, n]$ and

$$|Az^2p'(z) + B(z)zp'(z) + C(z)p^2(z) + D(z)p(z)| < M$$

then

$$|p(z)| < N.$$

References

- [1] S. S. Miller and P. T. Mocanu, *Differential Subordinations. Theory and Applications*, Marcel Dekker Inc., New York, Basel, 2000.
- [2] Gh. Oros and Georgia Irina Oros, *On a particular first-order nonlinear differential subordination I* (submitted).

Faculty of Mathematics and Computer Sciences

Babeş-Bolyai University

3400 Cluj-Napoca, Romania