

ON SOME PROPERTIES OF SOLUTIONS OF SECOND  
ORDER LINEAR FUNCTIONAL DIFFERENTIAL  
EQUATIONS

I. KIGURADZE

ABSTRACT. The properties of solutions of the equation  $u''(t) = p_1(t)u(\tau_1(t)) + p_2(t)u'(\tau_2(t))$  are investigated where  $p_i : [a, +\infty[ \rightarrow R$  ( $i = 1, 2$ ) are locally summable functions,  $\tau_1 : [a, +\infty[ \rightarrow R$  is a measurable function and  $\tau_2 : [a, +\infty[ \rightarrow R$  is a nondecreasing locally absolutely continuous one. Moreover,  $\tau_i(t) \geq t$  ( $i = 1, 2$ ),  $p_1(t) \geq 0$ ,  $p_2^2(t) \leq (4 - \varepsilon)\tau_2'(t)p_1(t)$ ,  $\varepsilon = \text{const} > 0$  and  $\int_a^{+\infty} (\tau_1(t) - t)p_1(t)dt < +\infty$ . In particular, it is proved that solutions whose derivatives are square integrable on  $[a, +\infty[$  form a one-dimensional linear space and for any such solution to vanish at infinity it is necessary and sufficient that  $\int_a^{+\infty} tp_1(t)dt = +\infty$ .

Consider the differential equation

$$u''(t) = p_1(t)u(\tau_1(t)) + p_2(t)u'(\tau_2(t)), \quad (1)$$

where  $p_i : [a, +\infty[ \rightarrow R$  ( $i = 1, 2$ ) are locally summable functions,  $\tau_i : [a, +\infty[ \rightarrow R$  ( $i = 1, 2$ ) are measurable functions and

$$\tau_i(t) \geq t \quad \text{for } t \geq a \quad (i = 1, 2). \quad (2)$$

We say that a solution  $u$  of the equation (1) is a *Kneser-type* solution if it satisfies the inequality  $u'(t)u(t) \leq 0$  for  $t \geq a_0$  for some  $a_0 \in [a, +\infty[$ . A set of such solutions is denoted by  $K$ . By  $W$  we denote a space of solutions of (1) that satisfy  $\int_a^{+\infty} u'^2(t)dt < +\infty$ . The results of [1, 2] imply that if  $p_1(t) \geq 0$  for  $t \geq a$  and the condition

$$(i) \quad \tau_i(t) \equiv t, \quad (i = 1, 2), \quad \int_a^{+\infty} |p_2(t)|dt < +\infty,$$

or

$$(ii) \quad p_2(t) \leq 0, \quad \text{for } t \geq 0, \quad \int_a^{+\infty} s p_1(s) ds < +\infty, \quad \int_a^{+\infty} \frac{s}{\tau_2(s)} |p_2(s)| ds < +\infty,$$

holds, then  $W \supset K$  and  $K$  is a one-dimensional linear space. The case when the conditions (i) and (ii) are violated, the matter of dimension of  $K$  and  $W$  and their interconnection has actually remained unstudied. An attempt is made in this note to fill up this gap to a certain extent.

**Theorem 1.** *Let  $\tau_i(t) \geq t$  ( $i = 1, 2$ ),  $p_1(t) \geq 0$  for  $t \geq a$ ,*

$$\int_a^{+\infty} [\tau_1(t) - t] p_1(t) dt < +\infty, \quad (3)$$

and let  $\tau_2$  be a nondecreasing locally absolutely continuous function satisfying

$$p_2^2(t) \leq (4 - \varepsilon) \tau_2'(t) p_1(t) \quad \text{for } t \geq a, \quad (4)$$

where  $\varepsilon = \text{const} > 0$ . Then

$$W \subset K, \quad \dim W = 1. \quad (5)$$

Before proceeding to the proof of the theorem we shall give two auxiliary statements.

**Lemma 1.** *Let the conditions of Theorem 1 be fulfilled and let  $a_0 \in [a, +\infty[$  be large enough for the equality*

$$\int_{a_0}^{+\infty} [\tau_1(s) - s] p_1(s) ds \leq 4\delta^2, \quad (6)$$

where  $\delta = \frac{1}{4}[2 - (4 - \varepsilon)^{1/2}]$ , to hold. Then any solution  $u$  of the equation (1) satisfies

$$\begin{aligned} \delta \int_t^x [u'^2(s) + p_1(s) u^2(s)] ds &\leq u'(x)u(x) - u'(t)u(t) + \\ &+ (1 - \delta) \int_x^{\tau(x)} u'^2(s) ds \quad \text{for } a_0 \leq t \leq x < +\infty, \end{aligned} \quad (7)$$

where  $\tau(x) = \text{ess sup}_{a_0 \leq t \leq x} [\max_{1 \leq i \leq 2} \tau_i(x)]$ . Moreover, if  $u \in W$ , then

$$u'(t)u(t) \leq -\delta \int_t^{+\infty} [u'^2(s) + p_1(s)u^2(s)]ds \quad \text{for } t \geq a_0 \tag{8}$$

and

$$2\delta \int_t^{+\infty} (s-t)[u'^2(s) + p_1(s)u^2(s)]ds \leq u^2(t) \quad \text{for } t \geq a_0. \tag{9}$$

*Proof.* Let  $u$  be any solution of the equation (1). Then

$$-u''(t)u(t) + p_1(t)u^2(t) = p_1(t)u(t) \int_{\tau_1(t)}^t u'(s)ds - p_2(t)u'(\tau_2(t))u(t).$$

Integrating this equality from  $t$  to  $x$ , we obtain

$$\begin{aligned} & u'(t)u(t) - u'(x)u(x) + \int_t^x [u'^2(s) + p_1(s)u^2(s)]ds = \\ & = \int_t^x [p_1(s)u(s) \int_{\tau_1(s)}^s u'(y)dy]ds - \int_t^x p_2(s)u'(\tau_2(s))u(s)ds. \end{aligned}$$

However, in view of (4) and (6),

$$\begin{aligned} & \int_t^x [p_1(s)u(s) \int_{\tau_1(s)}^s u'(y)dy]ds \leq \delta \int_t^x p_1(s)u^2(s)ds + \\ & + \frac{1}{4\delta} \left[ \int_t^x [\tau_1(s) - s]p_1(s)ds \right] \left[ \int_t^{\tau(x)} u'^2(s)ds \right] \leq \\ & \leq \delta \int_t^x p_1(s)u^2(s)ds + \delta \int_t^{\tau(x)} u'^2(s)ds \quad \text{for } a_0 \leq t \leq x < +\infty \end{aligned}$$

and

$$-\int_t^x p_2(s)u'(\tau_2(s))u(s)ds \leq$$

$$\begin{aligned}
&\leq 2(1-2\delta) \int_t^x \left[ p_1(s)u^2(s) \right]^{1/2} \left[ \tau_2'(s)u'^2(\tau_2(s)) \right]^{1/2} ds \leq \\
&\leq (1-2\delta) \int_t^x p_1(s)u^2(s)ds + (1-2\delta) \int_t^x \tau_2'(s)u'^2(\tau_2(s))ds \leq \\
&\leq (1-2\delta) \int_t^x p_1(s)u^2(s)ds + (1-2\delta) \int_t^{\tau(x)} u'^2(s)ds \\
&\quad \text{for } a_0 \leq t \leq x < +\infty.
\end{aligned}$$

Therefore

$$\begin{aligned}
&u'(t)u(t) - u'(x)u(x) + \int_t^x [u'^2(s) + p_1(s)u^2(s)]ds \leq \\
&\leq (1-\delta) \int_t^x [u'^2(s) + p_1(s)u^2(s)]ds + (1-\delta) \int_x^{\tau(x)} u'^2(s)ds \\
&\quad \text{for } a_0 \leq t \leq x < +\infty
\end{aligned}$$

and thus the inequality (7) holds.

Suppose now that  $u \in W$ . Then, as one can easily verify,

$$\liminf_{x \rightarrow +\infty} |u'(x)u(x)| = 0.$$

So (7) immediately implies (8). Integrating both sides of (8) from  $t$  to  $+\infty$ , we obtain the estimate (9).  $\square$

**Lemma 2.** *Let the conditions of Lemma 1 be fulfilled and there exist  $b \in ]a_0, +\infty[$  such that*

$$p_i(t) = 0 \quad \text{for } t \geq b \quad (i = 1, 2). \quad (10)$$

*Then for any  $c \in R$  there exists a unique solution of the equation (1) satisfying*

$$u(a_0) = c, \quad u'(t) = 0 \quad \text{for } t \geq b. \quad (11)$$

*Proof.* In view of (2) and (10), for any  $\alpha \in R$  the equation (1) has a unique solution  $v(\cdot; \alpha)$  satisfying  $v(t; \alpha) = \alpha$  for  $b \leq t < +\infty$ . Moreover,  $v(t; \alpha) = \alpha v(t; 1)$ . On the other hand, by Lemma 1 the function  $v(\cdot; 1) : [a_0, +\infty[ \rightarrow R$  is non increasing and  $v(a_0; 1) \geq 1$ . Therefore the function  $u(\cdot) = \frac{c}{v(a_0; 1)} v(a_0; \cdot)$  is a unique solution of (1), (11).  $\square$

*Proof of Theorem 1.* First of all we shall prove that for any  $c \in R$  the equation (1) has at least one solution satisfying

$$u(a_0) = c, \quad \int_{a_0}^{+\infty} u'^2(s)ds < +\infty. \tag{12}$$

For any natural  $k$  put

$$p_{ik}(t) = \begin{cases} p_i(t) & \text{for } a_0 \leq t \leq a_0 + k \\ 0 & \text{for } t > a_0 + k \end{cases} \quad (i = 1, 2). \tag{13}$$

According to Lemma 2, for any  $k$  the equation  $u''(t) = p_{1k}(t)u(\tau_1(t)) + p_{2k}(t)u'(\tau_2(t))$  has a unique solution  $u_k$  satisfying

$$u_k(a_0) = c, \quad u'_k(t) = 0 \quad \text{for } t \geq a_0 + k. \tag{14}$$

On the other hand, by Lemma 1

$$|u_k(t)| \leq |c| \quad \text{for } t \geq a_0, \quad 2\delta \int_{a_0}^{+\infty} (s - a_0)u'^2_k(s)ds \leq c^2. \tag{15}$$

Taking (2) and (13)–(15) into account, it is easy to show that the sequences  $(u_k)_{k=1}^{+\infty}$  and  $(u'_k)_{k=1}^{+\infty}$  are uniformly bounded and equicontinuous on each closed subinterval of  $[a_0, +\infty[$ . Therefore, by the Arzela-Ascoli lemma, we can choose a subsequence  $(u_{k_m})_{m=1}^{+\infty}$  out of  $(u_k)_{k=1}^{+\infty}$ , which is uniformly convergent alongside with  $(u'_{k_m})_{m=1}^{+\infty}$  on each closed subinterval of  $[a, +\infty[$ . By (13)–(15) the function  $u(t) = \lim_{m \rightarrow +\infty} u_{k_m}(t)$  for  $t \geq a$  is a solution of the problem (1), (12).

We have thus proved that  $\dim W \geq 1$ . On the other hand, by Lemma 1 any solution  $u \in W$  satisfies (8) and is therefore a Kneser-type solution. To complete the proof it remains only to show that  $\dim W \leq 1$ , i.e., that the problem (1), (12) has at most one solution for any  $c \in R$ . Let  $u_1$  and  $u_2$  be two arbitrary solutions of this problem and  $u_0(t) = u_2(t) - u_1(t)$ . Since  $u_0 \in W$  and  $u_0(a_0) = 0$ , by Lemma 1

$$2 \int_{a_0}^{+\infty} (s - a_0)u'^2_0(s)ds = 0 \quad \text{and} \quad u_0(t) = 0 \quad \text{for } t \geq a_0,$$

i.e.,  $u_1(t) \equiv u_2(t)$ .  $\square$

*Remark 1.* The condition (4) of Theorem 1 cannot be replaced by the condition

$$p^2_2(t) \leq (4 + \varepsilon)\tau'_2(t)p_1(t) \quad \text{for } t \geq a. \tag{16}$$

Indeed, consider the equation

$$u''(t) = \frac{1}{(4 + \varepsilon)t^2}u(t) - \frac{1}{t}u'(t), \quad (17)$$

satisfying all conditions of Theorem 1 except (4), instead of which the condition (16) is fulfilled. On the other hand, the equation (17) has the solutions  $u_i(t) = t^{\lambda_i}$  ( $i = 1, 2$ ), where  $\lambda_i = (-1)^i(4 + \varepsilon)^{-\frac{1}{2}}$  ( $i = 1, 2$ ). Clearly,  $u_i \in W$  ( $i = 1, 2$ ). Therefore in our case instead of (5) we have  $K \subset W$ ,  $\dim W = 2$ .

**Corollary 1.** *Let the conditions of Theorem 1 be fulfilled. Let, moreover,*

$$p_2(t) \leq 0 \quad \text{for } t \geq a. \quad (18)$$

Then

$$K = W, \quad \dim K = 1. \quad (19)$$

*Proof.* Let  $u \in K$ . Then by virtue of (18) and the non-negativity of  $p_1$  there exists  $t_0 \in [a, +\infty[$  such that  $u(t)u'(t) \leq 0$ ,  $u''(t)u(t) \geq 0$  for  $t \geq t_0$ . Hence

$$\int_{t_0}^{+\infty} u'^2(s)ds \leq |u(t_0)u'(t_0)|.$$

Therefore  $u \in W$ . Thus we have proved that  $W \supset K$ . This fact, together with (5), implies (19).  $\square$

A solution  $u$  of the equation (1) will be called *vanishing at infinity* if

$$\lim_{t \rightarrow +\infty} u(t) = 0. \quad (20)$$

**Theorem 2.** *Let the conditions of Theorem 1 be fulfilled. Then for any solution  $u \in W$  to vanish at infinity it is necessary and sufficient that*

$$\int_a^{+\infty} sp_1(s)ds = +\infty. \quad (21)$$

*Proof.* Let  $u \in W$ . Then by Lemma 1  $u^2(t) \geq \eta$  for  $t \geq a_0$ , where  $\eta = \lim_{t \rightarrow +\infty} u^2(t)$ , and  $\int_{a_0}^{+\infty} (s - a_0)p_1(s)u^2(s)ds \leq u^2(a_0)/2\delta$ . Hence it follows that (21) implies  $\eta = 0$ , i.e.,  $u$  is a vanishing solution at infinity.

To complete the proof it is enough to establish that if

$$\int_a^{+\infty} sp_1(s)ds < +\infty, \quad (22)$$

then any nontrivial solution  $u \in W$  tends to a nonzero limit as  $t \rightarrow +\infty$ . Let us assume the contrary: the equation (1) has a nontrivial solution  $u \in W$  vanishing at infinity. Then by Lemma 1

$$u(t)u'(t) \leq 0, \quad \rho(t) \leq \eta^2 u^2(t) \quad \text{for } t \geq a_0, \tag{23}$$

where

$$\rho(t) = \int_t^{+\infty} (s-t)[u'^2(s) + p_1(s)u^2(s)] ds, \quad \eta = (2\delta)^{-\frac{1}{2}}.$$

On the other hand, by (4), (20) and (22) we have

$$\begin{aligned} |u(t)| &= \left| \int_t^{+\infty} (s-t)[p_1(s)u(\tau_1(s)) + p_2(s)u'(\tau_2(s))] ds \right| \leq \\ &\leq \left[ \int_t^{+\infty} (s-t)p_1(s) ds \right]^{1/2} \left[ \int_t^{+\infty} (s-t)p_1(s)u^2(\tau_1(s)) ds \right]^{1/2} + \\ &\quad + 2 \int_t^{+\infty} (s-t)[p_1(s)]^{1/2} [\tau_2'(s)]^{1/2} |u'(\tau_2(s))| ds \leq \\ &\leq \left[ \int_t^{+\infty} (s-t)p_1(s) ds \right]^{1/2} \left[ \int_t^{+\infty} (s-t)p_1(s)u^2(\tau_1(s)) ds \right]^{1/2} + \\ &\quad + 2 \left[ \int_t^{+\infty} (s-t)p_1(s) ds \right]^{1/2} \left[ \int_t^{+\infty} (s-t)\tau_2'(s)u'^2(\tau_2(s)) ds \right]^{1/2} \\ &\quad \text{for } t \geq a_0. \end{aligned}$$

Hence by (2) and (23) we find

$$\begin{aligned} |u(t)| &\leq \left[ \int_t^{+\infty} (s-t)p_1(s) ds \right]^{1/2} \left[ \int_t^{+\infty} (s-t)p_1(s)u^2(s) ds \right]^{1/2} + \\ &\quad + 2 \left[ \int_t^{+\infty} (s-t)p_1(s) ds \right]^{1/2} \left[ \int_t^{+\infty} (s-t)u'^2(s) ds \right]^{1/2} \leq \\ &\leq 3\eta \left[ \int_t^{+\infty} (s-t)p_1(s) ds \right]^{1/2} |u(t)| \quad \text{for } t \geq a_0 \end{aligned}$$

and therefore  $u(t) = 0$  for  $t \geq a_1$ , where  $a_1$  is a sufficiently large number. By virtue of (2) the last equality implies  $u(t) = 0$  for  $t \geq a$ . But this is impossible, since by our assumption  $u$  is a nontrivial solution. The obtained contradiction proves the theorem.  $\square$

## REFERENCES

1. I.T. Kiguradze, On the non-negative nonincreasing solutions of nonlinear second order differential equations. *Ann. mat. pura ed appl.* **81**(1969), 169-192.
2. I.T. Kiguradze and D.I. Chichua, Kneser's problem for functional differential equations. (Russian) *Differentsial'nye Uravneniya* **27**(1992), No. 11, 1879-1892.

(Received 03.08.1993)

Author's address:  
A. Razmadze Mathematical Institute  
Georgian Academy of Sciences  
1, Z. Rukhadze St., Tbilisi 380093  
Republic of Georgia