

## ON SOME CONVEXITY PROPERTIES OF GENERALIZED CESÁRO SEQUENCE SPACES

SUTHEP SUANTAI

**Abstract.** We define a generalized Cesáro sequence space and consider it equipped with the Luxemburg norm under which it is a Banach space, and we show that it is locally uniformly rotund.

**2000 Mathematics Subject Classification:** 46E30, 46E40, 46B20.

**Key words and phrases:** Generalized Cesáro sequence spaces, Luxemburg norm, extreme point, locally uniformly rotund point, property (H), convex modular.

### 1. PRELIMINARIES

For a Banach space  $X$ , we denote by  $S(X)$  and  $B(X)$  the unit sphere and unit ball of  $X$ , respectively. A point  $x_0 \in S(X)$  is called

a) an *extreme point* if for every  $x, y \in S(X)$  the equality  $2x_0 = x + y$  implies  $x = y$ ;

b) a *locally uniformly rotund point* (LUR-point for short) if for any sequence  $(x_n)$  in  $B(X)$  such that  $\|x_n + x\| \rightarrow 2$  as  $n \rightarrow \infty$  there holds  $\|x_n - x\| \rightarrow 0$  as  $n \rightarrow \infty$ ;

c) an *H-point* if for any sequence  $(x_n)$  in  $X$  such that  $\|x_n\| \rightarrow 1$  as  $n \rightarrow \infty$ , the weak convergence of  $(x_n)$  to  $x_0$  (write  $x_n \xrightarrow{w} x_0$ ) implies that  $\|x_n - x_0\| \rightarrow 0$  as  $n \rightarrow \infty$ .

A Banach space  $X$  is said to be *rotund* (R) if every point of  $S(X)$  is an extreme point.

If every  $x \in S(X)$  is a LUR-point, then  $X$  is said to be *locally uniformly rotund* (LUR).

$X$  is said to possess property (H) provided every point of  $S(X)$  is an *H-point*.

For these geometric notions and their role in Mathematics we refer to the monographs [1], [6], [12] and [13]. Some of them were studied for Orlicz spaces in [1], [7], [8], [12] and [14].

Let  $X$  be a real vector space. A functional  $\varrho : X \rightarrow [0, \infty]$  is called a *modular* if it satisfies the conditions

- (i)  $\varrho(x) = 0$  if and only if  $x = 0$ ;
- (ii)  $\varrho(\alpha x) = \varrho(x)$  for all scalar  $\alpha$  with  $|\alpha| = 1$ ;
- (iii)  $\varrho(\alpha x + \beta y) \leq \varrho(x) + \varrho(y)$  for all  $x, y \in X$  and all  $\alpha, \beta \geq 0$  with  $\alpha + \beta = 1$ .  
The modular  $\varrho$  is called *convex* if
- (iv)  $\varrho(\alpha x + \beta y) \leq \alpha \varrho(x) + \beta \varrho(y)$  for all  $x, y \in X$  and all  $\alpha, \beta \geq 0$  with  $\alpha + \beta = 1$ .

If  $\varrho$  is a modular in  $X$ , we define

$$X_\varrho = \{x \in X : \lim_{\lambda \rightarrow 0^+} \varrho(\lambda x) = 0\},$$

$$\text{and } X_\varrho^* = \{x \in X : \varrho(\lambda x) < \infty \text{ for some } \lambda > 0\}.$$

It is clear that  $X_\varrho \subseteq X_\varrho^*$ . If  $\varrho$  is a convex modular, for  $x \in X_\varrho$  we define

$$\|x\| = \inf \left\{ \lambda > 0 : \varrho\left(\frac{x}{\lambda}\right) \leq 1 \right\}. \quad (1.1)$$

Orlicz [13] proved that if  $\varrho$  is a convex modular in  $X$ , then  $X_\varrho = X_\varrho^*$  and  $\|\cdot\|$  is a norm on  $X_\varrho$  for which it is a Banach space. The norm  $\|\cdot\|$  defined as in (1.1) is called the Luxemburg norm.

A modular  $\varrho$  on  $X$  is called

- (a) *right-continuous* if  $\lim_{\lambda \rightarrow 1^+} \varrho(\lambda x) = \varrho(x)$  for all  $x \in X_\varrho$ ;
- (b) *left-continuous* if  $\lim_{\lambda \rightarrow 1^-} \varrho(\lambda x) = \varrho(x)$  for all  $x \in X_\varrho$ ;
- (c) *continuous* if it is both left-continuous and right-continuous.

The following known results gave some relationships between the modular  $\varrho$  and the Luxemburg norm  $\|\cdot\|$  on  $X_\varrho$ .

**Theorem 1.1.** *Let  $\varrho$  be a convex modular on  $X$  and let  $x \in X_\varrho$  and  $(x_n)$  a sequence in  $X_\varrho$ . Then  $\|x_n - x\| \rightarrow 0$  as  $n \rightarrow \infty$  if and only if  $\varrho(\lambda(x_n - x)) \rightarrow 0$  as  $n \rightarrow \infty$  for every  $\lambda > 0$ .*

*Proof.* See [11, Theorem 1.3]. □

**Theorem 1.2.** *Let  $\varrho$  be a convex modular on  $X$  and  $x \in X_\varrho$ .*

- (i) *If  $\varrho$  is right-continuous, then  $\|x\| < 1$  if and only if  $\varrho(x) < 1$ .*
- (ii) *If  $\varrho$  is left-continuous, then  $\|x\| \leq 1$  if and only if  $\varrho(x) \leq 1$ .*
- (iii) *If  $\varrho$  is continuous, then  $\|x\| = 1$  if and only if  $\varrho(x) = 1$ .*

*Proof.* See [11, Theorem 1.4]. □

Let us denote by  $l^0$  the space of all real sequences. For  $1 \leq p < \infty$ , the Cesàro sequence space ( $ces_p$ , for short) is defined by

$$ces_p = \left\{ x \in l^0 : \sum_{n=1}^{\infty} \left( \frac{1}{n} \sum_{i=1}^n |x(i)| \right)^p < \infty \right\}$$

equipped with the norm

$$\|x\| = \left( \sum_{n=1}^{\infty} \left( \frac{1}{n} \sum_{i=1}^n |x(i)| \right)^p \right)^{\frac{1}{p}}.$$

This space was introduced by J.S. Shue [16]. It is useful in the theory of matrix operators and others (see [9] and [10]). Some geometric properties of the Cesàro sequence space  $ces_p$  were studied by many mathematicians. It is known that  $ces_p$  is LUR and possesses property (H) (see [10]). Y. A. Cui and H. Hudzik [2] proved that  $ces_p$  has the Banach-Saks property, and it was shown in [5] that  $ces_p$  has property  $(\beta)$ .

Now let  $p = (p_k)$  be a sequence of positive real numbers with  $p_k \geq 1$  for all  $k \in \mathbb{N}$ . The Nakano sequence space  $l(p)$  is defined by

$$l(p) = \{x \in l^0 : \sigma(\lambda x) < \infty \text{ for some } \lambda > 0\},$$

where  $\sigma(x) = \sum_{i=1}^{\infty} |x(i)|^{p_i}$ . We consider the space  $l(p)$  equipped with the norm

$$\|x\| = \inf \left\{ \lambda > 0 : \sigma\left(\frac{x}{\lambda}\right) \leq 1 \right\}$$

under which it is a Banach space. If  $p = (p_k)$  is bounded, we have

$$l(p) = \left\{ x \in l^0 : \sum_{i=1}^{\infty} |x(i)|^{p_i} < \infty \right\}.$$

Several geometric properties of  $l(p)$  were studied in [1] and [4].

The generalized Cesàro sequence space  $ces(p)$  is defined by

$$ces(p) = \{x \in l^0 : \varrho(\lambda x) < \infty \text{ for some } \lambda > 0\},$$

where  $\varrho(x) = \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{i=1}^n |x(i)|\right)^{p_n}$ . We consider this space equipped with the so-called Luxemburg norm

$$\|x\| = \inf \left\{ \lambda > 0 : \varrho\left(\frac{x}{\lambda}\right) \leq 1 \right\}$$

under which it is a Banach space. If  $p = (p_k)$  is bounded, we have

$$ces(p) = \left\{ x = x(i) : \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{i=1}^n |x(i)|\right)^{p_n} < \infty \right\}.$$

W. Sanhan [15] proved that  $ces(p)$  is nonsquare when  $p_k > 1$  for all  $k \in \mathbb{N}$ . In this paper, we show that the Cesàro sequence space  $ces(p)$  equipped with the Luxemburg norm is  $LUR$  and has property  $(H)$  when  $p = (p_k)$  is bounded with  $p_k > 1$  for all  $k \in \mathbb{N}$ .

Throughout this paper we assume that  $p = (p_k)$  is bounded with  $p_k > 1$  for all  $k \in \mathbb{N}$ , and  $M = \sup_k p_k$ .

## 2. MAIN RESULTS

We begin by giving some basic properties of the modular  $\varrho$  on the space  $ces(p)$ . By the convexity of the function  $t \rightarrow |t|^{p_k}$ , for every  $k \in \mathbb{N}$  we have that  $\varrho$  is a convex modular. So we have the following proposition.

**Proposition 2.1.** *The functional  $\varrho$  on the Cesàro sequence space  $ces(p)$  is a convex modular.*

**Proposition 2.2.** *For  $x \in ces(p)$ , the modular  $\varrho$  on  $ces(p)$  satisfies the following properties:*

- (i) if  $0 < a < 1$ , then  $a^M \varrho\left(\frac{x}{a}\right) \leq \varrho(x)$  and  $\varrho(ax) \leq a\varrho(x)$ ,
- (ii) if  $a \geq 1$ , then  $\varrho(x) \leq a^M \varrho\left(\frac{x}{a}\right)$ ,
- (iii) if  $a \geq 1$ , then  $\varrho(x) \leq a\varrho(x) \leq \varrho(ax)$ .

*Proof.* All assertions are clearly obtained by the definition of  $\varrho$ .  $\square$

**Proposition 2.3.** *The modular  $\varrho$  on  $ces(p)$  is continuous.*

*Proof.* For  $\lambda > 1$ , by Proposition 2.2 (ii) and (iii), we have

$$\varrho(x) \leq \lambda\varrho(x) \leq \varrho(\lambda x) \leq \lambda^M \varrho(x). \quad (2.1)$$

By taking  $\lambda \rightarrow 1^+$  in (2.1), we have  $\lim_{\lambda \rightarrow 1^+} \varrho(\lambda x) = \varrho(x)$ . Thus  $\varrho$  is right-continuous. If  $0 < \lambda < 1$ , by Proposition 2.2 (i), we have

$$\lambda^M \varrho(x) \leq \varrho(\lambda x) \leq \lambda\varrho(x) \quad (2.2)$$

By taking  $\lambda \rightarrow 1^-$  in (2.2), we have that  $\lim_{\lambda \rightarrow 1^-} \varrho(\lambda x) = \varrho(x)$ , hence  $\varrho$  is left-continuous. Thus  $\varrho$  is continuous.  $\square$

Next, we give some relationships between the modular  $\varrho$  and the Luxemburg norm on  $ces(p)$ .

**Proposition 2.4.** *For any  $x \in ces(p)$ , we have*

- (i) *if  $\|x\| < 1$ , then  $\varrho(x) \leq \|x\|$ ,*
- (ii) *if  $\|x\| > 1$ , then  $\varrho(x) \geq \|x\|$ ,*
- (iii)  *$\|x\| = 1$  if and only if  $\varrho(x) = 1$ ,*
- (iv)  *$\|x\| < 1$  if and only if  $\varrho(x) < 1$ ,*
- (v)  *$\|x\| > 1$  if and only if  $\varrho(x) > 1$ ,*
- (vi) *if  $0 < a < 1$  and  $\|x\| > a$ , then  $\varrho(x) > a^M$ , and*
- (vii) *if  $a \geq 1$  and  $\|x\| < a$ , then  $\varrho(x) < a^M$ .*

*Proof.* If  $\|x\| \leq 1$ , it follows by the convexity and continuity of  $\varrho$  that  $\varrho(x) = \varrho\left(\|x\| \frac{x}{\|x\|}\right) \leq \|x\| \varrho\left(\frac{x}{\|x\|}\right) \leq \|x\|$ . So (i) is obtained. If  $\|x\| > 1$ , then there is  $\varepsilon_0 > 0$  such that  $\|x\| - \varepsilon > 1$  for all  $\varepsilon \in (0, \varepsilon_0)$ . Consequently,  $\varrho(x) = \varrho\left((\|x\| - \varepsilon) \frac{x}{\|x\| - \varepsilon}\right) \geq (\|x\| - \varepsilon) \varrho\left(\frac{x}{\|x\| - \varepsilon}\right) > \|x\| - \varepsilon$ , so (ii) is satisfied. It is clear that (iii), (iv) and (v) follow by Theorem 1.2, and properties (vi) and (vii) follow by Proposition 2.2.  $\square$

**Proposition 2.5.** *Let  $(x_n)$  be a sequence in  $ces(p)$ .*

- (i) *If  $\|x_n\| \rightarrow 1$  as  $n \rightarrow \infty$ , then  $\varrho(x_n) \rightarrow 1$  as  $n \rightarrow \infty$ .*
- (ii)  *$\|x_n\| \rightarrow 0$  as  $n \rightarrow \infty$  if and only if  $\varrho(x_n) \rightarrow 0$  as  $n \rightarrow \infty$ .*

*Proof.* (i) Suppose  $\|x_n\| \rightarrow 1$  as  $n \rightarrow \infty$ . Let  $\epsilon \in (0, 1)$ . Then there exists  $N \in \mathbb{N}$  such that  $1 - \epsilon < \|x_n\| < 1 + \epsilon$  for all  $n \geq N$ . By Proposition 2.4 (vi) and (vii), we have  $(1 - \epsilon)^M < \varrho(x_n) < (1 + \epsilon)^M$  for all  $n \geq N$ , which implies that  $\varrho(x_n) \rightarrow 1$  as  $n \rightarrow \infty$ .

(ii) It follows from Theorem 1.1 that if  $\|x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ , then  $\varrho(x_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Conversely, suppose  $\|x_n\| \not\rightarrow 0$  as  $n \rightarrow \infty$ . Then there is  $\epsilon \in (0, 1)$  and a subsequence  $(x_{n_k})$  of  $(x_n)$  such that  $\|x_{n_k}\| > \epsilon$  for all  $k \in \mathbb{N}$ . By Proposition 2.4 (vi), we have  $\varrho(x_{n_k}) > \epsilon^M$  for all  $k \in \mathbb{N}$ . This implies  $\varrho(x_n) \not\rightarrow 0$  as  $n \rightarrow \infty$ .  $\square$

**Proposition 2.6.** *Let  $(x_n) \subseteq B(l(p))$  and  $(y_n) \subseteq B(l(p))$ . If  $\sigma\left(\frac{x_n + y_n}{2}\right) \rightarrow 1$ , then  $x_n(i) - y_n(i) \rightarrow 0$  as  $n \rightarrow \infty$  for all  $i \in \mathbb{N}$ .*

*Proof.* We first note that if  $x \in B(l(p))$ , then  $\sigma(x) \leq 1$ . Suppose that  $x_n(i) - y_n(i) \not\rightarrow 0$  as  $n \rightarrow \infty$  for some  $i \in \mathbb{N}$ . Without loss of generality we may assume that  $i = 1$ , and then assume without loss of generality (passing to a subsequence if necessary) that, for some  $\epsilon > 0$ ,

$$|x_n(1) - y_n(1)|^{p_1} \geq \epsilon \quad \forall n \in \mathbb{N}.$$

Thus

$$2^{p_1}(|x_n(1)|^{p_1} + |y_n(1)|^{p_1}) \geq \epsilon \quad \forall n \in \mathbb{N}. \tag{2.3}$$

Since the function  $t \rightarrow |t|^{p_1}$  is uniformly convex, there exists  $\delta > 0$  such that

$$\left| \frac{x_n(1) + y_n(1)}{2} \right|^{p_1} \leq (1 - \delta) \left( \frac{|x_n(1)|^{p_1} + |y_n(1)|^{p_1}}{2} \right) \quad \forall n \in \mathbb{N}. \tag{2.4}$$

It follows from (2.3) and (2.4) that for each  $n \in \mathbb{N}$ ,

$$\begin{aligned} \sigma\left(\frac{x_n + y_n}{2}\right) &= \sum_{i=1}^{\infty} \left| \frac{x_n(i) + y_n(i)}{2} \right|^{p_i} \\ &= \left| \frac{x_n(1) + y_n(1)}{2} \right|^{p_1} + \sum_{i=2}^{\infty} \left| \frac{x_n(i) + y_n(i)}{2} \right|^{p_i} \\ &\leq (1 - \delta) \left( \frac{|x_n(1)|^{p_1} + |y_n(1)|^{p_1}}{2} \right) + \frac{1}{2} \sum_{i=2}^{\infty} |x_n(i)|^{p_i} + \frac{1}{2} \sum_{i=2}^{\infty} |y_n(i)|^{p_i} \\ &= \frac{1}{2} \sigma(x_n) + \frac{1}{2} \sigma(y_n) - \delta \left( \frac{|x_n(1)|^{p_1} + |y_n(1)|^{p_1}}{2} \right) \\ &\leq \frac{1}{2} + \frac{1}{2} - \delta \frac{\epsilon}{2^{p_1+1}} = 1 - \delta \frac{\epsilon}{2^{p_1+1}}. \end{aligned}$$

This implies that  $\sigma\left(\frac{x_n + y_n}{2}\right) \not\rightarrow 1$  as  $n \rightarrow \infty$ , a contradiction, which finishes the proof.  $\square$

**Proposition 2.7.** *Let  $(x_n) \subseteq B(ces(p))$  and  $x \in S(ces(p))$ . If  $\varrho\left(\frac{x_n + x}{2}\right) \rightarrow 1$  as  $n \rightarrow \infty$ , then  $x_n(i) \rightarrow x(i)$  as  $n \rightarrow \infty$  for all  $i \in \mathbb{N}$ .*

*Proof.* For each  $n \in \mathbb{N}$  and  $i \in \mathbb{N}$ , let

$$s_n(i) = \begin{cases} \operatorname{sgn}(x_n(i) + x(i)) & \text{if } x_n(i) + x(i) \neq 0, \\ 1 & \text{if } x_n(i) + x(i) = 0. \end{cases}$$

Hence we have

$$\begin{aligned} 1 \leftarrow \varrho\left(\frac{x_n + x}{2}\right) &= \sum_{k=1}^{\infty} \left(\frac{1}{k} \sum_{i=1}^k \left|\frac{x_n(i) + x(i)}{2}\right|\right)^{p_k} \\ &= \sum_{k=1}^{\infty} \left(\frac{1}{k} \sum_{i=1}^k s_n(i) \frac{x_n(i)}{2} + \frac{1}{k} \sum_{i=1}^k s_n(i) \frac{x(i)}{2}\right)^{p_k}. \end{aligned} \quad (2.5)$$

Let  $a_n(k) = \frac{1}{k} \sum_{i=1}^k s_n(i)x_n(i)$  and  $b_n(k) = \frac{1}{k} \sum_{i=1}^k s_n(i)x(i)$  for all  $n, k \in \mathbb{N}$ . Then  $(a_n) \in l(p)$  and  $(b_n) \in l(p)$ , and from (2.5) we have

$$\sigma\left(\frac{a_n + b_n}{2}\right) \rightarrow 1 \quad \text{as } n \rightarrow \infty.$$

Form Proposition 2.6 we have

$$a_n(i) - b_n(i) \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (2.6)$$

for all  $i \in \mathbb{N}$ . Now we shall show that  $x_n(k) \rightarrow x(k)$  as  $n \rightarrow \infty$  for all  $k \in \mathbb{N}$ . From (2.6) we have

$$s_n(1)x_n(1) - s_n(1)x(1) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This implies  $x_n(1) \rightarrow x(1)$  as  $n \rightarrow \infty$ . Assume that  $x_n(i) \rightarrow x(i)$  as  $n \rightarrow \infty$  for all  $i \leq k-1$ . Then we have

$$s_n(i)(x_n(i) - x(i)) \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (2.7)$$

for all  $i \leq k-1$ . Since  $s_n(k)(x_n(k) - x(k)) = k(a_n(k) - b_n(k)) - \sum_{i=1}^{k-1} s_n(i)(x_n(i) - x(i))$ , it follows from (2.6) and (2.7) that  $s_n(k)(x_n(k) - x(k)) \rightarrow 0$  as  $n \rightarrow \infty$ . This implies  $x_n(k) \rightarrow x(k)$  as  $n \rightarrow \infty$ . So we have by induction that  $x_n(k) \rightarrow x(k)$  as  $n \rightarrow \infty$  for all  $k \in \mathbb{N}$ .  $\square$

**Theorem 2.8.** *The space  $ces(p)$  is LUR.*

*Proof.* Let  $(x_n) \subseteq B(ces(p))$  and  $x \in S(ces(p))$  be such that  $\|x_n + x\| \rightarrow 2$  as  $n \rightarrow \infty$ . Then  $\left\|\frac{x_n + x}{2}\right\| \rightarrow 1$  as  $n \rightarrow \infty$ . By Proposition 2.5 (i) we have  $\varrho\left(\frac{x_n + x}{2}\right) \rightarrow 1$  as  $n \rightarrow \infty$ . By Proposition 2.7 we have  $x_n(i) \rightarrow x(i)$  as  $n \rightarrow \infty$  for all  $i \in \mathbb{N}$ .

Now let  $\epsilon > 0$  be given. Then there exist  $k_0 \in \mathbb{N}$  and  $n_0 \in \mathbb{N}$  such that

$$\sum_{k=k_0+1}^{\infty} \left(\frac{1}{k} \sum_{i=1}^k |x(i)|\right)^{p_k} < \frac{\epsilon}{3} \frac{1}{2^{M+1}}, \quad (2.8)$$

$$\sum_{k=1}^{k_0} \left(\frac{1}{k} \sum_{i=1}^k |x_n(i) - x(i)|\right)^{p_k} < \frac{\epsilon}{3} \quad \text{for all } n \geq n_0, \quad (2.9)$$

$$\sum_{k=1}^{k_0} \left(\frac{1}{k} \sum_{i=1}^k |x_n(i)|\right)^{p_k} > \sum_{k=1}^{k_0} \left(\frac{1}{k} \sum_{i=1}^k |x(i)|\right)^{p_k} - \frac{\epsilon}{3} \frac{1}{2^M} \quad \text{for all } n \geq n_0. \quad (2.10)$$

By Proposition 2.4 (i) and (iii) we have  $\varrho(x_n) \leq 1$  for all  $n \in \mathbb{N}$  and  $\varrho(x) = 1$ . From these together with (2.8), (2.9), (2.10) and the fact that  $(a + b)^{p_k} \leq 2^{p_k}(a^{p_k} + b^{p_k})$  for  $a, b \geq 0$  we have that for all  $n \geq n_0$ ,

$$\begin{aligned} \varrho(x_n - x) &= \sum_{k=1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x_n(i) - x(i)| \right)^{p_k} \\ &= \sum_{k=1}^{k_0} \left( \frac{1}{k} \sum_{i=1}^k |x_n(i) - x(i)| \right)^{p_k} + \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x_n(i) - x(i)| \right)^{p_k} \\ &< \frac{\epsilon}{3} + 2^M \left( \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x_n(i)| \right)^{p_k} + \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} \right) \\ &= \frac{\epsilon}{3} + 2^M \left( \varrho(x_n) - \sum_{k=1}^{k_0} \left( \frac{1}{k} \sum_{i=1}^k |x_n(i)| \right)^{p_k} + \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} \right) \\ &\leq \frac{\epsilon}{3} + 2^M \left( 1 - \sum_{k=1}^{k_0} \left( \frac{1}{k} \sum_{i=1}^k |x_n(i)| \right)^{p_k} + \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} \right) \\ &< \frac{\epsilon}{3} + 2^M \left( 1 - \sum_{k=1}^{k_0} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} + \frac{\epsilon}{3} \frac{1}{2^M} + \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} \right) \\ &= \frac{\epsilon}{3} + 2^M \left( \varrho(x) - \sum_{k=1}^{k_0} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} + \frac{\epsilon}{3} \frac{1}{2^M} + \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} \right) \\ &= \frac{\epsilon}{3} + 2^M \left( \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} + \frac{\epsilon}{3} \frac{1}{2^M} + \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} \right) \\ &= \frac{\epsilon}{3} + 2^M \left( 2 \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} + \frac{\epsilon}{3} \frac{1}{2^M} \right) \\ &= \frac{\epsilon}{3} + 2^{M+1} \sum_{k=k_0+1}^{\infty} \left( \frac{1}{k} \sum_{i=1}^k |x(i)| \right)^{p_k} + \frac{\epsilon}{3} < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon. \end{aligned}$$

This shows that  $\varrho(x_n - x) \rightarrow 0$  as  $n \rightarrow \infty$ . By Proposition 2.5(ii) we have  $\|x_n - x\| \rightarrow 0$  as  $n \rightarrow \infty$ . This completes the proof of the theorem.  $\square$

It is known in general that a locally uniformly rotund space has property (H). So we have the following result.

**Corollary 2.9.** *The space  $ces(p)$  possesses property (H).*

ACKNOWLEDGEMENTS

The author would like to thank the Thailand Research Fund for the financial support and is very indebted to the referee for his valuable comments and suggestions.

## REFERENCES

1. S. T. CHEN, Geometry of Orlicz spaces. With a preface by Julian Musielak. *Dissertationes Math. (Rozprawy Mat.)* **356**(1996), 1–204.
2. Y. A. CUI and H. HUDZIK, On the Banach-Saks and weak Banach-Saks properties of some Banach sequence spaces. *Acta Sci. Math. (Szeged)* **65**(1999), No. 1-2, 179–187.
3. Y. A. CUI, H. HUDZIK and C. MENG, On some local geometry of Orlicz sequence spaces equipped the Luxemburg norms. *Acta Math. Hungar.* **80**(1998), No. 1-2, 143–154.
4. Y. A. CUI, H. HUDZIK and R. PLICIENNIK, Banach-Saks property in some Banach sequence spaces. *Ann. Polon. Math.* **65**(1997), No. 2, 193–202.
5. Y. A. CUI and C. MENG, Banach-Sak property and property  $(\beta)$  in Cesáro sequence spaces. *Southeast Asian Bull. Math.* **24**(2000), No. 2, 201–210.
6. J. DIESTEL, Geometry of Banach Spaces – Selected Topics. *Lecture Notes in Mathematics*, 485. *Springer-Verlag, Berlin–New York*, 1975.
7. H. HUDZIK, Orlicz spaces without strongly extreme points and without  $H$ -points. *Canad. Math. Bull.* **36**(1993), No. 2, 173–177.
8. H. HUDZIK and D. PALLASCHKE, On some convexity properties of Orlicz sequence spaces. *Math. Nachr.* **186**(1997), 167–185.
9. P. Y. LEE, Cesáro sequence spaces. *Math. Chronicle* **13**(1984), 29–45.
10. Y. Q. LIU, B. E. WU, and Y. P. LEE, Method of sequence spaces. (Chinese) *Guangdong of Science and Technology Press*, 1996.
11. L. MALIGRANDA, Orlicz spaces and interpolation. *Seminários de Matemática*. 5, 1–206. *Campinas, SP: Univ. Estadual de Campinas, Dep. de Matemática*, 1989.
12. J. MUSIELAK, Orlicz spaces and modular spaces. *Lecture Notes in Math.* 1034, *Springer-Verlag, Berlin*, 1983.
13. W. ORLICZ, A note on modular spaces I. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **9**(1961), 157–162.
14. R. PLUCIENNIK, T.F WANG and Y. L. ZHANG,  $H$ -points and Denting Points in Orlicz Spaces. *Comment. Math. Prace Mat.* **33**(1993), 135–151.
15. W. SANHAN, On geometric properties of some Banach sequence spaces. *Thesis for the degree of Master of Science in Mathematics, Chiang Mai University*, 2000.
16. J.-S. SHIUE, On the Cesáro sequence spaces. *Tamkang J. Math.* **1**(1970), 19–25.

(Received 20.11.2001; revised 27.05.2002)

Author's address:

Department of Mathematics

Faculty of Science

Chiang Mai University, Chiang Mai, 50200

Thailand

E-mail: suantai@yahoo.com