

ON ARU-RESOLUTIONS OF UNIFORM SPACES

V. BALADZE

Abstract. In this paper theorems which give conditions for a uniform space to have an *ARU*-resolution are proved. In particular, a finitistic uniform space admits an *ARU*-resolution if and only if it has trivial uniform shape or it is an absolute uniform shape retract.

2000 Mathematics Subject Classification: 54C56, 54E15.

Key words and phrases: Uniform shape, uniform resolution, absolute uniform shape retract.

1. INTRODUCTION

The definition of resolution of topological spaces was introduced by S. Mardešić [8] and it was used to define and study the shapes of topological spaces [10]. More recently, J. Segal, S. Spiez and B. Günter [14] defined a uniform version of resolution and proved that every finitistic uniform space has *ANRU*-resolution. They and T. Miyata [11] showed its usefulness in uniform shape theory.

The natural question is to determine which uniform spaces admit *ARU*-resolutions, i.e. uniform resolutions $\mathbf{p} : X \rightarrow \mathbf{X} = (X_\lambda, p_{\lambda\lambda'}, \Lambda)$, where each X_λ is an *ARU*-space. In [5] A. Koyama, S. Mardešić and T. Watanabe proved that a topological space X has an *AR*-resolution if and only if X has trivial shape.

In this paper we prove that for every finitistic uniform space X the following assertions are equivalent:

- (i) X admits an *ARU*-resolution.
- (ii) X has trivial uniform shape.
- (iii) X is an absolute uniform shape retract.

Without specific references we use results from [4], [10], [11] and [14]. A uniform space X is said to be an *A(N)RU*-space if whenever X is embedded in a uniform space Y , then there is a uniform retraction of Y (some uniform neighborhood of X in Y) onto X , equivalently, if whenever A is a uniform subspace of Y , then every uniform map $f : A \rightarrow X$ extends over Y (some uniform neighborhood of A in Y). We say that two uniform maps $f, g : X \rightarrow Y$ are semi-uniformly homotopic maps and write $f \simeq_u g$ if there exists a uniform map $H : X * I \rightarrow Y$ such that $H_0 = f$ and $H_1 = g$. Here $X * I$ is the semi-uniform product of X and $I = [0, 1]$ (see [4, p. 44]).

A uniform space X is said to be semi-uniform contractible [11] if the identity uniform map $1_X : X \rightarrow X$ and some constant uniform map $c : X \rightarrow X$ are semi-uniformly homotopic maps. It is clear that a uniform space is semi-uniform

contractible if and only if it has the semi-uniform homotopy type of a uniform space consisting of one point.

Let **Unif** denote the category of uniform spaces and uniform maps. By **A(N)RU** we denote the full subcategory of **Unif** whose objects are $A(N)RU$ -spaces. Also, we write **H(Unif)** and **H(A(N)RU)** for the semi-uniform homotopy category of **Unif** and **A(N)RU**, respectively.

There have been uniform shape theories defined by many mathematicians ([1], [2], [3], [12], [13]), based on the uniform homotopy [4]. Recently, T. Miyata [11] introduced a new uniform shape theory by the inverse system approach, using the semi-uniform homotopy which is weaker than the uniform homotopy. Here we consider T. Miyata's uniform shape theory which is convenient for our purpose. The uniform shape category **uSh** is the abstract shape category **Sh**(\mathcal{T}, \mathcal{P}) [10], where $\mathcal{T} = \mathbf{H}(\mathbf{Unif})$ and $\mathcal{P} = \mathbf{H}(\mathbf{ANRU})$. The uniform shape morphism $F : X \rightarrow Y$ of category **uSh** is given by a triple $(\mathbf{p}, \mathbf{q}, \mathbf{f})$, where $\mathbf{p} : X \rightarrow \mathbf{X} = (X_\lambda, p_{\lambda\lambda'}, \Lambda) \in \mathbf{pro} - \mathbf{H}(\mathbf{ANRU})$ and $\mathbf{q} : Y \rightarrow \mathbf{Y} = (Y_\mu, q_{\mu\mu'}, M) \in \mathbf{pro} - \mathbf{H}(\mathbf{ANRU})$ are **H(ANRU)**-expansions of X and Y , respectively, and $\mathbf{f} : \mathbf{X} \rightarrow \mathbf{Y}$ is a morphism of category **pro-H(ANRU)**. Two uniform spaces X and Y are said to have the same uniform shape denoted by $ush(X) = ush(Y)$, if X and Y are isomorphic objects of category **uSh**. We say that a uniform space X has trivial uniform shape and write $ush(X) = 0$ if X has the uniform shape of one point $\{*\}$.

2. ARU-RESOLUTION

A uniform resolution $\mathbf{p} = (p_\lambda) : X \rightarrow \mathbf{X} = (X_\lambda, p_{\lambda\lambda'}, \Lambda)$ of a uniform space X consists of an inverse system \mathbf{X} in **Unif** and of a morphism \mathbf{p} in **pro-Unif**, which has the property that for every $ANRU$ -space P and every uniform covering \mathcal{V} of P the following two conditions are satisfied:

(UR1) For every uniform map $f : X \rightarrow P$ there exist $\lambda \in \Lambda$ and a uniform map $h : X_\lambda \rightarrow P$ such that the maps $h \cdot p_\lambda$ and f are \mathcal{V} -near, i.e. every $x \in X$ admits $V \in \mathcal{V}$ such that $h \cdot p_\lambda(x), f(x) \in V$.

(UR2) There exists a uniform covering \mathcal{V}' of P such that if for $\lambda \in \Lambda$ and for two uniform maps $h, h' : X_\lambda \rightarrow P$ the maps $h \cdot p_\lambda, h' \cdot p_\lambda$ are \mathcal{V}' -near, then there exists $\lambda' \geq \lambda$ such that the maps $h \cdot p_{\lambda\lambda'}, h' \cdot p_{\lambda\lambda'}$ are \mathcal{V}' -near.

A uniform resolution is said to be cofinite if its index set Λ is cofinite. If all the terms $X_\lambda, \lambda \in \Lambda$, are $ANRU(ARU)$ -spaces, then we speak of an $ANRU(ARU)$ -resolution.

In [14] it is proved that every finitistic uniform space admits an $ANRU$ -resolution [14, Theorem 1]. A. Koyama, S. Mardešić and T. Watanabe raised the question of determining which spaces admit an AR -resolution, i.e. a resolution where each term X_λ is an absolute retract for metric spaces [5]. In this section we prove

Theorem 1. *Let X be a uniform space. Then the following assertions hold:*
 (i) *If X admits an ARU -resolution, then X has trivial uniform shape.*

(ii) *If X is a finitistic space and has trivial uniform shape, then X admits an ARU-resolution.*

The proof of this theorem is based on the following lemmas.

Lemma 2. *Let $\mathbf{X} = (X_\lambda, p_{\lambda\lambda'}, \Lambda)$ and $\mathbf{Y} = (Y_\mu, q_{\mu\mu'}, M)$ be the inverse systems of uniform spaces. If all Y_μ are ARU-spaces, then any two morphisms $(f_\mu, \varphi), (f'_\mu, \varphi') : \mathbf{X} \rightarrow \mathbf{Y}$ of category $\mathbf{inv} - \mathbf{Unif}$ are equivalent in the sense of morphisms [10].*

Proof. For each index $\mu \in M$ there exists an index $\lambda \in \Lambda$ such that $\lambda \geq \varphi(\mu), \varphi'(\mu)$. Let $Z = X_\lambda \times \{0\} \cup X_\lambda \times \{1\}$. Let $h : Z \rightarrow Y_\mu$ be a map defined by the formulas

$$h|_{X_\lambda \times \{0\}} = f_\mu \cdot p_{\varphi(\mu)\lambda}, \quad h|_{X_\lambda \times \{1\}} = f'_\mu \cdot p_{\varphi'(\mu)\lambda}.$$

It is clear that h is a uniform map. This map has a uniform extension $h' : X_\lambda * I \rightarrow Y_\mu$. Hence, $f_\mu \cdot p_{\varphi(\mu)\lambda}$ and $f'_\mu \cdot p_{\varphi'(\mu)\lambda}$ are semi-uniformly homotopic maps. Consequently, $(f_\lambda, \varphi) \sim (f'_\lambda, \varphi')$. \square

Lemma 3. *Let $f : X \rightarrow Y$ be a uniform map from a uniform space X into an ANRU-space Y . If the map f is semi-uniformly homotopic to a constant map, then there exist an ARU-space Z and a uniform map $r : Z \rightarrow Y$ such that $X \subseteq Z$ and $r|_X = f$.*

Proof. Consider X as a uniform subspace of an ARU-space Z . Since f is semi-uniform homotopic to a constant map $c : X \rightarrow Y$ and since c has a uniform extension on Z , the semi-uniform homotopy extension theorem (see [Theorem 1.6][11]) proves that $f : X \rightarrow Y$ too extends to a uniform map $r : Z \rightarrow Y$. \square

Lemma 4. *If a semi-uniform contractible space Y is an ANRU-space, then Y is an ARU-space.*

Proof. By the assumption of the lemma, there exist a point $y_0 \in Y$ and a semi-uniform homotopy $H : Y * I \rightarrow Y$ such that $H(y, 0) = y$ and $H(y, 1) = y_0$ for every $y \in Y$.

Let $f : A \rightarrow Y$ be a uniform map defined on a uniform subspace A of a uniform space X . Since Y is an ANRU-space, f has a uniform extension $f' : U \rightarrow Y$ over some uniform neighborhood U of A in X . There is a uniform covering \mathcal{U} of X such that $St(A, \mathcal{U}) \cap St(X \setminus U, \mathcal{U}) = \emptyset$. By the uniform version of Urysohn's lemma there exists a uniform map $e : X \rightarrow I$ such that

$$e(x) = \begin{cases} 0, & x \in A, \\ 1, & x \in X \setminus St(A, \mathcal{U}). \end{cases}$$

Then we define a map $g : X \rightarrow Y$ by

$$g(x) = \begin{cases} H(f'(x), e(x)), & x \in U, \\ y_0, & x \in X \setminus St(A, \mathcal{U}). \end{cases}$$

Let \mathcal{V} be an arbitrary uniform covering of X . There exists a uniform covering \mathcal{W} of X which is a refinement of the intersection $\mathcal{U} \wedge \mathcal{V}$ of uniform coverings

\mathcal{U} and \mathcal{V} . Let $W \in \mathcal{W}$ be an element of \mathcal{W} such that $W \cap U \neq \emptyset$ and $W \cap (X \setminus St(A, \mathcal{U})) \neq \emptyset$. Now show that $W \cap (U \cap (X \setminus St(A, \mathcal{U}))) \neq \emptyset$. First assume that $W \cap (X \setminus U) = \emptyset$. Then $W \subset U$ and, consequently, $W \cap (U \cap (X \setminus St(A, \mathcal{U}))) = W \cap (X \setminus St(A, \mathcal{U})) \neq \emptyset$. Now assume that $W \cap (X \setminus U) \neq \emptyset$. There exists an element $G \in \mathcal{U}$ such that $W \subset G$. Note that $G \cap (X \setminus U) \neq \emptyset$. Hence, $G \subset St(X \setminus U, \mathcal{U})$. Since $St(A, \mathcal{U}) \cap St(X \setminus U, \mathcal{U}) = \emptyset$, we have $G \cap St(A, \mathcal{U}) = \emptyset$. Consequently, $G \subset X \setminus St(A, \mathcal{U})$. We obtained that $G \cap (X \setminus St(A, \mathcal{U})) = G$. Note that $W \cap (X \setminus St(A, \mathcal{U})) = W$. This proves that $W \cap (U \cap (X \setminus St(A, \mathcal{U}))) = W \cap U \neq \emptyset$.

By Lemma 7 of [6] the map g is a uniform map. For each $a \in A$ we have

$$g(a) = H(f'(a), e(a)) = H(f'(a), 0) = f'(a) = f(a),$$

i.e. g is an extension of a uniform map $f : A \rightarrow Y$. Hence $Y \in ARU$. \square

Proof of Theorem 1. We can easily prove the first assertion by following the argument of (i) \Rightarrow (ii) in [5]. Here we use our Lemma 2 in the place of [5, Lemma 1]. Indeed, by the assumption (i) X admits an ARU -resolution $\mathbf{p} = (p_\lambda) : X \rightarrow \mathbf{X} = (X_\lambda, p_{\lambda\lambda'}, \Lambda)$. The space $Y = \{*\}$ consisting of one point $*$ also admits an ARU -resolution $\mathbf{q} = (q_\mu) : Y \rightarrow \mathbf{Y} = (Y_\mu, q_{\mu\mu'}, M)$, where $M = \{\mu_0\}$, $q_{\mu\mu'} = q_{\mu_0\mu_0} = 1_{Y_{\mu_0}} = 1_Y$, $Y_\mu = Y_{\mu_0} = Y$, $q_\mu = q_{\mu_0} = 1_Y$.

Let $f : X \rightarrow Y$ and $g : Y \rightarrow X$ be constant uniform maps. Let $(f_\mu, \varphi) : \mathbf{X} \rightarrow \mathbf{Y}$ and $(g_\lambda, \psi) : \mathbf{Y} \rightarrow \mathbf{X}$ be morphisms induced by f and g , respectively. By Lemma 3 we have

$$(f_\mu, \varphi) \cdot (g_\lambda, \psi) \sim (1_{Y_\mu}, 1_M), \quad (g_\lambda, \psi) \cdot (f_\mu, \varphi) \sim (1_{X_\lambda}, 1_\Lambda).$$

Let $F : X \rightarrow Y$ and $G : Y \rightarrow X$ be uniform shape morphisms with representatives $\mathbf{f} = [(f_\mu, \varphi)] : \mathbf{X} \rightarrow \mathbf{Y}$ and $\mathbf{g} = [(g_\lambda, \psi)] : \mathbf{Y} \rightarrow \mathbf{X}$. It is clear that $F \cdot G = I_Y$ and $G \cdot F = I_X$, where I_X and I_Y are the identity uniform shape morphisms. Hence $ush(X) = ush(Y)$, i.e. $ush(X) = 0$.

Now we will prove the second assertion. Since X is finitistic, there is an antisymmetric cofinite $ANRU$ -resolution $\mathbf{p} = (p_\lambda) : X \rightarrow \mathbf{X} = (X_\lambda, p_{\lambda\lambda'}, \Lambda)$ of X [14]. Using this $ANRU$ -resolution, the construction in the argument (iii) \Rightarrow (i) of [5] and our Lemma 3 in the place of [5, Lemma 2] and also Lemma 4 in next, we can define the ARU -system $\mathbf{Z} = (Z_\lambda, q_{\lambda\lambda'}, \Lambda^*)$ and the morphism $\mathbf{q} = (q_\lambda) : X \rightarrow \mathbf{Z}$.

Following the argument in [5] that the obtained resolution satisfies properties (R1) and (R2), we can easily show that \mathbf{q} has properties (UR1) and (UR2). \square

Remark. Following the approach by [9], we can construct the coherent semi-uniform homotopy category and, hence, uniform strong shape theory for uniform spaces by the inverse system approach based on semi-uniform homotopy. It is possible to find a condition in terms of uniform strong shape for a uniform finitistic space to have an ARU -resolution.

3. UNIFORM SHAPE RETRACTS

In this section we generalize the notion of an absolute shape retract [7] to the uniform spaces. Let A be a subspace of uniform space X . A uniform shape retraction is a uniform shape morphism $R : X \rightarrow A$ such that $R \cdot J = I_A$, where $J : A \rightarrow X$ is the uniform shape morphism induced by the inclusion uniform map $j : A \rightarrow X$. In this case the subspace A is called a uniform shape retract of X . Notice that if a uniform shape morphism $R : X \rightarrow A$ is induced by a uniform retraction $r : X \rightarrow A$, then R is a uniform shape retraction. If $A \subseteq Y \subseteq X$ and A is a uniform shape retract of X , then A is also a uniform shape retract of Y . Moreover, if A is a uniform shape retract of Y and Y is a uniform shape retract of X , then A is a uniform shape retract of X .

A uniform space X is called an absolute uniform shape retract provided for every uniform space Y with $X \subset Y$, X is a uniform shape retract of Y .

We have

Theorem 5. *Let X be a uniform space. Then the following assertions are equivalent:*

- (i) X is an absolute uniform shape retract.
- (ii) X has trivial uniform shape.

We first establish

Lemma 6. *Let X and Y be uniform spaces. If Y has trivial uniform shape, then there exists a unique uniform shape morphism $F : X \rightarrow Y$.*

Proof. Let $ush(Y) = 0$. The uniform shape morphism $C : Y \rightarrow \{*\}$, induced by the constant map $c : Y \rightarrow \{*\}$, has an inverse uniform shape morphism $G : \{*\} \rightarrow Y$, $G \cdot C = I_Y$. For each uniform shape morphism $F : X \rightarrow Y$ we have $F = G \cdot C \cdot F = G \cdot C'$, where $C' = C \cdot F : X \rightarrow \{*\}$ is the only uniform shape morphism into $\{*\}$. \square

Proof of Theorem 5. Implication (i) \Rightarrow (ii). Let X be an absolute uniform shape retract. We can consider X as a uniform subspace of an ARU-space Z . Note that Z is a semi-uniform contractible space. Consequently, Z has the semi-uniform homotopy type of a point, and $ush(Z) = 0$. By condition, there exists a uniform shape retraction $R : Z \rightarrow X$, $R \cdot J = I_X$, where $J : X \rightarrow Z$ is the uniform shape morphism induced by the inclusion uniform map $j : X \rightarrow Z$. From Lemma 6 it follows that $J \cdot R = I_Z$. Thus $ush(X) = ush(Z) = 0$.

Implication (ii) \Rightarrow (i). Let $ush(X) = 0$ and $X \subset Y$. By Lemma 6 there exist unique uniform shape morphisms $R : Y \rightarrow X$ and $I_X : X \rightarrow X$. Therefore $R \cdot J = I_X$, i.e. X is an absolute uniform shape retract. \square

Corollary 7. *Let $F : A \rightarrow Y$ be a uniform shape morphism from a uniform subspace A of a uniform space X into an absolute uniform shape retract Y . Then there exists a uniform shape morphism $F' : X \rightarrow Y$ such that $F' \cdot J = F$, where $J : A \rightarrow X$ is the inclusion uniform shape morphism induced by the inclusion uniform map $j : A \rightarrow X$.*

Proof. From Theorem 6 we obtain $ush(Y) = 0$. Then by Lemma 6 there are unique shape morphisms $F : A \rightarrow Y$ and $F' : X \rightarrow Y$. Consequently, the uniform shape morphism $F' \cdot J : A \rightarrow Y$ coincides with $F : A \rightarrow Y$. \square

Consider a uniform space X as a uniform subspace of an ARU -space Z . The set $\Lambda = \{U\}$ of all uniform neighborhoods of X in Z is directed by the order \leq defined by inclusion:

$$U \leq U' \Leftrightarrow U' \subseteq U.$$

Let $i_U : X \rightarrow U$ and $i_{UU'} : U' \rightarrow U$ be the inclusion uniform maps. The family $\mathbf{i} = ([i_U]) : X \rightarrow \mathbf{X} = (U, [i_{UU'}], \Lambda)$ forms a morphism of category $\mathbf{pro-H(ANRU)}$. In [11] it is proved that $\mathbf{i} : X \rightarrow \mathbf{X}$ is an $\mathbf{H(ANRU)}$ -expansion.

Theorem 8. *A uniform subspace X of an ARU -space Z has trivial uniform shape if and only if for every uniform neighborhood U of X in Z there exists a uniform neighborhood U' of X in Z such that $U' \subseteq U$ and the inclusion uniform map $i_{UU'} : U' \rightarrow U$ is semi-uniformly homotopic to a constant map.*

Proof. We can prove this as in S. Mardešić [7, Theorem 5], using Lemma 7. \square

Theorem 9. *For every finitistic uniform space X the following statements are equivalent:*

- (i) X admits an ARU -resolution.
- (ii) X has trivial uniform shape.
- (iii) X is an absolute uniform shape retract.

Proof. This is an immediate consequence of Theorems 1 and 5. \square

ACKNOWLEDGEMENT

The author expresses his gratitude to the referee for his helpful remarks.

REFERENCES

1. V. V. AGARONJAN, Shape classification of uniform spaces. (Russian) *Dokl. Akad. Nauk SSSR* **228**(1976), 1017–1020; translation in *Soviet Mat. Dokl.* **17**(1976), 848–851.
2. V. V. AGARONJAN and YU. M. SMIRNOV, The shape theory for uniform spaces and the shape uniform invariants. *Comment. Math. Univ. Carolin.* **19**(1978), No. 2, 351–357.
3. D. DOČINOV, On the uniform shape of metric spaces. (Russian) *Dokl. Akad. Nauk SSSR* **226**(1976), No. 2, 257–260; translation in *Soviet Mat. Dokl.* **17**(1976), 86–89.
4. J. R. ISBELL, Uniform spaces. *Mathematical Surveys*, No. 12. *American Mathematical Society, Providence, R.I.*, 1964.
5. A. KOYAMA, S. MARDEŠIĆ and T. WATANABE, Spaces which admit AR -resolutions. *Proc. Amer. Math. Soc.* **102**(1988), No. 3, 749–752.
6. V. I. KUZMINOV and I. A. ŠVEDOV, Cohomology groups of uniform spaces. *Sibirsk. Mat. Zh.* **5**(1964), 565–595.
7. S. MARDEŠIĆ, Retracts in shape theory. *Glas. Mat. Ser. III* **6(26)**(1971), 153–163.
8. S. MARDEŠIĆ, Aproximate polyhedra, resolutions of maps and shape fibrations. *Fund. Math.* **114**(1981), No. 1, 53–78.
9. S. MARDEŠIĆ, Strong shape and homology. *Springer Monographs in Mathematics. Springer-Verlag, Berlin*, 2000.

10. S. MARDEŠIĆ and J. SEGAL, Shape theory. *The inverse system approach. North-Holland Mathematical Library, 26. North-Holland Publishing Co., Amsterdam-New York, 1982.*
11. T. MIYATA, Uniform shape theory. *Glas. Mat. Ser. III* **29(49)**(1994), No. 1, 123–168.
12. NGUEN ANH KIET, Uniform fundamental classification of complete metric spaces and uniformly continuous mappings. *Bull. Acad. Pol. Sci. Ser. Sci. Mat. Astronom. Phys.* **23**(1975), 55–59.
13. NGUEN TO NHU, Shape of metric space in the category of metric spaces and uniformly continuous maps. *Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys.* **27**(1979), 929–934.
14. J. SEGAL, S. SPIEZ and B. GÜNTHER, Strong shape of uniform spaces. *Topology Appl.* **49**(1993), No. 3, 237–249.

(Received 21.06.2002)

Author's addresses:

Department of Mechanics and Mathematics
I. Javakhishvili Tbilisi State University
2, University St., Tbilisi 0143
Georgia

Department of Physics and Mathematics
Batumi State University
35, Ninoshvili St., Batumi 6016
Georgia
E-mail: vbaladze@viam.hepi.edu.ge