

ON TWO-POINT BOUNDARY VALUE PROBLEMS FOR TWO-DIMENSIONAL DIFFERENTIAL SYSTEMS WITH SINGULARITIES

S. MUKHIGULASHVILI

Abstract. For a differential system

$$\frac{du_1}{dt} = h_0(t, u_1, u_2)u_2, \quad \frac{du_2}{dt} = -h_1(t, u_1, u_2)u_1^{-\lambda} - h_2(t, u_1, u_2),$$

where $\lambda \in]0, 1[$ and $h_i :]a, b[\times]0, +\infty[\times \mathbb{R} \rightarrow [0, +\infty[$ ($i = 0, 1, 2$) are continuous functions, we have established sufficient conditions for the existence of at least one solution satisfying one of the two boundary conditions

$$\lim_{t \rightarrow a} u_1(t) = 0, \quad \lim_{t \rightarrow b} u_1(t) = 0$$

and

$$\lim_{t \rightarrow a} u_1(t) = 0, \quad \lim_{t \rightarrow b} u_2(t) = 0.$$

2000 Mathematics Subject Classification: 34B16.

Key words and phrases: Two-dimensional nonlinear differential system with singularities, two-point boundary value problem, existence theorem.

Two-point boundary value problems for second order ordinary differential equations with singularities with respect to one of the phase variable frequently occur in applications (see, for example, [1, 3, 4, 10, 11, 12]) and are investigated with sufficient thoroughness in the works of S. Taliafero [13], J. E. Bouillet and S. M. Gomes [2], Yu. A. Klovov and A. I. Lomakina [7], A. G. Lomtadze [8, 9], I. Kiguradze and B. Shekhter [6] and others. However analogous problems for two-dimensional singular differential systems still remain little studied. In this paper we investigate the first and second boundary value problems for nonlinear differential system with singularities with respect to both independent and/or one of the phase variables.

We consider the differential system

$$\frac{du_1}{dt} = h_0(t, u_1, u_2)u_2, \quad \frac{du_2}{dt} = -h_1(t, u_1, u_2)u_1^{-\lambda} - h_2(t, u_1, u_2), \quad (1)$$

with the boundary conditions

$$\lim_{t \rightarrow a} u_1(t) = 0, \quad \lim_{t \rightarrow b} u_1(t) = 0 \quad (2)$$

or

$$\lim_{t \rightarrow a} u_1(t) = 0, \quad \lim_{t \rightarrow b} u_2(t) = 0. \quad (3)$$

Here $\lambda \in]0, 1[$, $h_i :]a, b[\times]0, +\infty[\times \mathbb{R} \rightarrow [0, +\infty[$ ($i = 0, 1, 2$) are continuous functions. Moreover, there exist a continuous function $h :]a, b[\rightarrow [0, +\infty[$ and

positive numbers l, l_0 such that $0 < \int_a^b h(t) dt < \infty$ and the inequalities

$$h(t) \leq h_0(t, x, y) \leq l h(t), \quad h_1(t, x, y) + h_2(t, x, y) \geq l_0 h(t) \tag{4}$$

hold on the set $]a, b[\times]0, +\infty[\times \mathbb{R}$.

Solutions of problems (1), (2) and (1), (3) are sought for in the class of continuously differentiable vector functions $(u_1, u_2) :]a, b[\rightarrow]0, +\infty[\times \mathbb{R}$.

We set

$$\begin{aligned} \delta_1(t) &= \int_a^t h(s) ds \int_t^b h(s) ds, \quad \delta_2(t) = \int_a^t h(s) ds, \\ \delta(t) &= l \delta_1(t) \left(\int_a^b h(s) ds \right)^{-1}. \end{aligned}$$

Theorem 1. *Let, along with (4) the conditions*

$$\delta_1(t) > 0 \quad \text{for } a < t < b \tag{5}$$

and

$$\begin{aligned} \limsup_{\rho \rightarrow +\infty} \left[(2ll_0^{-1})^\lambda \rho^{\lambda^2-1} \int_a^b \delta_1^{1-\lambda}(t) h_1^*(t, \rho) dt + \rho^{-1} \int_a^b \delta_1(t) h_2^*(t, \rho) dt \right] \\ < l^{-2} \int_a^b h(t) dt \end{aligned} \tag{6}$$

be satisfied, where

$$h_i^*(t, \rho) = \sup \{ h_i(t, x, y) : 0 < x < l, \delta(t)|y| < \rho \} \quad (i = 1, 2). \tag{7}$$

Then problem (1), (2) is solvable.

Theorem 2. *Let, along with (4), the conditions*

$$\delta_2(t) > 0 \quad \text{for } a < t \leq b$$

and

$$\limsup_{\rho \rightarrow +\infty} \left[\left(\frac{2}{l_0 \delta_2(b)} \right)^\lambda \rho^{\lambda^2-1} \int_a^b \delta_2^{1-\lambda}(t) h_1^*(t, \rho) dt + \rho^{-1} \int_a^b \delta_2(t) h_2^*(t, \rho) dt \right] < \frac{1}{l}$$

hold, where

$$h_i^*(t, \rho) = \sup \{ h_i(t, x, y) : 0 < x < \rho, \delta_2(t)|y| < \rho \} \quad (i = 1, 2).$$

Then problem (1), (3) is solvable.

As an example, consider the differential system

$$\begin{aligned} \frac{du_1}{dt} &= (t-a)^\alpha (b-t)^\beta \rho_0(t, u_1, u_2) u_2, \\ \frac{du_2}{dt} &= -\frac{p_1(t, u_1, u_2)}{(t-a)^{\alpha_1} (b-t)^{\beta_1}} u_1^{-\lambda} - \frac{p_2(t, u_1, u_2)}{(t-a)^{\alpha_2} (b-t)^{\beta_2}}, \end{aligned} \tag{8}$$

where $\lambda \in]0, 1[$, and

$$\alpha > -1, \quad \beta > -1, \quad \alpha_i \geq -\alpha, \quad \beta_i \geq -\beta \quad (i = 1, 2),$$

and $p_i :]a, b[\times]0, +\infty[\times \mathbb{R} \rightarrow]r_1, r_2[$ ($i = 0, 1, 2$) are continuous functions, $r_i = \text{const} > 0$ ($i = 1, 2$).

Theorems 1 and 2 give rise to

Corollary 1. *The inequalities*

$$\alpha_1 < (\alpha + 1)(1 - \lambda) + 1, \quad \beta_1 < (\beta + 1)(1 - \lambda) + 1, \quad \alpha_2 < \alpha + 2, \quad \beta_2 < \beta + 2$$

guarantee the solvability of problem (1), (2), and the inequalities

$$\alpha_1 < (\alpha + 1)(1 - \lambda) + 1, \quad \beta_1 < 1, \quad \alpha_2 < \alpha + 2, \quad \beta_2 < 1$$

guarantee the solvability of problem (1), (3).

The above example shows that Theorem 1 (Theorem 2) covers the case where the functions h_1 and h_2 have singularities of arbitrary order at $t = a$ and $t = b$ (at $t = a$).

To prove the formulated theorems, we need two lemmas from [5] on the representation of a solution of the differential system

$$\frac{dv_1}{dt} = p(t) v_2, \quad \frac{dv_2}{dt} = q(t) \tag{9}$$

under the boundary conditions

$$v_1(a) = 0, \quad v_1(b) = 0 \tag{10}$$

or

$$v_1(a) = 0, \quad v_2(b) = 0. \tag{11}$$

Lemma 1. *Let $p, q :]a, b[\rightarrow \mathbb{R}$ be continuous functions such that*

$$\int_a^b |p(t)| dt < +\infty, \quad \int_a^b \left(\int_a^t |p(\tau)| d\tau \int_t^b |p(\tau)| d\tau \right) |q(t)| dt < +\infty \quad (i = 1, 2),$$

$$\gamma_0 \stackrel{\text{def}}{=} \int_a^b p(t) dt \neq 0. \tag{12}$$

Then problem (9), (10) has a unique solution $v = (v_1, v_2)$ and the representations

$$v_i(t) = \int_a^b g_i(t, s) q(s) ds \quad (i = 1, 2) \tag{13}$$

are valid, where

$$g_1(t, s) = \begin{cases} -\frac{1}{\gamma_0} \int_a^s p(\tau) d\tau \int_t^b p(\tau) d\tau & \text{for } s \leq t, \\ -\frac{1}{\gamma_0} \int_a^t p(\tau) d\tau \int_s^b p(\tau) d\tau & \text{for } s > t, \end{cases}$$

$$g_2(t, s) = \begin{cases} \frac{1}{\gamma_0} \int_a^s p(\tau) d\tau & \text{for } s \leq t, \\ -\frac{1}{\gamma_0} \int_s^b p(\tau) d\tau & \text{for } s > t. \end{cases}$$

Lemma 2. Let $p, q \in]a, b[\rightarrow \mathbb{R}$ be continuous functions such that conditions (12) and

$$\int_a^b p(t) dt < +\infty, \quad \int_a^b \left(\int_a^t |p(\tau)| d\tau \right) q(t) dt < +\infty$$

are fulfilled. Then problem (9), (11) has a unique solution $v = (v_1, v_2)$ and representations (13) are valid, where

$$g_1(t, s) = \begin{cases} - \int_a^s p(\tau) d\tau & \text{for } s \leq t, \\ - \int_a^t p(\tau) d\tau & \text{for } s > t, \end{cases} \quad g_2(t, s) = \begin{cases} 0 & \text{for } s \leq t, \\ -1 & \text{for } s > t. \end{cases}$$

Proof of Theorem 1. By virtue of condition (6), the constant ρ can be chosen so that the inequalities

$$\rho > 1 + \frac{l_0}{2l} \left(\int_a^b h(s) ds \right)^2, \tag{14}$$

$$\left(\frac{2l}{l_0} \right)^\lambda \rho^{\lambda^2} \int_a^b \delta_1^{1-\lambda}(t) h_1^*(t, \rho) dt + \int_a^b \delta_1(t) h_1^*(t, \rho) dt < \frac{\rho}{l^2} \int_a^b h(t) dt \tag{15}$$

be fulfilled.

Let \mathbb{B} be a Banach space of two-dimensional continuous vector functions $v = (v_1, v_2) : [a, b] \rightarrow \mathbb{R}^2$ with the norm

$$\|v\| = \max \{ |v_1(t)| + |v_2(t)| : a \leq t \leq b \}$$

and \mathbb{B}_ρ be a set of all $v = (v_1, v_2) \in \mathbb{B}$ satisfying the conditions

$$\frac{l_0}{2l} \rho^{-\lambda} \delta_1(t) \leq v_1(t) < \rho, \quad |v_2(t)| \leq \rho \quad \text{for } a \leq t \leq b.$$

In view of inequality (14), \mathbb{B}_ρ is a nonempty closed convex subset of the space \mathbb{B} .

For arbitrary $v = (v_1, v_2) \in \mathbb{B}_\rho$ we set

$$\gamma(v_1, v_2) = \int_a^b h_0 \left(\tau, v_1(\tau), \frac{v_2(\tau)}{\sigma(\tau)} \right) d\tau,$$

$$q(v_1, v_2)(t) = -h_1 \left(t, v_1(t), \frac{v_2(t)}{\sigma(t)} \right) v_1^{-\lambda}(t) - h_2 \left(t, v_1(t), \frac{v_2(t)}{\sigma(t)} \right),$$

$$g_1(v_1, v_2)(t, s) \cdot \gamma(v_1, v_2) = \begin{cases} - \int_a^s h_0 \left(\tau, v_1(\tau), \frac{v_2(\tau)}{\sigma(\tau)} \right) d\tau \\ \quad \times \int_t^b h_0 \left(\tau, v_1(\tau), \frac{v_2(\tau)}{\sigma(\tau)} \right) d\tau & \text{for } s \leq t, \\ - \int_a^t h_0 \left(\tau, v_1(\tau), \frac{v_2(\tau)}{\sigma(\tau)} \right) d\tau \\ \quad \times \int_s^b h_0 \left(\tau, v_1(\tau), \frac{v_2(\tau)}{\sigma(\tau)} \right) d\tau & \text{for } s > t, \end{cases}$$

$$g_2(v_1, v_2)(t, s) \cdot \gamma(v_1, v_2) = \begin{cases} \int_a^s h_0\left(\tau, v_1(\tau), \frac{v_2(\tau)}{\sigma(\tau)}\right) d\tau & \text{for } s \leq t, \\ -\int_s^b h_0\left(\tau, v_1(\tau), \frac{v_2(\tau)}{\sigma(\tau)}\right) d\tau & \text{for } s > t. \end{cases}$$

Let us now define the continuous operators $W_i : \mathbb{B} \rightarrow C([a, b])$ ($i = 1, 2$) and $W : \mathbb{B} \rightarrow C([a, b]) \times C([a, b])$ by the equalities

$$W_i(v_1, v_2)(t) = (\sigma(t))^{i-1} \int_a^b g_i(v_1, v_2)(t, s) q(v_1, v_2)(s) ds \quad (i = 1, 2)$$

and

$$W(v_1, v_2)(t) = (W_1(v_1, v_2)(t), W_2(v_1, v_2)(t)).$$

Then if problem (1), (2) has a solution $u = (u_1, u_2)$ satisfying the conditions

$$\frac{l_0}{2l} \rho^{-\lambda} \delta_1(t) \leq u_1(t) \leq \rho, \quad \sigma(t)|u_2(t)| \leq \rho \quad \text{for } a \leq t \leq b, \quad (16)$$

$$\lim_{t \rightarrow a} \sigma_1(t) u_2(t) = 0, \quad \lim_{t \rightarrow b} \sigma_1(t) u_2(t) = 0, \quad (17)$$

the vector function $v = (v_1, v_2)$ with the components

$$v_1(t) = u_1(t), \quad v_2(t) = \sigma(t) u_2(t) \quad (18)$$

belongs to the set \mathbb{B}_ρ and, by virtue of Lemma 1, is a solution of the operator equation

$$v(t) = W(v)(t). \quad (19)$$

Conversely, if equation (19) has a solution $v = (v_1, v_2) \in \mathbb{B}_\rho$, then, again by Lemma 1, the function $u = (u_1, u_2)$, the components of which are defined from equalities (18), is a solution of problem (1), (2) satisfying conditions (17). Thus to prove the theorem it is sufficient to establish that the operator W has a fixed point on the set \mathbb{B}_ρ .

In the first place, we will show that

$$W(\mathbb{B}_\rho) \subset \mathbb{B}_\rho. \quad (20)$$

Let $(v_1, v_2) \in \mathbb{B}_\rho$. Then by virtue of (7)

$$|q(v_1, v_2)(t)| \leq \varphi(t) \quad \text{for } a < t < b, \\ \varphi(t) = \left(\frac{2l}{l_0}\right)^\lambda \rho^{\lambda^2} \delta_1^{-\lambda}(t) h_1^*(t, \rho) + h_2^*(t, \rho),$$

and, also, by virtue of the second of equalities (4)

$$|q(v_1, v_2)(t)| \geq \rho^{-\lambda} l_0 h(t) \quad \text{for } a < t < b.$$

Taking into account the latter estimates and the first of equalities (4), we obtain

$$W_1(v_1, v_2)(t) \geq \frac{l_0}{l} \rho^{-\lambda} \left(\int_a^b h(\tau) d\tau\right)^{-1} \left[\int_a^b h(\tau) d\tau \int_a^t \left(\int_a^s h(\tau) d\tau\right) h(s) ds \right. \\ \left. + \int_a^t h(\tau) d\tau \int_t^b \left(\int_s^b h(\tau) d\tau\right) h(s) ds \right]$$

$$= \frac{l_0}{2l} \rho^{-\lambda} \delta_1(t) \text{ for } a \leq t \leq b, \quad (21)$$

$$\begin{aligned} W_1(v_1, v_2)(t) &\leq l^2 \left(\int_a^b h(\tau) d\tau \right)^{-1} \left[\int_t^b h(\tau) d\tau \int_a^t \left(\int_a^s h(\tau) d\tau \right) \varphi(s) ds \right. \\ &\quad \left. + \int_a^t h(\tau) d\tau \int_t^b \left(\int_t^b h(\tau) d\tau \right) \varphi(s) ds \right] \text{ for } a \leq t \leq b \end{aligned} \quad (22)$$

and

$$\begin{aligned} W_2(v_1, v_2)(t) &\leq l^2 \left(\int_a^b h(\tau) d\tau \right)^{-2} \delta_1(t) \left[\int_a^t \left(\int_a^s h(\tau) d\tau \right) \varphi(s) ds \right. \\ &\quad \left. + \int_t^b \left(\int_s^b h(\tau) d\tau \right) \varphi(s) ds \right] \text{ for } a \leq t \leq b. \end{aligned} \quad (23)$$

From equalities (21), (23) with (15) taken into account we obtained

$$\begin{aligned} W_i(v_1, v_2)(t) &\leq l^2 \left(\int_a^b h(\tau) d\tau \right)^{-1} \int_a^b \delta_1(\tau) \varphi(\tau) d\tau < \rho \quad (i = 1, 2) \\ &\text{for } a \leq t \leq b. \end{aligned} \quad (24)$$

This and (21) imply that inclusion (20) is valid.

Now note that from the definition of the operators W_i ($i = 1, 2$) and Lemma 1 it follows that the functions $W_i(v_1, v_2)(t)$ ($i = 1, 2$) are continuously differentiable on the interval $]a, b[$ for any $v = (v_1, v_2) \in \mathbb{B}_\rho$ and

$$\begin{aligned} \frac{d}{dt} W_1(v_1, v_2)(t) &= h_0 \left(t, v_1, \frac{v_2}{\sigma} \right) \delta^{-1}(t) W_2(v_1, v_2)(t), \\ \frac{d}{dt} W_2(v_1, v_2)(t) &= - \left(h_1 \left(t, v_1, \frac{v_2}{\sigma} \right) W_1^{-\lambda}(v_1, v_2)(t) + h_2(t, v_1, v_2) \right) \delta(t) \\ &\quad + \frac{\delta'(t)}{\delta(t)} W_2(v_1, v_2)(t). \end{aligned}$$

From this, taking into account estimates (21) and (24), we obtain

$$\left| \frac{d}{dt} W_i(v_1, v_2) \right| \leq \eta_i(t) \quad (i = 1, 2) \quad \text{for } a < t < b, \quad (25)$$

where $\eta_i :]a, b[\rightarrow [0, +\infty[$ are continuous functions defined by the equalities

$$\eta_1(t) = \rho \frac{h(t)}{\delta(t)}, \quad \eta_2(t) = \delta(t) \varphi(t) + \rho \frac{h(t)}{\delta_1(t)} \int_a^b h(\tau) d\tau.$$

Analogously, from estimates (22) and (23) we obtain

$$|W_1(v_1, v_2)(t)| + |W_2(v_1, v_2)(t)| \leq \varepsilon_0(t) \quad \text{for } a \leq t \leq b, \quad (26)$$

where the continuous function $\varepsilon_0 : [a, b] \rightarrow [0, +\infty[$ is defined by the equality

$$\begin{aligned} \varepsilon_0(t) &= 2l^2 \left(\int_a^b h(\tau) d\tau \right)^{-1} \left[\int_t^b h(\tau) d\tau \int_a^t \left(\int_a^s h(\tau) d\tau \right) \varphi(\tau) d\tau \right. \\ &\quad \left. + \int_a^t h(\tau) d\tau \int_t^b \left(\int_s^b h(\tau) d\tau \right) \varphi(\tau) d\tau \right], \end{aligned}$$

and therefore

$$\varepsilon_0(a) = 0, \quad \varepsilon_0(b) = 0. \quad (27)$$

From (25)–(27) we conclude that the set of functions $W(\mathbb{B}_\rho)$ is equicontinuous and uniformly bounded, i.e., the continuous operator W transforms the bounded convex set \mathbb{B}_ρ into its compact subset and, by virtue of the Schauder's principle of fixed point, equation (19) has at least one solution. \square

Theorem 2 is proved in an analogous manner with the only difference that we use Lemma 2.

ACKNOWLEDGEMENT

The work was supported by a fellowship from the INTAS YS 2001-2/80 as for a visiting professor at the University of Udine, Italy.

REFERENCES

1. J. A. ACKROYD, On the laminar compressible boundary layer with stationary origin on a moving flat wall. *Proc. Cambridge Phil. Soc.* **63**(1967), 871–888.
2. J. E. BOUILLET and S. M. GOMES, An equation with a singular nonlinearity related to diffusion problems in one dimension. *Quart. Appl. Math.* **42**(1985), No. 4, 395–402.
3. A. J. CALLEGARY and M. B. FRIEDMAN, An analytical solution of a nonlinear singular boundary value problem in the theory of viscous fluids. *J. Math. Anal. Appl.* **21**(1968), No. 3, 510–529.
4. A. J. CALLEGARY and A. NACHMAN, Some singular, nonlinear equations arising in boundary-layer theory. *J. Math. Anal. Appl.* **64**(1978), No. 1, 96–105.
5. I. T. KIGURADZE, On boundary value problems for linear differential systems with singularities. (Russian) *Differentsial'nye Uravneniya* **39**(2002), No. 2, 198–209.
6. I. T. KIGURADZE and B. L. SHEKHTER, Singular boundary value problems for second-order ordinary differential equations. (Russian) *Itogi Nauki i Tekhniki, Current problems in mathematics. Newest results, Vol. 30 (Russian)*, 105–201, *Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow*, 1987; English translation: *J. Sov. Math.* **43**(1988), No. 2, 2340–2417.
7. YU. A. KLOKOV and A. I. LOMAKINA, A certain boundary value problem with singularities at the endpoints of the interval. (Russian) *Latvian mathematical yearbook, 17 (Russian)*, 179–186. *Izdat. "Zinatne", Riga*, 1976.
8. A. G. LOMTATIDZE, Solvability of boundary value problems for second-order nonlinear ordinary differential equations with singularities. (Russian) *Reports of the extended sessions of a seminar of the I. N. Vekua Institute of Applied Mathematics, Vol. I, No. 3 (Russian) (Tbilisi)*, 1985, 85–92, 170–171, *Tbilis. Gos. Univ., Tbilisi*, 1985.
9. A. G. LOMTATIDZE, On positive solutions of singular boundary value problems for nonlinear second order ordinary differential equations. (Russian) *Differentsial'nye Uravneniya* **22**(1986), No. 6, p. 1092.
10. N. F. MOROZOV, On the analytic structure of the solution of the membrane equation. (Russian) *Dokl. Akad. Nauk SSSR* **152**(1963), No. 1, 78–80.
11. N. P. MOROZOV and L. S. SRUBSHCHIK, Application of Chaplygin's method to the study of the membrane equation. (Russian) *Differentsial'nye Uravneniya* **2**(1966), No. 3, 425–427.

12. L. S. SRUBSHCHIK and V. I. YUDOVICH, The asymptotics of the equation for large bending of a round symmetrically loaded lamina. (Russian) *Sibirsk. Mat. Zh.* **4**(1963), No. 3, 657–672.
13. S. TALIAFERRO, A nonlinear singular boundary value problem. *Nonlinear Anal.* **3**(1979), No. 6, 897–904.

(Received 1.05.200)

Author's address:

A. Razmadze Mathematical Institute
Georgian Academy of Sciences
1, M. Aleksidze St., Tbilisi 0193
Georgia
E-mail: mukhig@rmi.acnet.ge