

**THE WEIGHTED BMO CONDITION AND A
CONSTRUCTIVE DESCRIPTION OF CLASSES OF
ANALYTIC FUNCTIONS SATISFYING THIS CONDITION**

L. CHIKVINIDZE

ABSTRACT. The problem of local polynomial approximation of analytic functions prescribed in finite domains with a quasiconformal boundary is investigated in weighted plane integral metrics; a constructive description of the class of analytic functions satisfying a weak version of the known BMO condition is obtained.

The First results of the investigation of the problem (formulated by V. I. Belyi) dealing with a local polynomial approximation of analytic functions prescribed in finite domains with quasiconformal boundary have been described in [1, 2] for weighted plane integral metrics. This problem is investigated in [3] for the nonweighted case, where a constructive description of Hölder classes as well as of some other classes of analytic functions has been obtained. In the present paper we continue the investigation of the above-mentioned problem for the weighted case; moreover, a constructive description of one more class of analytic functions in weighted plane integral metrics is obtained.

1. NOTATION AND DEFINITIONS. THE BASIC RESULTS

Let G be the domain with a quasiconformal boundary $\partial G = \Gamma$, and let $y = y(\zeta)-$ be a quasiconformal reflection across the curve Γ [4]. We will be concerned only with the special, so-called canonical, quasiconformal reflection (see relations (2.1) and (2.2)). Let w be some weight function (i.e., nonnegative and measurable) defined in the domain Γ . Let us introduce the

1991 *Mathematics Subject Classification.* 30E10, 30C62, 41A10.

Key words and phrases. Domain with quasiconformal boundary, weighted local polynomial approximation, weak versions of the Muckenhoupt and the BMO conditions.

notation

$$H'(G) = \{f : f \text{ holomorphic in } G\},$$

$$L_p(G, w) = \{f : |f|^p w \in L_1(G)\}, \quad H'_p(G, w) = L_p(G, w) \cap H'(G) \quad (p \geq 1).$$

Furthermore, let σ be the plane Lebesgue measure, and let μ be the Borel measure defined by the equality

$$\mu(E) = \iint_E w(z) d\sigma_z \quad (E \subset G). \quad (1.1)$$

The integral with respect to the measure μ of the function f will be denoted by the symbols

$$\iint_E f(z) d\mu_z = \iint_E f d\mu.$$

If $\mu = \sigma$, then (in cases where this does not cause misunderstanding) we shall use the brief notation

$$\iint_E f(z) d\sigma_z = \iint_E f.$$

Let $Q = Q(z, r)$ be an open square with center at the point z , whose sides are parallel to the coordinate axes and have the length r , and let

$$F(G) = \{Q = Q(z, r) : z \in \Gamma, \quad r > 0\}.$$

For a square Q denote $|Q| = \sigma(Q)$.

Let the function f and the weight function w be defined in the domain G , let the measure μ be defined by equality (1.1), and

$$f_{\mu, Q \cap G} = \frac{1}{\mu(Q \cap G)} \iint_{Q \cap G} f d\mu. \quad (1.2)$$

We say that f satisfies the weighted BMO condition $\text{BMO}_p(G, w)$ (briefly, $f \in \text{BMO}_p(G, w)$), if

$$\sup_{Q \in F(G)} \left(\frac{1}{\mu(Q \cap G)} \iint_{Q \cap G} |f - f_{\mu, Q \cap G}|^p d\mu \right)^{\frac{1}{p}} \stackrel{\text{df}}{=} \|f\|_{\text{BMO}_p(G, w)} < \infty. \quad (1.3)$$

When $p = 1$ and $w(z) = 1$ everywhere in G , we shall use the usual notation $f \in \text{BMO}(G)$ and $\|f\|_{\text{BMO}(G)}$ respectively.

The $\text{BMO}(G)$ condition is a weaker analogue of the well-known BMO (bounded mean oscillation) condition (see, e.g., [5, Ch. VI]).

Next, we say that the weight function w given in the domain G satisfies the condition $A_p(F(G))$ ($1 < p < \infty$) (briefly $w \in A_p(F(G))$), if (assuming $0 \cdot \infty = 0$)

$$\sup_{Q \in F(G)} \left(\frac{1}{|Q \cap G|} \iint_{Q \cap G} w \right) \left(\frac{1}{|Q \cap G|} \iint_{Q \cap G} w^{-\frac{1}{p-1}} \right)^{p-1} < \infty.$$

The condition $A_p(F(G))$, introduced for the first time in [6] (with the unit circle as G), is a weaker analogue of the well-known Muckenhoupt condition (A_p) [7].

Let $z_0 \in \Gamma$, $\rho_n(z_0)$ ($n \in N$) be the distance from the point z_0 to the external level line $\Gamma_{1+1/n}$ of G , $u(z_0, r) = \{z : |z - z_0| < r\}$, $c_0 > 0$. Set

$$G(z_0, c_0) = \{z \in G : |\zeta - z| \geq c_0|\zeta - z_0| \quad \forall \zeta \in CG\},$$

$$G_n(z_0, c_0) = G(z_0, c_0) \cup \{u(z_0, \rho_n(z_0)) \cap G\}.$$

The set $G(z_0, c_0)$ is a kind of a “nontangential” subset of G with the vertex at the point z_0 .

In the sequel, for brevity we shall write

$$\left(\iint_{G_n(z_0, c_0)} |f(z) - P_n(z)|^p w(z) d\sigma_z \right)^{\frac{1}{p}} \stackrel{df}{=} \|f - P_n\|_{z_0, p, w}.$$

Let us now formulate the basic results in which G denotes a finite domain with a quasiconformal boundary Γ , the weight function $w \in A_p(F(G))$ ($1 < p < \infty$), and μ is the measure defined by equality (1.1).

Theorem 1. *For the function f to belong to the class $BMO_p(G, w) \cap H'(G)$ (neglecting its values on the set of measure zero), it is necessary and sufficient that a sequence of algebraic polynomials P_n of order not higher than n exist such that for all $z_0 \in \Gamma$ and $n \in N$ the relation*

$$\|f - P_n\|_{z_0, p, w} \leq c(c_0) \left(\mu\{u(z_0, \rho_n(z_0)) \cap G\} \right)^{\frac{1}{p}} \tag{1.4}$$

holds, where the constant $c(c_0)$ does not depend on z_0 and n .

Theorem 2. *Let $f \in H'(G)$. The following conditions are equivalent:*

- (a) $f \in BMO(G)$;
- (b) $f \in BMO_p(G, w)$.

This theorem is an analogue of the well-known John and Nirenberg theorem (see, e.g., [5]).

Obviously, Theorem 2 allows us to formulate Theorem 1 as follows:

Theorem 1*. *Theorem 1 remains valid when the class $BMO_p(G)$ is replaced by the class $BMO(G)$.*

Thus we have given the constructive description of the class of functions $BMO(G) \cap H'(G)$ in the weighted plane integral metrics.

2. AUXILIARY RESULTS

Let G be the domain with a quasiconformal boundary Γ , and let $0 \in G$, $y = y(\zeta)$ be a quasiconformal reflection across the curve Γ [4]. As follows from the Ahlfors theorem [4] (see also [8]), the reflection $y = y(\zeta)$ can always be chosen to be canonical in the sense that it is differentiable for $\zeta \notin \Gamma$, and for any fixed sufficiently small $\delta > 0$ it will satisfy the relations

$$|y_{\bar{\zeta}}(\zeta)| \asymp M, \quad |y_{\zeta}(\zeta)| \asymp M, \quad \delta < |\zeta| < 1/\delta, \quad \zeta \notin \Gamma, \quad (2.1)$$

$$|y_{\bar{\zeta}}(\zeta)| \asymp M|\zeta|^{-2}, \quad |y_{\zeta}(\zeta)| \asymp M|\zeta|^{-1}, \quad |\zeta| \leq \delta, \quad |\zeta| \geq \frac{1}{\delta}, \quad (2.2)$$

where $M = M(\delta, \Gamma)$ is a constant depending only on δ and Γ .

The symbol $A \asymp B$ for the numbers A and B depending on some parameters denotes that $A \leq cB$, where $c = \text{const} > 0$ does not depend on those parameters; the symbol $A \approx B$ means that $B \asymp A$; $A \asymp B$ if simultaneously $A \asymp B$ and $A \approx B$.

Let $w \in A_p(F(G))$ ($1 < p < \infty$), and let $y = y(\zeta)$ be a canonical quasiconformal reflection across the curve $\Gamma = \partial G$. Let us introduce the notation

$$w^*(z) = \begin{cases} w(z) & \text{for } z \in G, \\ w(y(z)) & \text{for } z \notin G, \end{cases} \quad \mu^*(E) = \iint_E w^* d\sigma. \quad (2.3)$$

It is clear that if $E \subset G$, then $\mu(E) = \mu^*(E)$. Suppose further that

$$\begin{aligned} \rho(E_1, E_2) &= \inf \{|z_1 - z_2| : z_1 \in E_1, z_2 \in E_2\}, \\ \text{diam } E &= \sup \{|z_1 - z_2| : z_1, z_2 \in E\}, \\ F(G, k) &= \{Q : \text{diam } Q \geq k\rho(Q, \Gamma), Q \cap G \neq \emptyset\} \quad (k > 0). \end{aligned} \quad (2.4)$$

Let $w \in A_p(F(G))$ ($1 < p < \infty$), and let $w^*(z)$ be the function defined by equality (2.3). Then owing to relations (2.1) and (2.2), we can conclude that for all $Q \in F(G, k)$, $\text{diam } Q < k_0$ ($k_0, k > 0$ are arbitrary fixed numbers) the inequality

$$\left(\frac{1}{|Q|} \iint_Q w^* \right) \left(\frac{1}{|Q|} \iint_Q w^{*-\frac{1}{p-1}} \right)^{p-1} \leq c(k, k_0) < \infty \quad (2.5)$$

holds, where $c(k, k_0)$ is a constant independent of Q .

Lemma 1 ([9], [10]). *Let $w \in A_p(F(G))$, and let $w^*(z)$ be the function defined by equality (2.3). There exist numbers $0 < \delta = \delta(k, k_0) < 1$, $0 < \varepsilon = \varepsilon(k, k_0) < 1$ such that for every $e \subset Q$ the inequality $|e| < \delta|Q|$ implies*

$$\iint_e w^* < \varepsilon \iint_Q w^*$$

for all $Q \in F(G, k)$, $\text{diam } Q < k_0$.

Next, by virtue of the Hölder inequality and relation (2.5), we find that for all $Q(z, r), Q(z, R) \in F(G, k)$ ($0 < r \leq R \leq k_0 < \infty$) the inequality

$$\left(\frac{r}{R}\right)^2 \leq c(k, k_0) \frac{\mu^*(Q(z, r))}{\mu^*(Q(z, R))} \tag{2.6}$$

holds, where μ^* is defined by equality (2.3).

In particular, it follows from (2.6) that μ^* satisfies the known “doubling” condition

$$\mu^*(2Q) \leq c \mu^*(Q), \quad (Q \in F(G, k), \text{diam } Q < k_0). \tag{2.7}$$

Let us now prove that for all squares $Q \in F(G)$ ($\text{diam } Q < k_0$) the relation

$$\mu(Q \cap G) \geq \mu^*(Q) \tag{2.8}$$

holds, where μ and μ^* are defined respectively by equalities (1.1) and (2.3).

Indeed, let $Q = Q(z_0, r)$ ($z_0 \in \Gamma, r > 0$), $\text{diam } Q < k_0$. Owing to relations (2.1) and (2.2), we get

$$\mu^*(Q(z_0, (1/M)r) \cap CG) \leq c(M) \mu^*(Q(z_0, r) \cap G) = c(M) \mu(Q \cap G),$$

where $M > 1$ is the constant from (2.1) and (2.2), and CG is the complement to the domain G . But then, using the “doubling” condition (2.7), we get

$$\begin{aligned} \mu^*(Q) &\leq \mu^*(Q(z_0, (1/M)r) \cap G) + \mu^*(Q(z_0, (1/M)r) \cap CG) \leq \\ &\leq c(M) \mu^*(Q(z_0, r) \cap G) \leq \mu(Q \cap G). \quad \square \end{aligned}$$

Let w be a weight function, and let μ^* be the measure defined by equality (2.3). Let f be a function given in the domain G , and let Q be a square. Introduce the notation

$$f^*(z) = \begin{cases} f(z) & \text{for } z \in G, \\ f(y(z)) & \text{for } z \notin G, \end{cases} \quad f_{\mu^*, Q}^* = \frac{1}{\mu^*(Q)} \iint_Q f^* d\mu^*. \tag{2.9}$$

In the case of the Lebesgue measure σ , we shall use f_Q^* instead of $f_{\sigma^*, Q}^*$.

Lemma 2. *Let w be some weight function, $p > 1$, $f \in \text{BMO}_p(G, w)$. Then for all squares $Q \in F(G, k)$, $\text{diam } Q < k_0$ ($k, k_0 > 0$ are fixed numbers) the relation*

$$\left(\frac{1}{\mu^*(Q)} \iint_Q |f^* - f_{\mu^*, Q}^*|^p d\mu^* \right)^{\frac{1}{p}} \leq c^* \|f\|_{\text{BMO}_p(G, w)} \quad (2.10)$$

holds, where c^* is a constant independent of Q , p , f , and w .

Proof. Assume first that $Q \in F(G)$, $\text{diam } Q < k_0$. Let $M > 1$ be the number from relations (2.1) and (2.2), and let MQ be the square obtained by an M -fold increase of the square Q . It follows from the ‘‘doubling’’ condition (2.7) that

$$\mu^*(Q) \geq c(M)\mu^*(MQ) \geq c(M)\mu(MQ \cap G). \quad (2.11)$$

On account of relations (2.1) and (2.2) we obtain

$$\left(\iint_{Q \cap CG} |f^* - f_{\mu, MQ}|^p d\mu^* \right)^{\frac{1}{p}} \leq M^2 \left(\iint_{MQ \cap G} |f - f_{\mu, MQ}|^p d\mu \right)^{\frac{1}{p}}. \quad (2.12)$$

Hence, using the Minkowsky inequality and relations (2.11), (2.12), and (1.3), we obtain

$$\begin{aligned} \left(\frac{1}{\mu^*(Q)} \iint_Q |f^* - f_{\mu^*, Q}^*|^p d\mu^* \right)^{\frac{1}{p}} &\leq \left(\frac{2}{\mu^*(Q)} \iint_Q |f^* - f_{\mu, MQ \cap G}|^p d\mu^* \right)^{\frac{1}{p}} \leq \\ &\leq \left(\frac{M^2 + 1}{c(M)} \frac{1}{\mu(MQ \cap G)} \iint_{MQ \cap G} |f - f_{\mu, MQ \cap G}|^p d\mu \right)^{\frac{1}{p}} \leq c^* \|f\|_{\text{BMO}_p(G, w)}. \end{aligned}$$

Thus we have proved that inequality (2.10) is true for all $Q \in F(G)$, $\text{diam } Q < k_0$. Using the ‘‘doubling’’ condition (2.7), it is not difficult to show that (2.10) holds for all $Q \in F(G, k)$ as well. \square

Lemma 3. *Let $f \in \text{BMO}(G)$ be an analytic function in the domain G , $\zeta \in G$, $Q = Q(\zeta, a) \notin F(G, k)$ ($k \leq 1$), $Q \subset G$. Then*

$$|f(\zeta) - f(z)| \leq 4\|f\|_{\text{BMO}(G)} \quad \forall (z \in Q(\zeta, a)).$$

Proof. Since $Q = Q(\zeta, a) \notin F(G, k)$ ($k \leq 1$), it follows from the definition of the set $F(G, k)$ (see (2.4)) that $\text{diam } Q < k\rho(Q, \Gamma)$ ($k \leq 1$). Hence

$$|\zeta - z| \leq \frac{1}{2} \text{diam } Q < \frac{1}{2} k\rho(Q, \Gamma) < \frac{1}{2} k\rho(\zeta, \Gamma) < \frac{1}{2} \rho(\zeta, \Gamma)$$

for all $z \in Q$. Then assuming for brevity that $\rho(\zeta, \Gamma) = \rho$, owing to the mean value theorem and the condition $f \in \text{BMO}(G)$, we obtain

$$\begin{aligned} |f(z) - f(\zeta)| &\leq \frac{1}{|u(z, \rho/2)|} \iint_{u(z, \rho/2)} |f(\xi) - f(\zeta)| d\sigma_\xi \leq \\ &\leq \frac{4}{|u(\zeta, \rho)|} \iint_{u(\zeta, \rho)} |f(\xi) - f_{u(\zeta, \rho)}| d\sigma_\xi \leq 4\|f\|_{\text{BMO}(G)}. \end{aligned}$$

for all $z \in Q(\zeta, a)$. \square

Lemma 4. *Let $f \in \text{BMO}(G)$ be an analytic function in the domain G , and let f^* and f_Q^* be defined by equalities (2.9) (the case $\mu = \sigma$), $Q \in F(G, k)$ ($k \leq 1$), $\alpha > c^*\|f\|_{\text{BMO}(G)}$ (c^* is a constant from (2.10)). Then there exists at most a countable set of nonintersecting squares $A = \{Q_j\}$ such that $Q_j \in F(G, k)$, and*

$$(1) |f^*(z) - f_Q^*| \leq 12\alpha \quad \forall (z \in Q \setminus Q_j, Q_j \in A); \tag{2.13}$$

$$(2) \alpha \leq \frac{1}{|Q_j|} \iint_{Q_j} |f^*(z) - f_Q^*| d\sigma_z < 4\alpha \quad \forall (Q_j \in A); \tag{2.14}$$

$$(3) \sum_{Q_j \in A} |Q_j| \leq \frac{1}{\alpha} \|f\|_{\text{BMO}(G)} |Q|. \tag{2.15}$$

Proof. It is obvious that the conditions (2.10) and $\alpha > c^*\|f\|_{\text{BMO}(G)}$ yield

$$\frac{1}{|Q|} \iint_Q |f^*(z) - f_Q^*| d\sigma_z < \alpha.$$

Let Q^* be the square obtained by partitioning Q into four equal squares.

In the case $Q^* \in F(G, k)$ we insert Q^* in A if $\iint_{Q^*} |f(z) - f_Q^*| d\sigma_z \geq \alpha|Q^*|$, while when the opposite inequality holds we again partition Q^* in four equal squares and argue as above.

Let us show that the squares $Q_j \in A$ obtained in such a way satisfy all the requirements of Lemma 2.

Let $\zeta \in \{Q \cap G\} \setminus \cup \{Q_j : Q_j \in A\}$. Then, obviously, there exist the squares Q_1 and Q_2 from the above-mentioned partitioning such that $\zeta \in Q_1 \subset Q_2$, $Q_1 \notin F(G, k)$, $Q_2 \in F(G, k)$, and

$$\frac{1}{|Q_2|} \iint_{Q_2} |f^*(z) - f_Q^*| < \alpha,$$

whence it follows that

$$\frac{1}{|Q_1|} \iint_{Q_1} |f^*(z) - f_Q^*| d\sigma_z < \frac{4}{|Q_2|} \iint_{Q_2} |f^*(z) - f_Q^*| d\sigma_z < 4\alpha.$$

Then, denoting by z_1 the center of the square Q_1 and using Lemma 3 and the mean value theorem, we obtain

$$\begin{aligned} |f(\zeta) - f_{Q_1}| &\leq |f(\zeta) - f(z_1)| + |f(z_1) - f_{Q_1}| \leq \\ &\leq 4\|f\|_{\text{BMO}(G)} + \frac{2}{|Q_1|} \iint_{Q_1} |f(z) - f_{Q_1}| d\sigma_z \leq 12\alpha. \end{aligned}$$

Thus the validity of the first requirement of Lemma 2 is proved.

Further, it is evident that the left-hand side of the “double” inequality (2.14) holds for all $Q_j \in A$. Let us show that the right-side of that inequality is also valid.

Let Q_j^* be a square whose partitioning into four equal squares gives the square $Q_j \in A$. Clearly, $Q_j^* \supset Q_j$, and

$$\frac{1}{|Q_j^*|} \iint_{Q_j^*} |f^*(z) - f_Q^*| d\sigma_z < \alpha.$$

Taking into account the above inequality, we obtain

$$\frac{1}{|Q_j|} \iint_{Q_j} |f^*(z) - f_Q^*| d\sigma_z < \frac{4}{|Q_j^*|} \iint_{Q_j^*} |f^*(z) - f_Q^*| d\sigma_z < 4\alpha.$$

Thus relation (2.14) is proved.

Finally, using the already proven relation (2.14) and inequality (2.10) (the case where $\mu = \sigma$ is the Lebesgue measure), we get

$$\begin{aligned} |\bigcup_A Q_j| &= \sum_{Q_j \in A} |Q_j| \leq \frac{1}{\alpha} \sum_{Q_j \in A} \iint_{Q_j} |f^*(z) - f_Q^*| d\sigma_z \leq \\ &\leq \frac{1}{\alpha} \iint_Q |f^*(z) - f_Q^*| d\sigma_z \leq \frac{1}{\alpha} |Q| \|f\|_{\text{BMO}(G)}. \quad \square \end{aligned}$$

The proof of the following lemma can be found in [11]. Let us formulate it in a way convenient for us.

Lemma 5. *Let G be a finite domain with a quasiconformal boundary Γ , $z_0 \in \Gamma$, $n, m \in \mathbb{N}$, $n > m$. Then*

$$\left(\frac{m}{n}\right)^2 \asymp \frac{\rho_n(z_0)}{\rho_m(z_0)} \asymp \left(\frac{m}{n}\right)^\beta, \quad (2.16)$$

where $\beta = \beta(G) > 0$ is a constant depending only on G .

In particular, from relation (2.16) we obtain the known inequality

$$\rho_n(z_0) \gtrsim \left(\frac{1}{n}\right)^2. \tag{2.17}$$

Lemma 6. *Let G be a finite domain with a quasi-conformal boundary Γ , $p > 1$, $w \in A_p(F(G))$, $z_0 \in \Gamma$, $u(z_0, r) = \{z : |z - z_0| < r\}$, let μ be a measure defined by equality (1.1), and let $\{\Pi_n(z)\}_{n=1}^\infty$ be a sequence of algebraic polynomials of order not higher than n such that*

$$\|\Pi_n\|_{z_0, p, w} \leq c_1 (\mu\{u(z_0, \rho_n(z_0))\})^{\frac{1}{p}},$$

where c_1 is a constant not depending on z_0 and n .

Then for all $z \in u(z_0, \rho_n(z_0))$ we have the inequality

$$|\Pi'_n(z)| \leq c_2 |\rho_n(z_0)|^{-1},$$

where c_2 is a constant not depending on z_0 and n .

This lemma is the analogue of the well-known theorem on the derivatives of algebraic polynomials [12, p.420], [13] which can be proved analogously to the result of [3, p.14].

3. PROOFS OF THE BASIC RESULTS

Proof of Theorem 1. For brevity we shall use the notation $\mu\{u(z_0, t) \cap G\} = \mu(z_0, t)$ ($t \geq 0$).

Let us prove first the necessity. Assume that $f \in \text{BMO}_p(G, w) \cap H'(G)$ and let us show that relation (1.4) holds.

Let $n \in N$, $z_0 \in \Gamma$, $Q = Q(z_0, \rho_n(z_0))$, and μ and μ^* be the measures defined by the equalities (1.1) and (2.3). Relations (2.8), (2.6), and (2.17) yield

$$\mu(Q \cap G) \gtrsim \mu(Q^*) \gtrsim [\rho_n(z_0)]^2 \gtrsim \left(\frac{1}{n}\right)^4,$$

but then, obviously, we shall have

$$f_{\mu, Q \cap G} \preccurlyeq \frac{1}{\mu(Q \cap G)} \iint_G |f| d\mu \preccurlyeq c(f, \mu) \left(\frac{1}{n}\right)^{-4}. \tag{3.1}$$

Clearly, $f \in H'_p(G, w)$. But then, repeating the arguments (and taking into account (3.1) cited in [2, pp. 174, 182]), we can see that there exists a

sequence of algebraic polynomials P_n of order not higher than n , such that

$$\begin{aligned} \|f - P_n\|_{z_0, p, w} &\leq c\rho_n(z_0) \left(\mu(z_0, \rho_n(z_0)) \right)^{\frac{1}{p}} \times \\ &\times \int_{\rho_n(z_0)}^{\infty} \frac{\sigma_p(f - f_{\mu, Q \cap G}, w, z_0, t)}{t^2 \mu^{1/p}(z_0, t)} dt. \end{aligned} \quad (3.2)$$

Now let us estimate the value $\sigma_p(f - f_{\mu, Q \cap G}, w, z_0, t)$ for $t \geq \rho_n(z_0)$.

Let $Q_m = Q(z_0, 2^m \rho_n(z_0))$ ($m \in N$), $Q_0 = Q$. Using relations (2.8) and (2.3) for all $m \geq 1$, we get

$$\mu(Q_{m-1} \cap G) \succcurlyeq \mu^*(Q_{m-1}) \succcurlyeq \mu^*(2Q_{m-1}) = \mu^*(Q_m) \geq \mu(Q_m \cap G).$$

Then, on account of (1.3), we have

$$|f_{\mu, Q_m \cap G} - f_{\mu, Q_{m-1} \cap G}| \preccurlyeq \frac{1}{\mu(Q_m \cap G)} \iint_{Q_m \cap G} |f - f_{\mu, Q_{m-1} \cap G}| d\mu \leq \|f\|_{\text{BMO}_p(G, w)}.$$

Thus, using the Minkowsky inequality and relation (1.3) for all $k \geq 1$, we obtain

$$\begin{aligned} \sigma_p(f - f_{\mu, Q \cap G}, w, z_0, 2^k \rho_n(z_0)) &\leq \left(\iint_{Q_k \cap G} |f - f_{\mu, Q_k \cap G}|^p d\mu \right)^{\frac{1}{p}} + \\ &+ \sum_{m=1}^k |f_{\mu, Q_m \cap G} - f_{\mu, Q_{m-1} \cap G}| \left(\iint_{Q_k \cap G} d\mu \right)^{\frac{1}{p}} \leq \\ &\leq \|f\|_{\text{BMO}_p(G, w)} (1+k) \left(\iint_{Q_k \cap G} d\mu \right)^{\frac{1}{p}} = \\ &= \|f\|_{\text{BMO}_p(G, w)} \left(1 + \log_2 \frac{2^k \rho_n(z_0)}{\rho_n(z_0)} \right) \cdot \mu^{\frac{1}{p}}(z_0, 2^k \rho_n(z_0)), \end{aligned}$$

whence, obviously,

$$\sigma_p(f - f_{\mu, Q \cap G}, w, z_0, t) \leq \|f\|_{\text{BMO}_p(G, w)} \left(1 + \log_2 \frac{t}{\rho_n(z_0)} \right) \cdot \mu^{\frac{1}{p}}(z_0, t)$$

for all $t \geq \rho_n(z_0)$.

Consequently, owing to (3.2), we have

$$\begin{aligned} \|f - P_n\|_{z_0, p, w} &\preccurlyeq \\ &\preccurlyeq \|f\|_{\text{BMO}_p(G, w)} \rho_n(z_0) \left(\mu(z_0, \rho_n(z_0)) \right)^{\frac{1}{p}} \int_{\rho_n(z_0)}^{\infty} \frac{(1 + \log_2 \frac{t}{\rho_n(z_0)})}{t^2} dt = \end{aligned}$$

$$\begin{aligned}
&= \|f\|_{\text{BMO}_p(G,w)} \left(\mu(z_0, \rho_n(z_0)) \right)^{\frac{1}{p}} \int_1^\infty \frac{(1 + \log_2 \tau)}{\tau^2} d\tau \preccurlyeq \\
&\preccurlyeq \|f\|_{\text{BMO}_p(G,w)} \left(\mu\{u(z_0, \rho_n(z_0)) \cap G\} \right)^{\frac{1}{p}}. \quad \square
\end{aligned}$$

Assume now that relation (1.4) is fulfilled for some function f given in G . Then, obviously, $f \in L_p(G, w)$. Moreover, $f \in H^1(G)$ if we neglect the values of the function f on the set of measure zero. Indeed, if $z \in G$ is an arbitrary point and $z^* \in \Gamma$ is a point such that $|z - z^*| = \rho(z, \Gamma)$, then it is easy to check that $u(z, \rho(z, \Gamma)) \subset G_n(z^*, c_0)$ ($c_0 \leq \frac{1}{2}$). Taking into account relation (1.4) it is not difficult to prove that the polynomials P_n converge uniformly on $u(z, \frac{1}{2}\rho(z, \Gamma))$. Clearly, the limiting analytic function coincides with the functions f a.e.

Further, let $z_0 \in \Gamma$, $r > 0$, $Q = Q(z_0, r)$, and let $n \in N$ be a number such that

$$\rho_{2^{n+1}}(z_0) < r \leq \rho_{2^n}(z_0). \quad (3.3)$$

Using the Minkowsky inequality, we can see that

$$\begin{aligned}
\left(\iint_{Q \cap G} |f(z) - f_{\mu, Q \cap G}|^p d\mu_z \right)^{\frac{1}{p}} &\leq \left(\iint_{Q \cap G} |f(z) - P_{2^n}(z)|^p d\mu_z \right)^{\frac{1}{p}} + \\
&\left(\iint_{Q \cap G} |P_{2^n}(z) - f_{\mu, Q \cap G}|^p d\mu_z \right)^{\frac{1}{p}} \stackrel{df}{=} I_1 + I_2.
\end{aligned}$$

By virtue of (1.4),

$$I_1 \leq \text{const } \mu^{1/p}(z_0, \rho_n(z_0)). \quad (3.4)$$

It remains to estimate I_2 . Evidently,

$$I_2 \preccurlyeq \mu^{1/p}(Q \cap G) \cdot \max_{z \in Q \cap G} |P_{2^n}(z) - f_{\mu, Q \cap G}|. \quad (3.5)$$

Then it is obvious that for all $z \in Q \cap G$

$$\begin{aligned}
|P_{2^n}(z) - f_{\mu, Q \cap G}| &\leq \frac{1}{\mu(Q \cap G)} \iint_{Q \cap G} |f(\zeta) - P_{2^n}(z)| d\mu_\zeta \leq \\
&\leq \frac{1}{\mu(Q \cap G)} \iint_{Q \cap G} |f(\zeta) - P_{2^n}(\zeta)| d\mu_\zeta + \\
&+ \max_{\zeta \in Q \cap G} |P_{2^n}(\zeta) - P_{2^n}(z)| \stackrel{df}{=} I_2' + I_2''.
\end{aligned}$$

Using the Hölder inequality and relation (1.4), we obtain

$$I'_2 \leq \frac{1}{\mu(Q \cap G)} \left(\iint_{Q \cap G} |f - P_{2^n}|^p d\mu \right)^{\frac{1}{p}} \left(\iint_{Q \cap G} d\mu \right)^{1-\frac{1}{p}} \preccurlyeq \text{const}. \quad (3.6)$$

To estimate I''_1 , let us consider the polynomial

$$\Pi_{2^k}(z) = P_{2^k}(z) - P_{2^{k-1}}(z) \quad (k \geq 1).$$

By (2.15), the Minkowsky inequality and relation (1.4) imply

$$\|\Pi_{2^k}\|_{z_0, p, w} \leq \|f - P_{2^k}\|_{z_0, p, w} + \|f - P_{2^{k-1}}\|_{z_0, p, w} \preccurlyeq \mu^{1/p}(z_0, \rho_n(z_0)).$$

But then, according to Lemma 6, we have

$$|\Pi'_{2^k}(z)| \preccurlyeq |\rho_{2^k}(z_0)|^{-1} \quad \forall (z \in u(z_0, \rho_{2^k}(z_0))), \quad k \geq 1).$$

Hence, taking into account (2.15), we obtain

$$\begin{aligned} & \left| P_{2^n}(\zeta) - P_{2^n}(z) \right| = \left| (P_1(\zeta) - P_1(z)) + \right. \\ & \left. + \sum_{k=1}^n \left(\Pi_{2^k}(\zeta) - \Pi_{2^k}(z) \right) \right| \preccurlyeq |\zeta - z| + \sum_{k=1}^n \int_{[\zeta, z]} |\Pi'_{2^k}(\xi)| |d\xi| \preccurlyeq \\ & \preccurlyeq |\zeta - z| \left(1 + \sum_{k=1}^n |\rho_{2^k}(z_0)|^{-1} \right) \preccurlyeq \left(\rho_{2^n}(z_0) + \sum_{k=1}^n \frac{\rho_{2^n}(z_0)}{\rho_{2^k}(z_0)} \right) \preccurlyeq \\ & \preccurlyeq \left(\rho_{2^n}(z_0) + \sum_{k=1}^n \left(\frac{1}{2^{n-k}} \right)^\beta \right) \leq \text{const} \end{aligned}$$

for all $z, \zeta \in Q \cap G$.

This means that $I''_2 \leq \text{const}$.

But then, taking into account (3.6) and (3.5), we get

$$I_2 \leq \text{const} \mu^{1/p}(z_0, \rho_n(z_0)),$$

which, with regard to (3.4), completes the proof of Theorem 1. \square

Proof of Theorem 2. Let us prove first that (b) \Rightarrow (a). Let $\omega \in A_p(F(G))$ ($1 < p < \infty$), $f \in \text{BMO}_p(G, \omega)$ and let us show that $f \in \text{BMO}(G)$.

Indeed,

$$\begin{aligned} & \frac{1}{|Q \cap G|} \iint_{Q \cap G} |f - f_{Q \cap G}| d\sigma \leq \\ & \leq \frac{2}{|Q \cap G|} \left(\iint_{Q \cap G} |f - f_{\mu, Q \cap G}|^p w d\sigma \right)^{\frac{1}{p}} \left(\iint_{Q \cap G} w^{-\frac{1}{p-1}} d\sigma \right)^{\frac{p-1}{p}} \preccurlyeq \end{aligned}$$

$$\begin{aligned} &\leq \left(\frac{1}{\mu(Q \cap G)} \iint_{Q \cap G} |f - f_{\mu, Q \cap G}|^p d\mu \right)^{\frac{1}{p}} \times \\ &\times \left(\frac{1}{|Q \cap G|} \iint_{Q \cap G} w \right)^{\frac{1}{p}} \left(\frac{1}{|Q \cap G|} \iint_{Q \cap G} w^{-\frac{1}{p-1}} \right)^{\frac{p-1}{p}} \leq \text{const} \|f\|_{\text{BMO}(G)}. \end{aligned}$$

It remains to prove that (a) \Rightarrow (b).

Let $f \in \text{BMO}(G)$ be an analytic function in the domain G , let f^* be a function defined by the equality (2.9), $\omega \in A_p(F(G))$ ($1 < p < \infty$), and let μ^* be the measure defined by (2.3). We prove first that for all $Q \in F(G)$ and $\lambda > 0$ the relation

$$\frac{1}{\mu^*(Q)} \mu^* \{z \in Q : |f^*(z) - f_Q^*| > \lambda\} \leq C \exp\left(\frac{-c\lambda}{\|f\|_{\text{BMO}(G)}}\right) \quad (3.7)$$

holds, where C and c are the constants independent of f^* , Q , and λ .

Choose a square $Q \in F(G)$. Let δ be the number from Lemma 1, and let c^* be the constant from (2.10). Without loss of generality we can assume that $\delta < 1/c^*$. In this case we can apply Lemma 4 to the function f^* and $\alpha = (1/\delta)\|f\|_{\text{BMO}(G)}$. Hence we get a family of disjoint squares $A_1 = \{Q_j^1 : Q_j^1 \in F(G, k)\}$ such that

$$|f^*(z) - f_Q^*| \leq 12\alpha$$

for all $z \in Q \setminus \bigcup_{A_1} Q_j^1$,

$$|f_{Q_j^1}^* - f_Q^*| < 4\alpha \quad (3.8)$$

according to (2.14), and by (2.15) we have

$$|\bigcup_{A_1} Q_j^1| = \sum_{A_1} |Q_j^1| \leq \frac{1}{\alpha} \|f\|_{\text{BMO}(G)} \cdot |Q|.$$

Since $(1/\alpha)\|f\|_{\text{BMO}(G)} = \delta$, by virtue of Lemma 1 we obtain

$$\mu^* \{ \bigcup_{A_1} Q_j^1 \} \leq \varepsilon \mu^*(Q). \quad (3.9)$$

Applying again Lemma 4 to the function f^* and $\alpha = (1/\delta)\|f\|_{\text{BMO}(G)}$, for every Q_j^1 we obtain a family of nonintersecting squares $A_2 = \{Q_j^2 : Q_j^2 \in F(G, k)\}$ such that each of these squares is contained in one of the Q_j^1 . Thus, by (3.8) and (2.13) the relation

$$|f^* - f_Q^*| \leq |f^* - f_{Q_j^1}^*| + |f_{Q_j^1}^* - f_Q^*| < 12\alpha + 4\alpha < 2 \cdot 12\alpha$$

is fulfilled on $Q \setminus \bigcup_{A_2} Q_j^2$, while owing to (2.14) and (3.8) we have that

$$|f_{Q_j^2}^* - f_Q^*| \leq |f_{Q_j^2}^* - f_{Q_j^1}^*| + |f_{Q_j^1}^* - f_Q^*| < 4\alpha + 4\alpha < 2 \cdot 12\alpha.$$

Finally, according to (2.15), we have

$$\left| \bigcup_{Q_j^2 \subset Q_j^1} Q_j^2 \right| = \sum_{Q_j^2 \subset Q_j^1} |Q_j^2| \leq \frac{1}{\alpha} \|f\|_{\text{BMO}(G)} \cdot |Q_j^1|$$

for every Q_j^1 .

Then again, by virtue of Lemma 1 and (3.9), we obtain

$$\mu^* \left\{ \bigcup_{A_2} Q_j^2 \right\} = \sum_{Q_j^1 \in A_1} \mu^* \left\{ \bigcup_{Q_j^2 \subset Q_j^1} Q_j^2 \right\} \leq \sum_{Q_j^1 \in A_1} \varepsilon \mu^*(Q_j^1) \leq \varepsilon^2 \mu^*(Q).$$

Continuing this process ad infinitum, we obtain at the step n a family of intersecting squares $A_n = \{Q_j^n\}$ such that

$$|f^* - f_{Q_j^n}^*| \leq 12\alpha \cdot n \quad \text{a.e. in } Q \setminus \bigcup_{A_n} Q_j^n \quad \text{and} \quad \mu^* \left\{ \bigcup_{A_n} Q_j^n \right\} \leq \varepsilon^n \mu^*(Q).$$

Assume now that $\lambda > 12\alpha$. Let $n \geq 1$ be a natural number such that $12\alpha n < \lambda \leq 12\alpha n + 12\alpha$. Then, obviously, we shall have

$$\begin{aligned} \mu^* \{z \in Q : |f^*(z) - f_Q^*| > \lambda\} &\leq \mu^* \{z \in Q : |f^*(z) - f_Q^*| > 12\alpha n\} \leq \\ &\leq \mu^* \left\{ \bigcup_{A_n} Q_j^n \right\} \leq \varepsilon^n \mu^*(Q) \leq \varepsilon^{\frac{\lambda}{12\alpha} - 1} \mu^*(Q) = \frac{1}{\varepsilon} \exp \left(\frac{-c\lambda}{\|f\|_{\text{BMO}(G)}} \right) \mu^*(Q) \end{aligned}$$

for $c = (1/12)\delta \cdot \ln(1/\varepsilon)$.

Hence, estimate (3.7) is valid for all $\lambda > 12\alpha$. But for all $0 < \lambda \leq 12\alpha$ we, obviously, have

$$\begin{aligned} \mu^* \{z \in Q : |f^*(z) - f_Q^*| > \lambda\} &\leq \mu^*(Q) = \exp \left(\frac{c\lambda}{\|f\|_{\text{BMO}(G)}} \right) \times \\ &\times \exp \left(\frac{-c\lambda}{\|f\|_{\text{BMO}(G)}} \right) \mu^*(Q) \leq \exp \left(\frac{12c}{\delta} \right) \exp \left(\frac{-c\lambda}{\|f\|_{\text{BMO}(G)}} \right) \mu^*(Q). \end{aligned}$$

Consequently, assuming $C = \max \left\{ \frac{1}{\varepsilon}, \exp \left(\frac{12c}{\delta} \right) \right\}$, we get estimate (3.7) for all $\lambda > 0$.

Relation (3.7) with regard to (2.8) implies that

$$\frac{1}{\mu(Q \cap G)} \mu \{z \in Q \cap G : |f(z) - f_Q^*| > \lambda\} \leq \exp \left(\frac{-c\lambda}{\|f\|_{\text{BMO}(G)}} \right). \quad (3.10)$$

The latter relation allows us to complete the proof of Theorem 2. Indeed, using first the Minkowsky inequality and then writing the corresponding

integral in terms of a distribution function, applying estimate (3.10), we obtain

$$\begin{aligned} & \left(\frac{1}{\mu(Q \cap G)} \iint_{Q \cap G} |f - f_{\mu, Q \cap G}|^p d\mu \right) \leq \left(\frac{2}{\mu(Q \cap G)} \iint_{Q \cap G} |f - f_Q^*|^p d\mu \right)^{\frac{1}{p}} = \\ & = \left(2p \int_0^\infty \lambda^{p-1} \frac{1}{\mu(Q \cap G)} \mu\{z \in Q \cap G : |f(z) - f_Q^*| > \lambda\} d\lambda \right)^{\frac{1}{p}} \asymp \\ & \asymp \left(2p \int_0^\infty \lambda^{p-1} \exp\left(\frac{-c\lambda}{\|f\|_{\text{BMO}(G)}}\right) d\lambda \right)^{\frac{1}{p}} \asymp c(p) (\|f\|_{\text{BMO}(G)})^{\frac{1}{p}}, \end{aligned}$$

which implies that $f \in \text{BMO}_p(G, w)$. \square

REFERENCES

1. L. N. Chikvinidze, Approximate functions in domains with quasi-conformal boundary in weighted plane integral metrics. (Russian) *Sibirskii Mat. J.* **XXX**(1989), No. 2, 172–184; English translation: *Siberian Math. J.* **30**(1989), No. 2, 306–316.
2. L. N. Chikvinidze, On approximation of functions in domains with quasiconformal boundary in weighted plane integral metrics. (Russian) *Dokl. Akad. Nauk SSSR* **304**(1989), No. 1, 41–43; English translation: *Soviet. Math. Dokl.* **39**(1989), No. 1, 32–34.
3. L. N. Chikvinidze, Local approximation of analytic functions and their derivatives in integral metrics in domains with a quasi-conformal boundary. *Proc. A. Razmadze Math. Inst.* **101**(1992), 5–26.
4. L. V. Ahlfors, Lectures on quasiconformal mappings. *Van Nostrand, Princeton, etc.*, 1966.
5. J. B. Garnett, Bounded analytic functions. *Academic Press, New York*, 1981.
6. D. Bekolle and A. Bonami, Inegalites a poids pour le noyau de Bergman. *C.R. Acad. Sci., Paris* **AB286**(1978), No. 18, 775–778.
7. B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal functions. *Trans. Amer. Math. Soc.* **165**(1972), 207–226.
8. I. M. Batchaev, Integral representations in domains with quasi-conformal boundary and some of their applications. (Russian) Abstract of Thesis for Cand. Sci. Degree. *Math. Inst. Acad. Sci. Azerb. SSR, Baku*, 1981.
9. L. N. Chikvinidze, On one variant of Hardy and Littlewood maximal function. (Russian) *Trudy Tbiliss. Mat. Inst. Razmadze* **86**(1987), 101–109.

10. L. N. Chikvinidze, On an analogue of the Muckenhoupt condition in domains with quasi-conformal boundary. (Russian) *Trudy Tbiliss. Mat. Inst. Razmadze* **88**(1989), 41–58.
11. V. V. Andrievsky, Geometrical properties of V. K. Dzyadyk domains. (Russian) *Ukrain. Math. Zh.* **33**(1981), No. 6, 723–727.
12. V. K. Dzyadyk, Introduction to the theory of uniform approximation of functions by polynomials. (Russian) *Nauka, Moscow*, 1977.
13. P. M. Tamrazov, Smoothness and polynomial approximations. (Russian) *Naukova Dumka, Kiev*, 1975.

(Received 21.09.1994)

Author's address:
A. Razmadze Mathematical Institute
Georgian Academy of Sciences
1, M. Alexidze St., Tbilisi 380093
Republic of Georgia