

OSCILLATORY CRITERIA FOR NONLINEAR n TH-ORDER DIFFERENTIAL EQUATIONS WITH QUASIDERIVATIVES

MIROSLAV BARTUŠEK

ABSTRACT. Sufficient conditions are given for the existence of oscillatory proper solutions of a differential equation with quasiderivatives $L_n y = f(t, L_0 y, \dots, L_{n-1} y)$ under the validity of the sign condition $f(t, x_1, \dots, x_n) x_1 \leq 0, f(t, 0, x_2, \dots, x_n) = 0$ on $\mathbb{R}_+ \times \mathbb{R}^n$.

1. INTRODUCTION

Consider the n th-order differential equation

$$L_n y(t) = f(t, L_0 y, L_1 y, \dots, L_{n-1} y) \quad \text{in } \mathbb{D} = \mathbb{R}_+ \times \mathbb{R}^n, \quad (1)$$

where $n \geq 2, \mathbb{R}_+ = [0, \infty], \mathbb{R} = (-\infty, \infty), L_i y$ is the i th quasiderivative of y defined as

$$L_0 y(t) = \frac{y(t)}{a_0(t)}, L_i y(t) = \frac{(L_{i-1} y(t))'}{a_i(t)}, \quad i = 1, 2, \dots, n-1, \quad (2)$$

$$L_n y(t) = (L_{n-1} y(t))',$$

functions $a_i \in C^\circ(\mathbb{R}_+)$ are positive, and $f : \mathbb{D} \rightarrow \mathbb{R}$ fulfills the local Carathéodory conditions.

Throughout the paper we assume that

$$f(t, x_1, \dots, x_n) x_1 \leq 0, f(t, 0, x_2, \dots, x_n) = 0 \quad \text{in } \mathbb{D}. \quad (3)$$

Definition. A function $y : [0, T) \rightarrow \mathbb{R}, T \in (0, \infty]$, is called a solution of (1) if (1) is valid for almost all $t \in [0, T)$. It is called noncontinuable if either $T = \infty$ or $T < \infty$, and

$$\limsup_{t \rightarrow T} \sum_{i=0}^{n-1} |L_i y(t)| = \infty.$$

1991 *Mathematics Subject Classification.* 34C10.

Key words and phrases. Oscillatory solutions, existence criteria, differential equation of n th order.

Let $y : [0, T) \rightarrow \mathbb{R}$, $T \leq \infty$, be a noncontinuable solution of (1). It is said to be proper if $T = \infty$ and $\sup_{\tau \leq t < \infty} |y(t)| > 0$ for all $\tau \in \mathbb{R}_+$. It is said to be singular of the first (second) kind if $t^* \in (0, \infty)$ exists such that

$$y \equiv 0 \text{ in } [t^*, \infty), \quad \sup_{0 \leq t \leq t^*} \sum_{i=0}^{n-1} |L_i y(t)| > 0$$

(if $T < \infty$). A proper solution y is said to be oscillatory if a sequence $\{t_k\}_0^\infty$ exists such that $t_k \in \mathbb{R}_+$, $\lim_{k \rightarrow \infty} t_k = \infty$ and $y(t_k) = 0$ holds. Otherwise, it is called nonoscillatory.

Many authors studied the problem of structure and properties of proper nonoscillatory solutions of (1) (see, e.g., [1]–[3]). But as regards proper oscillatory solutions, their existence is proved only in the cases where $n \geq 3$ and $a_i \equiv 1$ (see [4]–[6]), or $n = 3$ (see [1]).

Definition. Equation (1) has property A if every proper solution y is oscillatory for even n and it is either oscillatory or

$$\lim_{t \rightarrow \infty} L_i y(t) = 0 \text{ monotonically, } i = 0, 1, \dots, n-1, \quad (4)$$

for odd n .

Similarly to a differential equation without quasiderivatives ($a_i \equiv 1$), it is possible to use the following way to prove the existence of proper oscillatory solutions: If

- 1° there exists no singular solution of the 1st kind;
- 2° there exists no singular solution of the 2st kind;
- 3° (1) has Property A;
- 4° the initial conditions of y at zero are chosen such that (4) is not valid,

then y is oscillatory proper.

Sufficient conditions for the validity of relations 1°, 2°, 4° can be easily obtained similarly to the case $a_i \equiv 1$ (see later). Very profound results concerning 3° are given in [7].

In our paper we generalize the results which could be obtained by this approach. Especially, we shall weaken conditions 1° and 3°.

Sometimes, we will suppose that

$$a_n(t)|x_1|^\lambda \leq |f(t, x_1, \dots, x_n)| \text{ in } \mathbb{D}, \quad (5)$$

where $0 < \lambda \leq 1$, $a_n \in L_{\text{loc}}(\mathbb{R}_+)$, $a_n \geq 0$;

$$\int_0^\infty a_i(t) dt = \infty, \quad i = 1, 2, \dots, n - 1, \tag{6}$$

$$|f(t, x_1, \dots, x_n)| \leq h(t)\omega\left(\sum_{i=1}^n |x_i|\right) \quad \text{in } \mathbb{D}, \tag{7}$$

where $h \in L_{\text{loc}}(\mathbb{R}_+)$, $\omega \in C^\circ(\mathbb{R}_+)$, $\omega(x) > 0$ for $x > 0$, $\int_0^\infty \frac{dt}{\omega(t)} = \infty$;

$$|f(t, x_1, \dots, x_n)| \leq A(t)g(|x_1|) \quad \text{in } \mathbb{R}_+ \times [-\varepsilon, \varepsilon]^n, \tag{8}$$

where $\varepsilon > 0$, $A \in L_{\text{loc}}(\mathbb{R}_+)$, $g \in C^\circ[0, \varepsilon]$, $g(0) = 0$, $g(x) > 0$ for $x > 0$,

$$\int_0^\varepsilon \frac{dt}{g(t)} = \infty;$$

$$\left\{ \begin{array}{l} \text{let } \frac{a_1}{a_2} \in C^1(\mathbb{R}_+) \text{ for } n = 3, \\ a_1 \in C^1(\mathbb{R}_+), a_2 \in C^1(\mathbb{R}_+), \frac{a_3}{a_1} \in C^2(\mathbb{R}_+) \text{ for } n = 4 \\ \text{and let for } n > 4 \text{ an index } l \in \{1, 2, \dots, n - 4\} \text{ exist} \\ \text{such that } a_{l+j}, j = 1, 2, \text{ are absolutely continuous and} \\ a'_{l+j}, j = 1, 2, \text{ are locally bounded from below.} \end{array} \right. \tag{9}$$

Notation. If $b_i \in C^\circ(I)$, then

$$I^\circ(t) \equiv 1, \quad I^k(t, b_1, \dots, b_k) = \int_0^t b_1(s)I^{k-1}(s, b_2, \dots, b_k) ds, \quad t \in I.$$

Put $a_{n_j+i}(t) = a_i(t)$, $j \in \{\dots, -1, 0, 1, \dots\}$, $i \in \{0, 1, \dots, n\}$, $N = \{1, 2, \dots\}$.

2. MAIN RESULTS

Further, we shall investigate a solution y of (1) that satisfies the initial conditions

$$l \in \{0, 1, \dots, n - 1\}, \tau \in \{-1, 1\}, \tau L_i y(0) > 0, \quad i = 0, 1, \dots, l, \tag{10}$$

$$\tau L_j y(0) < 0, \quad j = l + 1, \dots, n - 1,$$

and we shall prove that this solution is oscillatory proper under the validity of certain assumptions.

Theorem 1. Let $\lambda \in (0, 1)$ and let (5), (7), and (9) be valid. Let

$$\int_0^\infty a_{i+1}(\tau_{i+1}) \int_0^{\tau_{i+1}} a_{i+2}(\tau_{i+2}) \int_0^{\tau_{i+2}} \cdots \int_0^{\tau_{n-1}} a_n(\tau_n) \left[\int_0^{\tau_n} a_{n+1}(\tau_{n+1}) \cdots \int_0^{\tau_{i+n-1}} a_{i+n}(\tau_{i+n}) d\tau_{i+n} \cdots d\tau_{n+1} \right]^\lambda \times d\tau_n \cdots d\tau_{i+1} = \infty, \quad (11)$$

$$i = 0, 1, \dots, n-1.$$

Then any solution y of (1) that fulfills the Cauchy initial conditions (10) is oscillatory proper.

Theorem 2. Let $\lambda = 1$, (5), (6), and (7) hold. Let

$$\limsup_{t \rightarrow \infty} I^1(a_{n-1}) \int_t^\infty \frac{I^{n-1}(s, a_1, \dots, a_{n-1})}{I^1(s, a_{n-1})} a_n(s) ds > 1. \quad (12)$$

Further, let either (9) or (8) hold.

Then any solution y of (1), that fulfills the Cauchy initial conditions (10) is oscillatory proper.

Theorem 3. Let (6), (7) be valid and let functions $a_n \in L_{loc}(\mathbb{R}_+)$, $b \in C^\circ(\mathbb{R}_+)$ exist such that $\int_0^\infty a_n(t) dt = \infty$, $b(0) = 0$, $b(x) > 0$ for $x > 0$, b is nondecreasing, and

$$a_n(t)b(|x_1|) \leq |f(t, x_1, \dots, x_n)| \quad \text{in } \mathbb{D}.$$

Further, let either (9) or (8) be valid. Then any solution y of (1) that fulfills (10) is oscillatory proper.

3. PROOF OF MAIN RESULTS

Let us define two special types of solutions of (1) that will be encountered later.

Type I (τ): $y : [0, \tau) \rightarrow \mathbb{R}$, $0 < \tau \leq \infty$ and sequences $\{t_k^i\}$, $\{\bar{t}_k^{n-1}\}$, $k \in \mathbb{N}$, $i \in \{0, 1, \dots, n-1\}$ exist such that $\lim_{k \rightarrow \infty} t_k^0 = \tau$,

$$0 \leq t_k^0 < t_k^{n-1} \leq \bar{t}_k^{n-1} < t_k^{n-2} \cdots < t_k^1 < t_{k+1}^0,$$

$L_i y(t_k^i) = 0, i = 0, 1, \dots, n - 2, L_{n-1} y(t) = 0$ for $t \in [t_k^{n-1}, \bar{t}_k^{n-1}], k \in \mathbb{N}$,

$$\begin{aligned} L_i y(t) L_0 y(t) &> 0 \quad \text{for } t \in (t_k^0, t_k^i), i = 0, 1, \dots, n - 1, \\ &< 0 \quad \text{for } t \in (t_k^i, t_{k+1}^0), i = 0, 1, \dots, n - 2, \\ &< 0 \quad \text{for } t \in (\bar{t}_k^{n-1}, t_{k+1}^0), i = n - 1, k \in \mathbb{N}. \end{aligned}$$

If $\tau < \infty$, then $\lim_{t \rightarrow \tau} L_i y(t) = 0, i = 0, 1, \dots, n - 1$.

Type II (s): $y : \mathbb{R}_+ \rightarrow \mathbb{R}, s \in \{0, 1, \dots, n - 1\}, \tau \in \mathbb{R}_+$,

$$\begin{aligned} L_j y(t) L_s y(t) &\geq 0 \quad \text{for } j \in \{0, 1, \dots, s\} \\ &\leq 0 \quad \text{for } j \in \{s + 1, \dots, n - 1\}, \\ L_m y(t) &\neq 0, m \in \{0, 1, \dots, n - 2\}, t \in [\tau, \infty). \end{aligned}$$

Remark. Any solution y of Type I (∞) (of Type II (s)) is oscillatory proper (nonoscillatory proper). If we define $y \equiv 0$ on $[\tau, \infty)$, then any solution y of Type I(τ), $\tau < \infty$ is singular of the first kind.

Lemma 1. Let $J = [t_1, t_2] \subset \mathbb{R}_+, t_1 < t_2$ and $y : J \rightarrow \mathbb{R}$ be a solution of (1).

- (a) If $j \in \{1, 2, \dots, n\}, L_j y(t) \geq 0$ (≤ 0) in J , then $L_{j-1} y$ is nondecreasing (nonincreasing) in J ;
- (b) if $j \in \{1, 2, \dots, n\}, L_j y(t) > 0$ (< 0) in J , then $L_{j-1} y$ is increasing (decreasing) in J ;
- (c) if $L_0 y(t) \geq 0$ (≤ 0) in J , then $L_{n-1} y$ is nonincreasing (nondecreasing) in J .

Proof.

- (a) Let $L_j y(t) \geq 0$ in J . Then according to (2) either $(L_{j-1} y(t))' = a_j(t) L_j y(t) \geq 0, j < n$ or $(L_{n-1} y(t))' = L_n y(t) \geq 0$ holds.
- (b), (c) The proof is similar, only (3) must be used instead of (2) in (c). \square

Lemma 2. Let $y : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a solution of (1) which satisfies (10). Then one of the following possibilities holds:

- (a) y is of Type I (∞)
- (b) there exists $\tau \in (0, \infty)$ such that y is of Type I(τ) in $[0, \tau)$.
- (c) there exists $i \in \{0, \dots, n - 1\}$ such that y is of Type II (i).

Proof. First suppose that y satisfies the Cauchy initial conditions

$$\sigma L_i y(0) > 0, \quad i = 0, 1, \dots, n - 1. \tag{13}$$

According to Lemma 1 $\sigma L_i y > 0, i = 0, 1, \dots, n - 1$, in some right neighborhood of $t = 0$, and $\sigma L_j y, j = 0, 1, \dots, n - 2$, are nondecreasing ($\sigma L_{n-1} y$

is nondecreasing) until $\sigma L_{j+1}y \geq 0$ ($\sigma L_0y \geq 0$). Thus either y is of Type II ($n-1$) or numbers t^n, \bar{t}^n exist such that

$$\begin{aligned} 0 < t^n \leq \bar{t}^n, \sigma L_j y(t) > 0 & \text{ in } [0, \bar{t}^n], j \in \{0, 1, \dots, n-2\}, \\ \sigma L_{n-1} y(t) > 0 & \text{ in } [0, t^n), \sigma L_{n-1} y(t) \equiv 0 & \text{ in } [t^n, \bar{t}^n], \\ \sigma L_j y(t) > 0, \sigma L_{n-1} y < 0 & \text{ in some right neighborhood of } t = \bar{t}^n. \end{aligned}$$

By the same procedure it can be proved that either y is of Type II (s), $s \in \{0, \dots, n-2\}$, or numbers $t^j, j \in \{0, 1, \dots, n-2\}$, exist such that

$$\begin{aligned} \bar{t}^{n-1} < t^{n-2} < \dots < t^0, \quad \sigma L_i y(t^i) = 0, \sigma L_i y > 0 & \text{ in } (t^{i+1}, t^i), \\ \sigma L_m y > 0, \sigma L_k y < 0 & \text{ in } (t^{i+1}, t^i), \\ m \in \{0, 1, \dots, i-1\}, k \in \{i+1, \dots, n-1\}, \end{aligned}$$

and

$$\sigma L_i y < 0, \quad i \in \{0, 1, \dots, n-1\} \text{ in some right neighborhood of } t^0. \quad (14)$$

Thus (13) is valid in this neighborhood and the statement follows by repeating the considerations in the case (13). Note that in the case Type I(τ), $\tau < \infty$, the relations $\lim_{t \rightarrow \tau} L_i y(t) = 0, \quad i = 0, 1, \dots, n-1$, must be valid because y is defined in \mathbb{R}_+ .

Further, let (10) be valid. By the use of (13), (14) we see that the same initial conditions are valid in some $t^*, t^* \in [0, t^0]$, in the previous part of the proof. Thus the statement of the lemma can be proved similarly. \square

Remark. Let $y : [0, \tau) \rightarrow \mathbb{R}, \tau < \infty$, be a noncontinuable solution. Then the statement of Lemma 2 is valid, too, if (a) is changed into

(a') y is of Type I(τ) with the exception of $\lim_{t \rightarrow \tau} L_i y(t) = 0, \quad i = 0, 1, \dots, n-1$, and if Type II (s) is defined only on $[0, \tau)$.

Lemma 3 ([6, Lemma 9.2]). Let $c_0 \geq 0, t_0 \in I \subset \mathbb{R}_+, h \in L_{loc}(I), h \geq 0, \omega \in C^0(\mathbb{R}_+), \omega(x) > 0$ for $x > c_0, \int_{c_0}^{\infty} \frac{ds}{\omega(s)} < \infty$. Then for every continuous function $x(t) : I \rightarrow \mathbb{R}_+$ which satisfies

$$x(t) \leq c_0 + \left[\int_{t_0}^t h(\tau) \omega(x(\tau)) d\tau \right] \text{sign}(t - t_0), \quad t \in I,$$

we have

$$x(t) \leq \Omega^{-1} \left(\left| \int_{t_0}^t h(\tau) d\tau \right| \right), \quad t \in I,$$

where Ω^{-1} is the inverse function of $\Omega(s) = \int_{c_0}^s \frac{d\tau}{\omega(\tau)}$.

Lemma 4. *Let (7) hold. Then there exists no singular solution of (1) of the second kind.*

The lemma can be proved analogously to Lemma 4 in [7].

Lemma 5 (see [7], **Lemma 1.5 and Consequence 1.2**). *Let $\omega : (0, \infty) \rightarrow \mathbb{R}_+$ be continuous, nondecreasing and $h \in L_{loc}(\mathbb{R}_+)$, $h \geq 0$, such that*

$$\int_0^\infty h(t) dt = \infty, \quad \int_0^1 \frac{dx}{\omega(x)} < \infty.$$

Then the differential inequality $u' + a(t)\omega(u) \leq 0$ has no proper positive solution in \mathbb{R}_+ .

Lemma 6. *Let (5) be valid and one of the following conditions hold:*

- (a) $\lambda = 1$, (6) and (12) hold
- (b) $\lambda \in (0, 1)$, (11) holds.

Then there exists no solution of (1) of Type II(i), $i = 0, 1, \dots, n - 1$.

Proof. (a) With respect to (6) no solution of (1) of Type II(i), $i = 0, 1, \dots, n - 2$, exists (see [3]). The fact that there exists no solution of Type II ($n - 1$) is proved by Chanturia [7] in the proof of Theorem 3.5.

(b) We prove indirectly that a solution of Type II(s), $s \in \{0, 1, \dots, n - 1\}$, does not exist. Thus suppose, without loss of generality, that a solution of (1) $y : \mathbb{R}_+ \rightarrow \mathbb{R}$ exists such that $T \in \mathbb{R}_+$,

$$\begin{aligned} L_i y(t) &\geq 0, \quad i = 0, 1, \dots, s; \quad L_j y(t) \leq 0, \quad j = s + 1, \dots, n - 1, \\ L_m y(t) &\neq 0, \quad m = 0, 1, \dots, n - 2, \quad t \geq T. \end{aligned} \tag{15}$$

Then according to Lemma 1 and (3)

$$\begin{aligned} |L_i y| &\text{ is nondecreasing for } i \in \{0, 1, \dots, n - 1\}, \quad i \neq s, \\ L_s y &\text{ is nonincreasing in } [T, \infty). \end{aligned} \tag{16}$$

Further, by the use of (2), (5), (15), (16)

$$\begin{aligned} |L_i y(t)| &\geq \int_T^\infty a_{i+1}(s) |L_{i+1} y(s)| ds, \quad i = 0, 1, \dots, n - 2, \\ |L_{n-1} y(t)| &\geq \int_T^\infty |L_n y(s)| ds \geq \int_T^\infty a_n(s) |L_0 y(s)|^\lambda ds, \\ -(L_s y(t))' &= a_{s+1}(t) |L_{s+1} y(t)| \quad \text{for } s \in \{0, 1, \dots, n - 2\}, \\ -(L_s y(t))' &= -L_n y(t) \geq a_n(t) (L_0 y(t))^\lambda \quad \text{for } s = n - 1. \end{aligned} \tag{17}$$

From this and (17) we have for $t \in [T, \infty)$

$$\begin{aligned}
& |L_{s+1}y(t)| \geq \\
& \geq \int_T^t a_{s+2}(\tau_{s+2}) \int_T^{\tau_{s+2}} a_{s+3}(\tau_{s+3}) \cdots \int_T^{\tau_{n-2}} a_{n-1}(\tau_{n-1}) |L_{n-1}y(\tau_{n-1})| \geq \\
& \geq \int_T^t a_{s+2}(\tau_{i+2}) \cdots \int_T^{\tau_{n-2}} a_{n-1}(\tau_{n-1}) \int_T^{\tau_{n-1}} a_n(\tau_n) \times \\
& \times \left[\int_T^{\tau_n} a_1(s_1) \int_T^{\tau_1} \cdots \int_T^{\tau_{s-1}} a_s(\tau_s) L_s y(\tau_s) \right]^\lambda d\tau_s \dots d\tau_1 d\tau_n \dots d\tau_{s+2} \leq \\
& \leq Z_s(t, T) (L_s y(t))^\lambda, \quad s = 0, 1, \dots, n-2, \\
& |L_0 y(t)| \geq Z_{n-1}(t, T) L_{n-1} y(t) \quad (\text{for } s = n-1),
\end{aligned}$$

where

$$\begin{aligned}
Z_s(t, T) &= \int_T^t a_{s+2}(\tau_{i+2}) \cdots \int_T^{\tau_{n-1}} a_n(\tau_n) \left[\int_T^{\tau_n} a_1(\tau_1) \cdots \right. \\
&\cdots \left. \int_T^{\tau_{s-1}} a_s(\tau_s) d\tau_s \dots d\tau_1 \right]^\lambda d\tau_n d\tau_{s+2}, \quad s = 0, 1, \dots, n-2, \\
Z_{n-1}(t, T) &= \int_T^t a_1(\tau_1) \int_T^{\tau_1} a_2(\tau_2) \cdots \int_T^{\tau_{n-2}} a_{n-1}(\tau_{n-1}) d\tau_{n-1} \dots d\tau_1 \\
&\quad (\text{for } s = n-1).
\end{aligned}$$

It follows from (17) that

$$(L_s y(t))' + a_{s+1}(t) Z_s^\beta(t, T) (L_s y(t))^\lambda \leq 0, \quad t \in [T, \infty),$$

where $\beta = 1$ for $s \in \{0, 1, \dots, n-2\}$, $\beta = \lambda$ for $s = n-1$.

As according to (11)

$$Z_s(\infty, T) = Z_s(\infty, 0) = \infty,$$

we get the contradiction to Lemma 5 if $L_s y(t) > 0$ in $[T, \infty)$. Thus with respect to (17)

$$s = n-1, \quad L_{n-1} y(t) \equiv 0 \quad \text{on } [\tau, \infty), \quad \tau \in [T, \infty),$$

is the last case which has to be considered. In that case, according to (17), (16),

$$\begin{aligned} 0 &= -(L_{n-1}y(t))' \geq a_n(t)(L_0y(t))^\lambda, \\ a_n(t) &= 0 \quad \text{for almost all } t \in [\tau, \infty). \end{aligned}$$

The contradiction to (11), $i = n - 1$, proves the statement of the lemma. \square

Remark.

(a) The idea of the proof (b) is due to Kiguradze [5] (for the n th-order differential equation); see [7], too.

(b) In [7] sufficient conditions for equation (1) to have Property A are given. For example, (1) has Property A if (5), (6), $\lambda = 1$,

$$\begin{aligned} &\limsup_{t \rightarrow \infty} \frac{I^{n-i}(t, a_{n-1}, \dots, a_i)}{I^{n-i-1}(t, a_{n-1}, \dots, a_{i+1})} \times \\ &\times \int_t^\infty \frac{I^{n-i-1}(s, a_{n-1}, \dots, a_{i+1})I^i(s, a_1, \dots, a_i)}{I^1(s, a_i)} a_n(s) ds > 1 \end{aligned} \quad (18)$$

for $i = 1, 2, \dots, n - 1$, $2|(i + n)$ and $\int_0^\infty I^{n-1}(t, a_{n-1}, \dots, a_1)a_n(t) dt = \infty$ holds.

It is evident that if (1) has Property A then solutions of Type II (i), $i = 0, 1, \dots, n - 1$, do not exist. Condition (12) is the same as (19) for $i = n - 1$. Assumptions of Lemma 6 are weaker see the following example. A similar situation exists for $0 < \lambda < 1$. Moreover, in [7] an extra assumption is made in this case.

Example. Consider equation (1) with (5) where $n = 6$, $a_0 = a_1 = a_2 = a_3 = a_4 = 1$, $a_5 = \frac{1}{t+1}$, $a_6 = \frac{1}{(t+1)^5}$. Then condition (11) is true, but (19) is not true for $i = 3$. Thus solutions of Type II(i), $i = 0, 1, \dots, 5$, do not exist; at the same time the above results of (5) do not guarantee Property A for (1).

Lemma 7. *Let (6) hold and functions $a_n \in L_{loc}(\mathbb{R}_+)$, $g \in C^0(\mathbb{R}_+)$ exist such that $g(0) = 0$, $g(x) > 0$ for $x > 0$, g is nondecreasing, $\int_0^\infty a_n(t) dt = \infty$, and*

$$a_n(t)g(|x_1|) \leq |f(t, x_1, \dots, x_n)| \quad \text{in } \mathbb{D}.$$

Then there exists no solution of (1) of Type II (i), $i = 0, 1, \dots, n - 1$.

Proof. According to [3] and (6) no solution of Type II (i), $i = 0, 1, \dots, n-2$, exists. Let y be a solution of (1) of Type II ($n-1$). Then according to Lemma 1 $|L_{n-1}y|$ is nonincreasing and

$$\begin{aligned} \infty &> |L_{n-1}y(\infty) - L_{n-1}y(T)| = \int_T^\infty |L_n y(s)| ds \geq \\ &\geq \int_T^\infty a_n(t)g(|L_0 y(s)|) ds \geq g(|L_0 y(T)|) \int_T^\infty a_n(s) ds = \infty. \end{aligned}$$

The contradiction proves the lemma. \square

Lemma 8. *Let (8) be valid. Then there exists no singular solution of (1) of the first kind.*

Proof. Let on the contrary a solution y of (1) of the first kind exist. Then numbers $\tau, \tau_1 \in \mathbb{R}_+$, $\tau_1 < \tau$, exist such that

$$\begin{aligned} \varrho(\tau_1) > 0, L_i y \equiv 0 \quad \text{on} \quad [\tau, \infty), \quad i = 0, 1, \dots, n-1, \\ \text{where} \quad \varrho(t) = \sum_{i=0}^{n-1} |L_i y(t)|. \end{aligned} \tag{19}$$

Then by the use of (2) and (8)

$$\begin{aligned} |L_i y(t)| &\leq \int_t^\tau a_{i+1}(s) |L_{i+1} y(s)| ds, \quad i = 0, 1, \dots, n-2, \\ |L_{n-1} y(t)| &\leq \int_t^\tau |L_n y(s)| ds, \\ |L_i y(t)| &\leq \\ &\leq \int_t^\tau a_{i+1}(s_{i+1}) \int_{s_{i+1}}^\tau a_{i+2} \cdots \int_{s_{n-2}}^\tau a_{n-1}(s_{n-1}) \int_{s_{n-1}}^\tau |L_n y(s_n)| ds_n \dots ds_{i+1} \leq \\ &\leq \left[\prod_{j=i+1}^{n-1} \int_{\tau_1}^\tau a_j(s) ds \right] \int_t^\tau |L_n y(s)| ds, \quad i = 0, 1, \dots, n-2, \\ \varrho(t) &\leq C \int_t^\tau |L_n y(s)| ds \leq C \int_t^\tau A(s)g(\varrho(s)) ds, \quad t \in [\tau_1, \tau], \end{aligned}$$

where

$$C = \sum_{i=0}^{n-2} \prod_{j=i+1}^{n-1} \int_{\tau_1}^{\tau} a_j(s) ds + 1.$$

Then it follows from Lemma 3 that

$$\int_0^{\varrho(\tau_1)} \frac{ds}{g(s)} \leq C \int_{\tau_1}^{\tau} A(s) ds < \infty,$$

which contradicts (8) and (19). \square

Lemma 9. *Let y be a solution of (1) defined in \mathbb{R}_+ that satisfies the initial conditions (10). Let (9) be valid. Then y is not of Type $I(\tau)$ for $\tau < \infty$.*

Proof. For $n = 3, 4$ the statement follows from [8] and [9]. Let $n > 4$. Let on the contrary a solution y of Type $I(\tau)$, $\tau < \infty$ exist. It follows from the assumptions of the lemma that an interval $\Lambda = [\tau_1, \tau]$, $\tau_1 < \tau$, exists, for which we have

$$\frac{\max_{t \in \Lambda} a_e \cdot \max_{t \in \Lambda} a_{e+1}}{\min_{t \in \Lambda} a_e \cdot \min_{t \in \Lambda} a_{e+1}} \leq \frac{5}{4}, a_{e+1}(t)a_{e+2}(t) + [a'_{e+1}(t)]_- \int_{\Lambda} a_{e+2}(s) ds > 0,$$

$$a_{e+2}(t)a_{e+3}(t) + [a'_{e+2}(t)]_- \int_{\Lambda} a_{e+3}(s) ds > 0,$$

where $[g(t)]_- = \min(0, g(t))$.

Use the same notation as in the definition of Type $I(\tau)$. According to $\lim_{t \rightarrow \tau} L_e y(t) = 0$ there exists $k_0 \in \mathbb{N}$ such that

$$|L_e y(t_{k_0}^{e+1})| > |L_e y(t_{k_0+1}^{e+1})| > 0, t_{k_0}^{e+1} > \tau_1. \tag{21}$$

Denote $t_{k_0}^{e+1} = t_1$, $t_{k_0}^e = t_2$, $t_{k_0}^{e-1} = t_3$, $\Lambda_1 = t_2 - t_1$, $\Lambda_2 = t_3 - t_2$. Then it follows from (21) and from the definition of Type $I(\tau)$ that (we choose

$L_{e-1}(t_2) > 0$ for simplicity)

$$\begin{aligned}
&L_{e-1}y > 0 \text{ in } [t_1, t_3), \quad L_{e-1}y(t_3) = 0, \\
&L_{e-1} \text{ is increasing (decreasing) in } [t_1, t_2] \text{ (in } [t_2, t_3]), \\
&L_e y > 0 \text{ in } [t_1, t_2), \quad L_e y(t_2) = 0, \quad L_e y < 0 \text{ in } (t_2, t_3], \\
&L_e \text{ is decreasing in } [t_0, t_3], \\
&L_{e+1}y(t_1) = 0, \quad L_{e+1}y < 0 \text{ in } (t_1, t_3], \\
&L_{e+1}y \text{ is decreasing in } [t_0, t_3] \\
&L_{e+j}y < 0 \text{ and } L_{e+j}y \text{ is decreasing in } [t_0, t_3], \quad j = 2, 3.
\end{aligned} \tag{22}$$

From this and (21), (22)

$$L_e y(t_1) > |L_e y(t_3)|, \tag{23}$$

$$\begin{aligned}
L_{e+1}y(t) &= \int_{t_1}^t a_{e+2}(s)L_{e+2}y(s) ds \geq L_{e+2}y(t) \int_{\Lambda} a_{e+2}(s) ds, \quad t \in [t_1, t_3], \\
[L_e y(t)]'' &= [a_{e+1}L_{e+1}y(t)]' = a_{e+1}(t)a_{e+2}(t)L_{e+2}y(t) + \\
&\quad + a'_{e+1}(t)L_{e+1}y(t) \leq a_{e+1}(t)a_{e+2}(t)L_{e+2}y(t) + \\
&\quad + [a'_{e+2}(t)]L_{e+1}y(t) \leq L_{e+2}y(t)[a_{e+1}(t)a_{e+2}(t) + [a'_{e+2}(t)] - \\
&\quad - \int_{\Lambda} a_{e+2}(s) ds] < 0, \quad t \in [t_1, t_3].
\end{aligned} \tag{24}$$

Thus

$$L_e y \text{ is concave in } [t_1, t_3]. \tag{25}$$

We can prove similarly that

$$L_{e+1}y \text{ is concave in } [t_1, t_3]. \tag{26}$$

Further, by the use of (23), (25)

$$\begin{aligned}
L_{e-1}y(t_2) &= \int_{t_2}^{t_3} a_e(s)|L_e y(s)| ds \leq \max_{s \in \Lambda} a_e(s)|L_e y(t_3)| \frac{\Lambda_2}{2}, \\
L_{e-1}y(t_2) &\geq L_{e-1}y(t_2) - L_{e-1}y(t_1) = \int_{t_1}^{t_2} a_e(s)L_e y(s) ds \geq \\
&\geq \min_{s \in \Lambda} a_e(s)L_e y(t_1) \frac{\Lambda_1}{2}.
\end{aligned}$$

Thus, according to (24)

$$1 \leq \frac{|L_e y(t_3)|}{L_e y(t_1)} \frac{\max_{s \in \Lambda} a_e(s)}{\min_{s \in \Lambda} a_e(s)} \frac{\Lambda_2}{\Lambda_1} < \frac{\max_{s \in \Lambda} a_e(s)}{\min_{s \in \Lambda} a_e(s)} \frac{\Lambda_2}{\Lambda_1}. \tag{27}$$

According to (23), (26)

$$L_e(t_1) = \int_{t_1}^{t_2} a_{e+1}(s) |L_{e+1} y(s)| ds \leq |L_{e+1} y(t_2)| \frac{\Lambda_1}{2} \max_{s \in \Lambda} a_{e+1}(s),$$

$$|L_e(t_3)| = \int_{t_2}^{t_3} a_{e+1}(s) |L_{e+1} y(s)| ds \geq |L_{e+1} y(t_2)| \Lambda_2 \min_{s \in \Lambda} a_{e+1}(s).$$

Thus, according to (24), (27) and (23)

$$1 < \frac{\Lambda_1}{2\Lambda_2} \frac{\max_{s \in \Lambda} a_{e+1}(s)}{\min_{s \in \Lambda} a_{e+1}(s)} \leq \frac{1}{2} \frac{\max_{s \in \Lambda} a_{e+1}(s) \max_{s \in \Lambda} a_e(s)}{\min_{s \in \Lambda} a_{e+1}(s) \min_{s \in \Lambda} a_e(s)} \leq \frac{5}{8}.$$

The contradiction proves the statement of the lemma. \square

Proof of Theorem 1. According to Lemmas 2, 6, and 9 y is of Type I(∞) and by the use of Lemma 4 it is proper. \square

Proof of Theorem 2. The statement is a consequence of Lemmas 2, 4, 6, 8, and 9. \square

Proof of Theorem 3. It follows from Lemmas 4, 8, and 9 that y is proper and according to Lemma 7 it is of Type I(∞). \square

ACKNOWLEDGEMENTS

M. Bartušek was partially supported by grant 201/93/0452 of the Czech Grant Agency.

REFERENCES

1. M. Bartušek and Z. Došla, Oscillatory criteria for nonlinear third-order differential equations with quasiderivatives. *Diff. Eq. and Dynam. Systems (to appear)*.
2. T. Kusano and W. F. Trench, Global existence of nonoscillatory solutions of perturbed general disconjugate equations. *Hiroshima Math. J.* **17**(1987), 415–431.
3. M. Švec, Behaviour of nonoscillatory solutions of some nonlinear differential equations. *Acta Math. Univ. Comenian.* **XXXIX**(1980), 115–129.

4. M. Bartušek, Asymptotic properties of oscillatory solutions of differential equations of the n th order. *Folia Fak. Sci. Natur. Univ. Masarykiana Brun. Math.* **3**(1992).
5. I. T. Kiguradze, Some singular boundary-value problems for ordinary differential equations. (Russian) *Tbilisi University Press, Tbilisi*, 1975.
6. I. T. Kiguradze and T. A. Chanturia, Asymptotic properties of solutions of nonautonomous ordinary differential equations. (Russian) *Nauka, Moscow*, 1990; *English translation: Kluwer Academic Publishers, Dordrecht*, 1993.
7. T. A. Chanturia, On oscillatory properties of a system of nonlinear ordinary differential equations. (Russian) *Proc. I. Vekua Inst. Appl. Math. (Tbiliss. Gos. Univ. Inst. Prikl. Math. Trudy)* **14**(1983), 163–204.
8. M. Bartušek, On the structure of solutions of a system of three differential inequalities. *Arch. Math.* **30**(1994), No. 2, 117–130.
9. M. Bartušek, On the structure of solutions of a system of four differential inequalities. *Georgian Math. J.* **2**(1995), No. 3, 225–236.

(Received 20.11.1994)

Author's address:
Faculty of Sciences
Masaryk University
Janačkovo nám. 2a, 662 95 Brno
Czech Republic