

**UPPER ESTIMATE OF THE INTERVAL OF EXISTENCE  
OF SOLUTIONS OF A NONLINEAR TIMOSHENKO  
EQUATION**

DRUMI BAINOV AND EMIL MINCHEV

ABSTRACT. Solutions to the initial-boundary value problem for a nonlinear Timoshenko equation are considered. Conditions on the initial data and nonlinear term are given so that solutions to the problem under consideration do not exist for all  $t > 0$ . An upper estimate of the  $t$ -interval of the existence of solutions is obtained. An estimate of the growth rate of the solutions is given.

1. INTRODUCTION

Let  $\Omega \subset \mathbb{R}^n$  be a bounded domain with boundary  $\partial\Omega$  and  $\bar{\Omega} = \Omega \cup \partial\Omega$ . It is assumed that the divergence theorem can be applied on  $\Omega$ . Let

$$L^q(\Omega) = \left\{ u: \|u\|_{q,\Omega} = \left( \int_{\Omega} |u(x)|^q dx \right)^{1/q} < +\infty \right\}.$$

Denote by  $\bar{u}$  the complex conjugate of  $u$ .

We consider the initial-boundary value problem (IBVP) for the nonlinear Timoshenko equation

$$u_{tt} - \varphi(\|\nabla u\|_{2,\Omega}^2)\Delta u + \Delta^2 u = |u|^{p-1}u, \quad t \geq 0, \quad x \in \Omega, \quad (1)$$

$$u(0, x) = u_0(x), \quad x \in \bar{\Omega}, \quad (2)$$

$$u_t(0, x) = u_1(x), \quad x \in \bar{\Omega}, \quad (3)$$

$$u(t, x) \Big|_{x \in \partial\Omega} = 0, \quad t \geq 0, \quad (4)$$

$$\Delta u(t, x) \Big|_{x \in \partial\Omega} = 0, \quad t \geq 0, \quad (5)$$

---

1991 *Mathematics Subject Classification.* 35B30, 35B35.

*Key words and phrases.* Nonlinear Timoshenko equation, initial-boundary problem, interval of existence, estimate of growth rate of solutions.

where  $p > 1$  is a constant,  $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ ,  $\mathbb{R}_+ = [0, +\infty)$ ,  $u_0, u_1: \overline{\Omega} \rightarrow \mathbb{C}$  are given functions.

Equations of type (1) appear in a variety of models describing nonlinear vibration of beams. This kind of equations is the object of long-standing interest. We should mention the papers [1]–[6], as well as the monograph [7].

In the present paper we investigate the blowing up of solutions [8] of the initial boundary value problem for the nonlinear Timoshenko equation.

## 2. MAIN RESULTS

**Theorem 1.** *Let the following conditions hold:*

1.  $u$  is a suitably smooth classical solution of the IBVP (1)–(5);
2.  $\varphi \in C(\mathbb{R}_+, \mathbb{R}_+)$  and  $m\Phi(s) \geq s\varphi(s)$ ,  $\forall s \geq 0$ , where  $\Phi(s) = \int_0^s \varphi(k) dk$ ,  
 $m \geq 1$ ;
3.  $\operatorname{Re} \int_{\Omega} u_0 \overline{u_1} dx > 0$ ;
4.  $E(0) = \|u_1\|_{2,\Omega}^2 + \Phi(\|\nabla u_0\|_{2,\Omega}^2) + \|\Delta u_0\|_{2,\Omega}^2 - \frac{2}{p+1} \|u_0\|_{p+1,\Omega}^{p+1} \leq 0$ ;
5.  $p > 2m - 1$ .

Then the maximal time interval of the existence  $[0, T_{\max})$  of  $u$  can be estimated by

$$T_{\max} \leq T_0 = \int_{\|u_0\|_{2,\Omega}^2}^{\infty} \left[ C_1 - 4mE(0)\xi + \frac{4C}{p+3} \xi^{\frac{p+3}{2}} \right]^{-\frac{1}{2}} d\xi < +\infty,$$

where

$$C_1 = \left[ 2 \operatorname{Re} \int_{\Omega} u_0 \overline{u_1} dx \right]^2 + 4mE(0)\|u_0\|_{2,\Omega}^2 - \frac{4C}{p+3} \|u_0\|_{2,\Omega}^{p+3},$$

$$C = 2 \frac{p+1-2m}{p+1} \left( \frac{1}{\operatorname{mes} \Omega} \right)^{\frac{p-1}{2}}$$

and if  $T_{\max} = T_0$ , then

$$\lim_{t \rightarrow T_0^-} \|u(t)\|_{2,\Omega}^2 = +\infty$$

and

$$\lim_{t \rightarrow T_0^-} (T_0 - t) \|u(t)\|_{2,\Omega}^{\frac{p-1}{2}} \leq \frac{2}{p-1} \sqrt{\frac{p+3}{C}}.$$

*Proof.* We multiply both sides of equation (1) by  $\bar{u}$ , take the real parts of both sides, and integrate to obtain

$$\frac{1}{2} \frac{d^2}{dt^2} \left( \|u\|_{2,\Omega}^2 \right) - \|u_t\|_{2,\Omega}^2 + \varphi(\|\nabla u\|_{2,\Omega}^2) \|\nabla u\|_{2,\Omega}^2 + \|\Delta u\|_{2,\Omega}^2 = \|u\|_{p+1,\Omega}^{p+1}. \quad (6)$$

On the other hand, if we multiply both sides of equation (1) by  $\bar{u}_t$ , take the real parts of both sides, and integrate, we get the energy identity

$$\|u_t\|_{2,\Omega}^2 + \Phi(\|\nabla u\|_{2,\Omega}^2) + \|\Delta u\|_{2,\Omega}^2 - \frac{2}{p+1} \|u\|_{p+1,\Omega}^{p+1} = E(0). \quad (7)$$

Thus, from (6), (7) and condition 2 of the theorem we conclude that

$$\begin{aligned} \frac{1}{2} \frac{d^2}{dt^2} \left( \|u\|_{2,\Omega}^2 \right) &\geq \|u_t\|_{2,\Omega}^2 - \|\Delta u\|_{2,\Omega}^2 + \|u\|_{p+1,\Omega}^{p+1} + m\|u_t\|_{2,\Omega}^2 + \\ &\quad + m\|\Delta u\|_{2,\Omega}^2 - \frac{2m}{p+1} \|u\|_{p+1,\Omega}^{p+1} - mE(0) \geq \\ &\geq -mE(0) + \frac{p+1-2m}{p+1} \|u\|_{p+1,\Omega}^{p+1} \geq \\ &\geq -mE(0) + \frac{p+1-2m}{p+1} \left( \frac{1}{\text{mes } \Omega} \right)^{\frac{p-1}{2}} \|u\|_{2,\Omega}^{p+1}, \end{aligned}$$

where we have used the Hölder inequality in the last step.

Denote  $X(t) = \|u(t, \cdot)\|_{2,\Omega}^2$ . Then we have the differential inequality

$$\frac{d^2 X}{dt^2} \geq -2mE(0) + CX^{\frac{p+1}{2}}, \quad (8)$$

and

$$X(0) = \|u_0\|_{2,\Omega}^2, \quad \frac{dX}{dt}(0) = 2 \operatorname{Re} \int_{\Omega} u_0 \bar{u}_1 dx.$$

It is easy to prove that  $X'(t) > 0$  wherever  $X(t)$  exists. If this conclusion is false then the set  $Q = \{t: X'(t) \leq 0\}$  is nonempty. Denote  $t_0 = \inf Q$ . By integrating, we obtain

$$0 = X'(t_0) \geq X'(0) - 2mE(0)t_0 + C \int_0^{t_0} X^{\frac{p+1}{2}}(\tau) d\tau > 0,$$

which is a contradiction. We multiply both sides of (8) by  $X'(t)$  and integrating twice we conclude that

$$T_{\max} \leq T_0 = \int_{\|u_0\|_{2,\Omega}^2}^{\infty} \left[ C_1 - 4mE(0)\xi + \frac{4C}{p+3} \xi^{\frac{p+3}{2}} \right]^{-\frac{1}{2}} d\xi < +\infty.$$

If  $T_{\max} = T_0$  then  $\lim_{t \rightarrow T_0^-} \|u(t)\|_{2,\Omega}^2 = +\infty$  and from (8) we have

$$\begin{aligned} T_0 - t &\leq \int_{X(t)}^{\infty} \left[ C_1 - 4mE(0)\xi + \frac{4C}{p+3} \xi^{\frac{p+3}{2}} \right]^{-\frac{1}{2}} d\xi \leq \\ &\leq \sqrt{\frac{p+3}{4C}} \frac{4}{p-1} \frac{1}{[X(t)]^{\frac{p-1}{4}}}, \quad t \in [T_0 - \varepsilon, T_0), \end{aligned}$$

where  $\varepsilon > 0$  is sufficiently small. Thus we obtain

$$\lim_{t \rightarrow T_0^-} (T_0 - t) \|u(t)\|_{2,\Omega}^{\frac{p-1}{2}} \leq \frac{2}{p-1} \sqrt{\frac{p+3}{C}}. \quad \square$$

*Remark 1.* When  $\varphi(s) = 1 + s$  we have  $m = 2$  and  $p > 3$ .

#### ACKNOWLEDGEMENTS

This investigation was partially supported by the Bulgarian Ministry of Education, Science and Technologies under Grant MM-422.

#### REFERENCES

1. M. A. Astaburuaga, C. Fernandez, and G. Perla Menzala, Local smoothing effects for a nonlinear Timoshenko type equation. *Nonlinear Analysis* **23**(1994), No. 9, 1091–1103.
2. P. D’Ancona and S. Spagnolo, Nonlinear perturbations of the Kirchhoff equation. *Comm. Pure Appl. Math.* **XLVII**(1994), No. 7, 1005–1029.
3. R. W. Dickey, Free vibrations and dynamic buckling of the extensible beam, *J. Math. Anal. Appl.* **29**(1970), 443–454.
4. L. A. Medeiros, On a new class of nonlinear wave equations. *J. Math. Anal. Appl.* **69**(1979), 252–262.
5. L. A. Medeiros and M. Milla Miranda, Solutions for the equations of nonlinear vibrations in Sobolev spaces of fractionary order. *Math. Apl. Comput.* **6**(1987), No. 3, 257–267.
6. N. Yoshida, Forced oscillations of extensible beams. *SIAM J. Math. Anal.* **16**(1985), No. 2, 211–220.
7. S. Timoshenko, D. H. Young, and W. Weaver, Jr., Vibration Problems in Engineering. *John Wiley, New York*, 1974.
8. H. Fujita, On the blowing-up of solutions of the Cauchy problem for  $u_t = \Delta u + u^{1+\alpha}$ . *J. Fac. Sci. Univ. Tokyo, Sect. IA* **13**(1966), 109–124.

(Received 19.02.1996)

Authors’ address:  
 Medical University of Sofia  
 P.O. Box 45, Sofia 1504, Bulgaria