

A REGULARITY CRITERION FOR SEMIGROUP RINGS

W. BRUNS AND J. GUBELADZE

ABSTRACT. An analogue of the Kunz–Frobenius criterion for the regularity of a local ring in a positive characteristic is established for general commutative semigroup rings.

Let S be a commutative semigroup (we always assume that S contains a neutral element), and K a field. For every $m \in \mathbb{Z}_+$ the assignment $x \mapsto x^m$, $x \in S$, induces a K -endomorphism π_m of the semigroup ring $R = K[S]$. Therefore we can consider R as an R -algebra via π_m , and especially as an R -module. Let $R^{[m]}$ denote R with its R -module structure induced by π_m . If S is finitely generated, then $R^{[m]}$ is obviously a finitely generated R -module.

In this note we want to give a regularity criterion for S in terms of the homological properties of $R^{[m]}$ that is analogous to Kunz’s [1] characterization of regular local rings of a characteristic $p > 0$ in terms of the Frobenius functor. Our criterion, which generalizes the result of Gubeladze [2, 10.2], requires only a mild condition on S and we provide a ‘pure commutative algebraic’ proof. (In [2] the result was stated for seminormal simplicial affine semigroup rings and derived from the main result of [2] that K_1 -regularity implies the regularity for such rings.)

Theorem 1. *Let S be a finitely generated semigroup, K a field, $R = K[S]$, and $m \in \mathbb{Z}_+$, $m > 0$. Suppose that S has no invertible element $\neq 1$ and is generated by irreducible elements. Then the following conditions are equivalent:*

- (a) $R^{[m]}$ has a finite projective dimension;
- (b) $R^{[m]}$ is a free module;
- (c) S is free, in other words, $S \cong \mathbb{Z}_+^n$ for some $n \in \mathbb{Z}_+$.

1991 *Mathematics Subject Classification.* Primary 13D05, 20M25; Secondary 13A35.
Key words and phrases. Kunz–Frobenius criterion, regularity, commutative semigroup rings.

Proof. It is obvious that (c) implies (b) and (b) implies (a). Now assume that (a) is satisfied. We first reduce the problem to a question of local algebra.

Set $T = S \setminus \{1\}$. The ideal $\mathfrak{m} = TR$ is a maximal ideal of R . Indeed, $R/\mathfrak{m} \cong K$. Furthermore the only prime ideal \mathfrak{q} of R such that $\mathfrak{m} = \pi_m^{-1}(\mathfrak{q})$ is \mathfrak{m} . Therefore $\pi_m \otimes R_{\mathfrak{m}}$ is an endomorphism of $R_{\mathfrak{m}}$ that makes $R_{\mathfrak{m}}$ a finitely generated $R_{\mathfrak{m}}$ -module of finite projective dimension. In particular, $R_{\mathfrak{m}}$ has the same depth considered as an $R_{\mathfrak{m}}$ -module via $\pi_m \otimes R_{\mathfrak{m}}$ as it has in its natural $R_{\mathfrak{m}}$ -module structure (for example, see [3, 1.2.26]). The Auslander–Buchsbaum formula [3, 1.3.3] thus implies that $R_{\mathfrak{m}}$ is a finite free module over itself via $\pi_m \otimes R_{\mathfrak{m}}$. The lemma below shows that $R_{\mathfrak{m}}$ is a regular local ring.

Let x_1, \dots, x_n be the irreducible elements of S . We claim that their images in $R_{\mathfrak{m}}$ form a minimal system of generators of the maximal ideal $\mathfrak{m}R_{\mathfrak{m}}$. Indeed, consider a presentation

$$R^r \xrightarrow{\varphi} R^n \xrightarrow{\psi} \mathfrak{m} \rightarrow 0,$$

where the i -th element e_i of the natural basis of R^n is mapped to x_i . We must show that all the entries of the matrix φ are in \mathfrak{m} . Suppose on the contrary that there is a relation

$$a_1x_1 + \dots + a_nx_n = 0$$

with, for example, $a_1 \notin \mathfrak{m}$. Then $a_1 = \alpha_1 + \alpha_2s_2 + \dots + \alpha_us_u$ with $\alpha_i \in K$, $\alpha_i \neq 0$, and $s_2, \dots, s_u \in T$. Writing a_2, \dots, a_n similarly, we see that there are only two possibilities, (i) $x_1 = s_ix_1$ for some i , or (ii) $x_1 = vx_j$ for some $v \in S$ and $j > 0$. Both cases are impossible because x_1 is irreducible.

However, $R_{\mathfrak{m}}$ is a regular local ring. Especially it is a factorial ring, in which the (images of the) x_i are pairwise non-associated prime elements. Therefore all the elements $x_1^{e_1} \cdots x_n^{e_n}$, $e_1, \dots, e_n \in \mathbb{Z}_+$ are pairwise different, and it follows that $S \cong \mathbb{Z}_+^n$. \square

Remark 2. (a) If we omit the hypothesis that S be generated by irreducible elements, then the proof above shows just the following: the sub-semigroup generated by $x_1, \dots, x_n \in S$ such that x_1, \dots, x_n form a minimal system of generators of the ideal $\mathfrak{m}R_{\mathfrak{m}}$ is free of rank n .

(b) One can weaken the hypothesis of the theorem by requiring only that the group S_0 of invertible elements of S be a free abelian group. Then $T = S \setminus S_0$ generates a prime ideal \mathfrak{p} in R , and part (c) of the theorem must be replaced by the condition that $S \cong \mathbb{Z}_+^n \times \mathbb{Z}^q$ for some $n, q \in \mathbb{Z}_+$.

The following lemma is just an abstract version of Herzog's argument [4] characterizing the modules of finite projective dimension in terms of the Frobenius functor.

Lemma 3. *Let R be a Noetherian local ring with maximal ideal \mathfrak{m} . If there exists an endomorphism π of R with $\pi(\mathfrak{m}) \subset \mathfrak{m}^2$ and such that R is a flat R -module via π , then R is a regular local ring.*

Proof. According to the criterion of Auslander–Buchsbaum–Serre [3, 2.2.7] we must show that $k = R/\mathfrak{m}$ has finite projective dimension as an R -module. Write R' for R considered as an R -module via π , and let \mathcal{P} be the functor that takes an R -module M to $M \otimes R'$ considered as an R -module via the identification $R = R'$. We choose a minimal free resolution \mathcal{F} of k ,

$$\mathcal{F}: \cdots \rightarrow F_{i+1} \xrightarrow{\varphi_{i+1}} F_i \rightarrow \cdots \rightarrow F_1 \rightarrow k \rightarrow 0.$$

One has $\mathcal{P}(R) = R$, $\mathcal{P}(F_i) = F_i$, and $\mathcal{P}(\mathcal{F})$ is the complex that we obtain from \mathcal{F} by replacing all entries in its matrices by their images under π . By hypothesis, $\mathcal{P}(\mathcal{F})$ is again exact, and the exactness is preserved by an e -fold iteration of this process. Especially, $\mathcal{P}^e(\mathcal{F})$ is a free resolution of $\mathcal{P}^e(k)$ for all $e > 0$.

Let $x_1, \dots, x_t \in \mathfrak{m}$ be a maximal R -sequence. Then $\bar{R} = R/(x_1, \dots, x_t)$ has projective dimension t , and so $\mathrm{Tor}_i^R(\bar{R}, \mathcal{P}^e(k)) = 0$ for all $i > t$ and $e > 0$. On the other hand, one can compute $\mathrm{Tor}_i^R(\bar{R}, \mathcal{P}^e(k))$ by tensoring $\mathcal{P}^e(\mathcal{F})$ with \bar{R} . Let B_i be the kernel of φ_i . Then for sufficiently large i and all $e > 0$ we have an exact sequence

$$0 \rightarrow \bar{R} \otimes \mathcal{P}^e(B_{i+1}) \rightarrow \bar{R} \otimes F_{i+1} \rightarrow \bar{R} \otimes \mathcal{P}^e(B_i) \rightarrow 0.$$

Since we have chosen a maximal R -sequence, $\mathrm{depth} \bar{R} \otimes F_{i+1} = 0$ if $F_{i+1} \neq 0$. On the other hand, for e sufficiently large, $\mathcal{P}^e(B_{i+1}) \subset \mathfrak{m}^{2^e} \bar{R} \otimes F_{i+1}$ and $\mathcal{P}^e(B_i) \subset \mathfrak{m}^{2^e} \bar{R} \otimes F_i$ have a positive depth or are zero according to [4, Lemma 3.2]. This is a contradiction. \square

ACKNOWLEDGEMENT

Joseph Gubeladze was supported by Alexander von Humboldt Foundation and CRDF grant #GM1–115.

REFERENCES

1. E. Kunz, Characterizations of regular local rings of characteristic p . *Amer. J. Math.* **91**(1969), 772–784.
2. J. Gubeladze, Nontriviality of $SK_1(R[M])$. *J. Pure Appl. Algebra* **104**(1995), 169–190.
3. W. Bruns and J. Herzog, Cohen–Macaulay rings. *Cambridge University Press*, 1993.

4. J. Herzog, Ringe der Charakteristik p und Frobeniusfunktoren. *Math. Z.* **140**(1974), 67–78.

(Received 25.06.1997)

Authors' addresses:

Winfried Bruns
Universität Osnabrück
Fachbereich Mathematik/Informatik
49069 Osnabrück
Germany

Joseph Gubeladze
A. Razmadze Mathematical Institute
Georgian Academy of Sciences
1, M. Aleksidze St., Tbilisi 380093
Georgia