

ON THE REPRESENTATION OF NUMBERS BY THE  
 DIRECT SUMS OF SOME QUATERNARY QUADRATIC  
 FORMS

N. KACHAKHIDZE

**abstract.** The systems of bases are constructed for the spaces of cusp forms  $S_{2m}(\Gamma_0(5), \chi^m)$  and  $S_{2m}(\Gamma_0(13), \chi^m)$  for an arbitrary integer  $m \geq 2$ . Formulas are obtained for the number of representations of a positive integers by the direct sums of some quaternary quadratic forms.

**2000 Mathematics Subject Classification:** 11E25, 11E20, 11F11, 11E45, 11F27.

**Key words and phrases:** Modular form, representation of integers by quadratic forms, space of cusp forms.

Let  $F_{2m}$  denote a direct sum of  $m$  quaternary quadratic forms  $F_2$  with the same positive discriminant  $q$  ( $q$  is prime  $\equiv 1 \pmod{4}$ ). We shall use the notions, notation and some results from [1].

In what follows let

$$Q_{\frac{q-1}{2}m}^{(q)} = Q_{\frac{q-1}{2}m}^{(q)}(y) = Q_{\frac{q-1}{2}m}^{(q)}(y_1, y_2, \dots, y_{(q-1)m}),$$

where

$$Q_{\frac{q-1}{2}m}^{(q)} = \sum_{k=1}^m Q_{\frac{q-1}{2}}^{(q)}(y_{(k-1)(q-1)+1}, \dots, y_{k(q-1)})$$

with

$$Q_{\frac{q-1}{2}}^{(q)}(u_1, \dots, u_{q-1}) = q \sum_{1 \leq i \leq j \leq q-2} u_i u_j + q \sum_{1 \leq i \leq q-2} u_i u_{q-1} + \frac{q-1}{2} u_{q-1}^2;$$

$$h^{(q)} = \left(1, 2, \dots, \frac{q-1}{2}, \frac{q+3}{2}, \frac{q+5}{2}, \dots, q-1, \frac{q(1-q)}{2}\right),$$

$$g^{(q,m)} = (g_1, g_2, \dots, g_{(q-1)m}) = (h^{(q)}, h^{(q)}, \dots, h^{(q)});$$

$y \equiv g^{(q,m)} \pmod{q}$  is to be understood elementwise;

$$\sigma_t^*(n) = \begin{cases} (q^{(t+1)/2} + q^{(t+1)/4} + 1)\sigma_t(n), & \text{if } q \nmid n, \\ (q^{(t+1)/2} + q^{(t+1)/4} + 1)\sigma_t(n) + q^{3(t+1)/4}\sigma_t\left(\frac{n}{q}\right), & \text{if } q|n, \end{cases}$$

where

$$\begin{aligned}\sigma_t(n) &= \sum_{d|n} d^t; \\ \rho_t^*(n) &= q^{(3t+1)/4} \sum_{\delta d=n} \left(\frac{\delta}{q}\right) d^t + (-1)^{(t+1)/2} \sum_{d|n} \left(\frac{d}{q}\right) d^t,\end{aligned}$$

where  $\left(\frac{\cdot}{q}\right)$  is the Jacobi symbol.

1. Let

$$\begin{aligned}F_2 &= x_1^2 + x_2^2 + x_3^2 + x_4^2 + x_1x_2 + x_1x_3 + x_2x_4, \\ \Phi_2 &= x_1^2 + x_2^2 + 2x_3^2 + 2x_4^2 + x_1x_2 + x_1x_3 + x_1x_4 + x_2x_4 + 2x_3x_4.\end{aligned}$$

$F_2$  and  $\Phi_2$  are the only reduced positive quaternary quadratic forms with determinants 5 and 25 respectively [2, pp. 146, 147]. In [3] it is shown, that  $F_{2m}$  is a quadratic form of type  $(2m, 5, \chi^m)$  ( $\chi = \chi(d) = \left(\frac{d}{5}\right)$ ).

To construct a spherical function of order  $\nu$ , we find a linear transformation which reduces the quadratic form to a sum of squares. Further, all possible monomials of order  $\nu$  are constructed of the variables of the quadratic form. The found linear transformation are next applied to these monomials. The obtained polynomial is subjected to the action of the Laplace operator and for the resulting polynomials we choose identically vanishing linear combinations. A linear combination of the corresponding monomials of order  $\nu$  is a spherical function of the same order with respect to the initial quadratic form.

Applying the Jacobi method it is easy to show that the linear transformation

$$\begin{aligned}x_1 &= y_1 - \frac{1}{\sqrt{3}}y_2 - \frac{2}{\sqrt{15}}y_3, & x_2 &= \frac{2}{\sqrt{3}}y_2 + \frac{1}{\sqrt{15}}y_3 - \frac{1}{\sqrt{5}}y_4, \\ x_3 &= \sqrt{\frac{3}{5}}y_3 - \frac{1}{\sqrt{5}}y_4, & x_4 &= \frac{2}{\sqrt{5}}y_4\end{aligned}\tag{1.1}$$

takes  $\Phi_2$  into a sum of squares.

**Lemma 1.** (a)  $\varphi = \varphi(x_1, \dots, x_4) = x_1^2 - 2x_4^2$  is a spherical function of order 2 with respect to  $\Phi_2$ ;

(b)  $\vartheta(\tau; \Phi_2, \varphi) = 4z - 16z^2 + 8z^3 + 32z^4 - 20z^5 + \dots \in S_4(\Gamma_0(5), 1)$ ;

(c)  $\text{ord}(\vartheta(\tau; \Phi_2, \varphi), i\infty, \Gamma_0(5)) = 1$ .

*Proof.* Using transformation (1.1) we get

$$\begin{aligned}\varphi(x_1, \dots, x_4) &= y_1^2 + \frac{1}{3}y_2^2 + \frac{4}{15}y_3^2 - \frac{8}{5}y_4^2 - \frac{2}{\sqrt{3}}y_1y_2 - \frac{4}{\sqrt{15}}y_1y_3 \\ &\quad + \frac{4}{3\sqrt{5}}y_2y_3 = \varphi(y_1, \dots, y_4), \\ \sum_{i=1}^4 \frac{\partial^2 \varphi}{\partial y_i^2} &= 2\left(1 + \frac{1}{3} + \frac{4}{15} - \frac{8}{5}\right) = 0.\end{aligned}$$

Consequently, by definition (see [4], pp. 849, 853),  $\varphi$  is a spherical function of order 2 with respect to  $\Phi_2$  and, by Lemma 3 from [1],  $\vartheta(\tau; \Phi_2, \varphi) \in S_4(\Gamma_0(5), 1)$ .

Having performed suitable calculations we get

$$\begin{aligned} \vartheta(\tau; \Phi_2, \varphi) &= \sum_{n=1}^{\infty} \left( \sum_{\Phi_2=n} x_1^2 - 2x_4^2 \right) z^n \\ &= 4z - 16z^2 + 8z^3 + 32z^4 - 20z^5 + \dots \end{aligned} \quad (1.2)$$

It follows from (1.2) and formula (1.1) of [1] that

$$\text{ord} \left( \vartheta(\tau; \Phi_2, \varphi), i\infty, \Gamma_0(5) \right) = 1. \quad \square$$

**Theorem 2.** *Let  $C = (c_{sr})$  ( $s = 1, 2, 3; r = 1, 2, \dots, m-1$ ) be the matrix whose elements are non-negative integers satisfying the conditions*

$$4c_{1r} + 2c_{2r} + 2c_{3r} = 2m \quad (1.3)$$

$$c_{1r} + c_{2r} = r, \quad c_{1r} > 0 \quad (1.4)$$

Then for  $m \geq 2$  the system of functions

$$\vartheta^{c_{1r}}(\tau; \Phi_2, \varphi) \vartheta^{c_{2r}}(\tau; Q_2^{(5)}, 1, h^{(5)}) \vartheta^{c_{3r}}(\tau, F_2) \quad (r = 1, 2, \dots, m-1) \quad (1.5)$$

is the basis of the space  $S_{2m}(\Gamma_0(5), \chi^m)$ .

*Proof.* By Lemma 1, (1.3) and Lemmas 3, 8, 9 from [1], functions (1.5) are cusp forms of type  $(2m, \Gamma_0(5), \chi^m)$ . By Lemma 1, (1.4) and Lemma 8 from [1], we have

$$\begin{aligned} \text{ord} \left( \vartheta^{c_{1r}}(\tau; \Phi_2, \varphi) \vartheta^{c_{2r}}(\tau; Q_2^{(5)}, 1, h^{(5)}) \vartheta^{c_{3r}}(\tau, F_2), i\infty, \Gamma_0(5) \right) &= r \\ &(r = 1, 2, \dots, m-1). \end{aligned}$$

Functions (1.5) are linearly independent because their orders at the cusp  $i\infty$  are different. Hence the theorem is proved as it is known (see [4], pp. 815, 816) that  $\dim S_{2m}(\Gamma_0(5), \chi^m) = m-1$ .  $\square$

**Corollary 3.** *For  $m = 2$  the function  $\vartheta(\tau; \Phi_2, \varphi)$  and for  $m \geq 3$  the system of functions*

$$\begin{aligned} &\vartheta^r(\tau; \Phi_2, \varphi) \vartheta^{m-2r}(\tau, F_2) \quad (r = 1, 2, \dots, [m/2]), \\ &\vartheta^{m-r}(\tau; \Phi_2, \varphi) \vartheta^{2r-m}(\tau; Q_2^{(5)}, 1, h^{(5)}) \quad (r = [m/2] + 1, \dots, m-1) \end{aligned}$$

is the basis of the space  $S_{2m}(\Gamma_0(5), \chi^m)$ .

By Lemma 6 from [1],  $\vartheta(\tau, F_{2m}) - E(\tau, F_{2m}) \in S_{2m}(\Gamma_0(5), \chi^m)$ , where  $E(\tau, F_{2m})$  is the Eisenstein series (see [4], pp. 875, 877). Hence, by Corollary 3, there are constants  $\alpha$  and  $\alpha_r^{(2m)}$  such that

$$\vartheta(\tau, F_4) = E(\tau, F_4) + \alpha \vartheta(\tau; \Phi_2, \varphi) \quad (1.6)$$

and for  $m \geq 3$

$$\begin{aligned} \vartheta(\tau, F_{2m}) &= E(\tau, F_{2m}) + \sum_{r=1}^{[m/2]} \alpha_r^{(2m)} \vartheta^r(\tau; \Phi_2, \varphi) \vartheta^{m-2r}(\tau, F_2) \\ &+ \sum_{r=[m/2]+1}^{m-1} \alpha_r^{(2m)} \vartheta^{m-r}(\tau; \Phi_2, \varphi) \vartheta^{2r-m}(\tau; Q_2^{(5)}, 1, h^{(5)}). \end{aligned} \quad (1.7)$$

Using the expansions of  $\vartheta(\tau, F_{2m})$ ,  $E(\tau, F_{2m})$  from [3], that of  $\vartheta(\tau; Q_2^{(5)}, 1, h^{(5)})$  from [1], Lemma 1, and equating, in both parts of (1.6) and (1.7), the coefficients of  $z$  and  $z, z^2, \dots, z^{m-1}$  we get the systems of linear equations, whence we find the constants  $\alpha$  and  $\alpha_r^{(2m)}$ . Equating, in both parts of (1.6) and (1.7), the coefficients of  $z^n$ , we obtain formulas for the arithmetical function  $r(n, F_{2m})$ .

The formulas for  $r(n, F_{2m})$  when  $m = 2, 3, \dots, 6$  have the form

$$\begin{aligned} r(n, F_4) &= \frac{20}{13} \sigma_3^*(n) - \frac{25}{13} \sum_{\Phi_2=n} x_1^2 - 2x_4^2, \\ r(n, F_6) &= \frac{5}{67} \rho_5^*(n) + \frac{225}{67} \sum_{\Phi_2 \oplus F_2=n} x_1^2 - 2x_4^2 \\ &- \frac{1250}{67} \sum_{\substack{\Phi_2 \oplus Q_2^{(5)}=n \\ y \equiv h^{(5)} \pmod{5}}} x_1^2 - 2x_4^2, \\ r(n, F_8) &= \frac{120}{13 \cdot 313} \sigma_7^*(n) + \frac{61850}{13 \cdot 313} \sum_{\Phi_2 \oplus F_4=n} x_1^2 - 2x_4^2 \\ &- \frac{545625}{13 \cdot 313} \sum_{\Phi_4=n} (x_1^2 - 2x_4^2)(x_5^2 - 2x_8^2) \\ &- \frac{1550000}{13 \cdot 313} \sum_{\substack{\Phi_2 \oplus Q_4^{(5)}=n \\ y \equiv g^{(5,2)} \pmod{5}}} x_1^2 - 2x_4^2, \\ r(n, F_{10}) &= \frac{25}{191 \cdot 2161} \rho_9^*(n) + \frac{9830500}{191 \cdot 2161} \sum_{\Phi_2 \oplus F_6=n} x_1^2 - 2x_4^2 \\ &- \frac{185896875}{2 \cdot 191 \cdot 2161} \sum_{\Phi_4 \oplus F_2=n} (x_1^2 - 2x_4^2)(x_5^2 - 2x_8^2) \\ &- \frac{596640625}{2 \cdot 191 \cdot 2161} \sum_{\substack{\Phi_4 \oplus Q_2^{(5)}=n \\ y \equiv h^{(5)} \pmod{5}}} (x_1^2 - 2x_4^2)(x_5^2 - 2x_8^2) \\ &- \frac{1228906250}{191 \cdot 2161} \sum_{\substack{\Phi_2 \oplus Q_6^{(5)}=n \\ y \equiv g^{(5,3)} \pmod{5}}} x_1^2 - 2x_4^2, \end{aligned}$$

$$\begin{aligned}
 r(n, F_{12}) &= \frac{20}{601 \cdot 691} \sigma_{11}^*(n) + \frac{12379975}{601 \cdot 691} \sum_{\Phi_2 \oplus F_8 = n} x_1^2 - 2x_4^2 \\
 &\quad - \frac{115155625}{601 \cdot 691} \sum_{\Phi_4 \oplus F_4 = n} (x_1^2 - 2x_4^2)(x_5^2 - 2x_8^2) \\
 &\quad - \frac{497140625}{601 \cdot 691} \sum_{\Phi_6 = n} (x_1^2 - 2x_4^2)(x_5^2 - 2x_8^2)(x_9^2 - 2x_{12}^2) \\
 &\quad - \frac{3886875000}{601 \cdot 691} \sum_{\substack{\Phi_4 \oplus Q_4^{(5)} = n \\ y \equiv g^{(5,2)} \pmod{5}}} (x_1^2 - 2x_4^2)(x_5^2 - 2x_8^2) \\
 &\quad - \frac{7737500000}{601 \cdot 691} \sum_{\substack{\Phi_2 \oplus Q_8^{(5)} = n \\ y \equiv g^{(5,4)} \pmod{5}}} x_1^2 - 2x_4^2.
 \end{aligned}$$

2. Let

$$\begin{aligned}
 F_2 &= x_1^2 + x_2^2 + x_3^2 + 2x_4^2 + x_1x_2 + x_1x_3 + x_2x_4, \\
 F_2^* &= 7x_1^2 + 6x_2^2 + 5x_3^2 + 2x_4^2 - 8x_1x_2 - 7x_1x_3 \\
 &\quad + 2x_1x_4 + 4x_2x_3 - 3x_2x_4 - x_3x_4, \\
 \Phi_2 &= x_1^2 + 2x_2^2 + 2x_3^2 + 4x_4^2 + x_1x_2 + x_1x_4 \\
 &\quad + x_2x_3 + x_2x_4 + 2x_3x_4.
 \end{aligned}$$

$F_2$  is the only reduced positive quaternary quadratic form with determinant 13 ([2], p. 141) and  $F_2^*$  is adjoint to  $F_2$ .  $F_2$  and  $F_2^*$  are quadratic forms of type  $(2, 13, \chi)$  [5], and  $\Phi_2$  is a quadratic form of type  $(2, 13, 1)$  [6], where  $\chi = \chi(d) = \left(\frac{d}{13}\right)$ .

Using Lemma 5 from [1], it is easy to show that  $F_{2m}$  is a quadratic form of type  $(2m, 13, \chi^m)$ .

The following Lemmas are proved exactly as Lemma 1.

**Lemma 4.** (a)  $\varphi_1 = \varphi_1(x_1, \dots, x_4) = x_1^2 - 2x_2^2$ ,  $\varphi_2 = \varphi_2(x_1, \dots, x_4) = x_2^2 - x_3^2$ ,  $\varphi_3 = \varphi_3(x_1, \dots, x_4) = 2x_1x_3 - x_4^2$  are spherical functions of order 2 with respect to  $\Phi_2$ ;

$$\begin{aligned}
 \text{(b) } \vartheta(\tau; \Phi_2, \varphi_1) &= 2z - 6z^2 - 2z^3 + 14z^4 - 10z^5 + 26z^6 \\
 &\quad - 30z^7 - 38z^8 + 56z^9 + 42z^{10} + \dots,
 \end{aligned}$$

$$\vartheta(\tau; \Phi_2, \varphi_2) = 2z^2 - 4z^3 + 2z^5 - 2z^6 + 18z^7 - 14z^8 - 24z^9 - 4z^{10} + \dots,$$

$$\vartheta(\tau; \Phi_2, \varphi_3) = 4z^3 - 4z^4 - 8z^6 - 8z^7 + 16z^8 + 12z^9 - 4z^{10} + \dots;$$

$$\text{(c) } \vartheta(\tau; \Phi_2, \varphi_r) \in S_4(\Gamma_0(13), 1);$$

$$\text{(d) } \text{ord}(\vartheta(\tau; \Phi_2, \varphi_r), i\infty, \Gamma_0(13)) = r \quad (r = 1, 2, 3).$$

**Lemma 5.** (a)  $\varphi_4 = \varphi_4(x_1, \dots, x_8) = x_4^2 - x_5^2$  is a spherical function of order 2 with respect to  $\Phi_2 \oplus F_2^*$ ;

$$(b) \vartheta(\tau; \Phi_2 \oplus F_2^*, \varphi_4) = 8z^4 + 24z^6 - 24z^7 - 24z^8 \\ - 48z^9 - 40z^{10} + \dots \in S_6(\Gamma_0(13), \chi);$$

$$(c) \text{ord}(\vartheta(\tau; \Phi_2 \oplus F_2^*, \varphi_4), i\infty, \Gamma_0(13)) = 4.$$

**Lemma 6.** (a)  $\varphi_5 = \varphi_5(x_1, \dots, x_4) = x_2^4 + x_3^4 - 6x_2^2x_3^2$  and  $\varphi_6 = \varphi_6(x_1, \dots, x_4) = x_1x_4^3 + x_3x_4^3 - 6x_1^2x_3x_4$  are spherical functions of order 4 with respect to  $F_2^*$ ;

$$(b) \vartheta(\tau; F_2^*, \varphi_5) = 8z^5 + 8z^6 - 32z^7 + 8z^8 + 0 \cdot z^{10} + \dots,$$

$$\vartheta(\tau; F_2^*, \varphi_6) = 12z^6 - 12z^7 - 12z^8 + 0 \cdot z^{10} + \dots;$$

$$(c) \vartheta(\tau; F_2^*, \varphi_r) \in S_6(\Gamma_0(13), \chi);$$

$$(d) \text{ord}(\vartheta(\tau; F_2^*, \varphi_r), i\infty, \Gamma_0(13)) = r \quad (r = 5, 6).$$

**Lemma 7.** (a)  $\varphi_7 = \varphi_7(x_1, \dots, x_8) = x_7^4 - x_3^4 + 6x_2^2x_3^2 - 6x_2^2x_7^2$  is a spherical function of order 4 with respect to  $F_4^*$ ;

$$(b) \vartheta(\tau; F_4^*, \varphi_7) = 48z^7 + 96z^9 - 96z^{10} + \dots \in S_8(\Gamma_0(13), 1);$$

$$(c) \text{ord}(\vartheta(\tau; F_4^*, \varphi_7), i\infty, \Gamma_0(13)) = 7.$$

**Theorem 8.** Let

$$\lambda = \begin{cases} 2m + \left\lceil \frac{m}{3} \right\rceil - 1, & \text{if } m \equiv 0 \text{ or } 2 \pmod{3}, \\ \frac{7}{3}(m-1), & \text{if } m \equiv 1 \pmod{3} \end{cases}$$

and let  $C = (c_{sr})$  ( $s = 1, 2, \dots, 9$ ;  $r = 1, 2, \dots, \lambda$ ) be the matrix whose elements are non-negative integers satisfying the conditions

$$4 \sum_{s=1}^8 c_{sr} + 2 \sum_{s=4}^9 c_{sr} + 2c_{7r} = 2m \quad (2.1)$$

$$\sum_{s=1}^7 s \cdot c_{sr} + 7c_{8r} = r \quad (r = 1, 2, \dots, \lambda). \quad (2.2)$$

$$c_{4r} = 0 \text{ or } 1, \quad \sum_{s=1}^7 c_{sr} > 0$$

Further, let  $\varphi^{(r)} = \prod_{s=1}^7 \varphi_s^{c_{sr}}$  denote the direct product of functions  $\varphi_s$ , defined by Lemmas 4–7 ( $\varphi_s^0 \equiv 1$ ). Then for each  $m \geq 2$ , the system of functions

$$\vartheta(\tau; \Phi_{2v_r} \oplus F_{2t_r}^*, \varphi^{(r)}) \vartheta^{c_{8r}}(\tau; Q_6^{(13)}, 1, h^{(13)}) \vartheta^{c_{9r}}(\tau, F_2^*) \quad (2.3) \\ (r = 1, 2, \dots, \lambda),$$

where

$$v_r = \sum_{s=1}^4 c_{sr}, \quad t_r = \sum_{s=4}^7 c_{sr} + c_{7r} \quad (2.4)$$

is the basis of the space  $S_{2m}(\Gamma_0(13), \chi^m)$ .

*Proof.* It follows from the definition of  $\varphi^{(r)}$ , (2.4) and ([1], Lemma 9) that

$$\begin{aligned} \vartheta\left(\tau; \Phi_{2v_r} \oplus F_{2t_r}^*, \varphi^{(r)}\right) &= \vartheta^{c_{1r}}(\tau; \Phi_2, \varphi_1) \\ &\quad \times \vartheta^{c_{2r}}(\tau; \Phi_2, \varphi_2) \vartheta^{c_{3r}}(\tau; \Phi_2, \varphi_3) \\ &\quad \times \vartheta^{c_{4r}}(\tau; \Phi_2 \oplus F_2^*, \varphi_4) \vartheta^{c_{5r}}(\tau; F_2^*, \varphi_5) \\ &\quad \times \vartheta^{c_{6r}}(\tau; F_2^*, \varphi_6) \vartheta^{c_{7r}}(\tau; F_4^*, \varphi_7). \end{aligned} \quad (2.5)$$

Therefore (2.5), (2.1), above Lemmas 4–7 and Lemmas 3, 8, 9 from [1] imply that functions (2.3) are cusp forms of type  $(2m, \Gamma_0(13), \chi^m)$ . By Lemmas 4–7, (2.2) and Lemma 8 from [1], we have

$$\begin{aligned} \text{ord}\left(\vartheta(\tau; \Phi_{2v_r} \oplus F_{2t_r}^*, \varphi^{(r)}) \vartheta^{c_{8r}}(\tau; Q_6^{(13)}, 1, h^{(13)}) \vartheta^{c_{9r}}(\tau, F_2^*), i\infty, \Gamma_0(13)\right) &= r \\ &\quad (r = 1, 2, \dots, \lambda). \end{aligned}$$

Functions (2.3) are linearly independent because their orders at the cusp  $i\infty$  are different. Hence the theorem is proved as it is known that  $\dim S_{2m}(\Gamma_0(13), \chi^m) = \lambda$  (see [4], pp. 815, 816).  $\square$

By ([1], Lemma 6)  $\vartheta(\tau, F_{2m}) - E(\tau, F_{2m}) \in S_{2m}(\Gamma_0(13), \chi^m)$ . Hence by Theorem 8 there are constants  $\alpha_r^{(2m)}$  such that

$$\begin{aligned} \vartheta(\tau, F_{2m}) &= E(\tau, F_{2m}) \\ &+ \sum_{r=1}^{\lambda} \alpha_r^{(2m)} \vartheta\left(\tau; \Phi_{2v_r} \oplus F_{2t_r}^*, \varphi^{(r)}\right) \vartheta^{c_{8r}}(\tau; Q_6^{(13)}, 1, h^{(13)}) \vartheta^{c_{9r}}(\tau, F_2^*). \end{aligned}$$

Using the expansion of  $\vartheta(\tau, F_2)$  from [5] we get the expansions of  $\vartheta(\tau, F_{2m}) = \vartheta^m(\tau, F_2)$ . From formulas (1.5)–(1.9) of [1], when  $q = 13$ , we get the expansions of  $E(\tau, F_{2m})$ . Then using the expansions of  $\vartheta(\tau, F_2^*)$  from [5] and  $\vartheta(\tau; Q_6^{(13)}, 1, h^{(13)})$  from [1], Lemmas 4–7, exactly as above we get formulas for  $r(n, F_{2m})$ . When  $m = 2, \dots, 5$  they have the form

$$\begin{aligned} r(n, F_4) &= \frac{12}{7 \cdot 17} \sigma_3^*(n) + \frac{330}{7 \cdot 17} \sum_{\Phi_2=n} x_1^2 - 2x_2^2 + \frac{1342}{7 \cdot 17} \sum_{\Phi_2=n} x_2^2 - x_3^2 \\ &\quad - \frac{1013}{7 \cdot 17} \sum_{\Phi_2=n} 2x_1x_3 - x_4^2, \\ r(n, F_6) &= \frac{13}{109 \cdot 307} \rho_5^*(n) + \frac{416694}{109 \cdot 307} \sum_{\Phi_2 \oplus F_2^*=n} x_1^2 - 2x_2^2 \\ &\quad + \frac{3425570}{109 \cdot 307} \sum_{\Phi_2 \oplus F_2^*=n} x_2^2 - x_3^2 + \frac{4662503}{109 \cdot 307} \sum_{\Phi_2 \oplus F_2^*=n} 2x_1x_3 - x_4^2 \\ &\quad - \frac{2905512}{109 \cdot 307} \sum_{\Phi_2 \oplus F_2^*=n} x_4^2 - x_5^2 - \frac{1698555}{109 \cdot 307} \sum_{F_2^*=n} x_2^4 + x_3^4 - 6x_2^2x_3^2 \end{aligned}$$

$$\begin{aligned}
& + \frac{2615880}{109 \cdot 307} \sum_{F_2^*=n} x_1 x_4^3 + x_3 x_4^3 - 6x_1^2 x_3 x_4, \\
r(n, F_8) & = \frac{24}{17 \cdot 14281} \sigma_7^*(n) + \frac{5481876}{17 \cdot 14281} \sum_{\Phi_2 \oplus F_4^*=n} x_1^2 - 2x_2^2 \\
& + \frac{83647460}{17 \cdot 14281} \sum_{\Phi_2 \oplus F_4^*=n} x_2^2 - x_3^2 \\
& + \frac{251775638}{17 \cdot 14281} \sum_{\Phi_2 \oplus F_4^*=n} 2x_1 x_3 - x_4^2 \\
& + \frac{239040860}{17 \cdot 14281} \sum_{\Phi_4=n} (2x_1 x_3 - x_4^2)(x_5^2 - 2x_6^2) \\
& + \frac{591029498}{17 \cdot 14281} \sum_{\Phi_4=n} (2x_1 x_3 - x_4^2)(x_6^2 - x_7^2) \\
& + \frac{877045926}{17 \cdot 14281} \sum_{\Phi_4=n} (2x_1 x_3 - x_4^2)(2x_5 x_7 - x_8^2) \\
& - \frac{315282279}{17 \cdot 14281} \sum_{F_4^*=n} x_7^4 - x_3^4 + 6x_2^2 x_3^2 - 6x_2^2 x_7^2, \\
r(n, F_{10}) & = \frac{1}{144547171} \rho_9^*(n) + \frac{4305040872}{144547171} \sum_{\Phi_2 \oplus F_6^*=n} x_1^2 - 2x_2^2 \\
& + \frac{106015990372}{144547171} \sum_{\Phi_2 \oplus F_6^*=n} x_2^2 - x_3^2 \\
& + \frac{541004173576}{144547171} \sum_{\Phi_2 \oplus F_6^*=n} 2x_1 x_3 - x_4^2 \\
& + \frac{1284103896202}{144547171} \sum_{\Phi_4 \oplus F_2^*=n} (x_1^2 - 2x_2^2)(2x_5 x_7 - x_8^2) \\
& + \frac{7148202390955}{144547171} \sum_{\Phi_4 \oplus F_2^*=n} (x_2^2 - x_3^2)(2x_5 x_7 - x_8^2) \\
& + \frac{11816421944265}{144547171} \sum_{\Phi_4 \oplus F_2^*=n} (2x_1 x_3 - x_4^2)(2x_5 x_7 - x_8^2) \\
& + \frac{8100018167088}{144547171} \sum_{\Phi_2 \oplus F_2^*=n} (x_1^2 - 2x_2^2)(x_5 x_8^3 + x_7 x_8^3 - 6x_5^2 x_7 x_8) \\
& + \frac{22777304972068}{144547171} \sum_{\Phi_2 \oplus F_2^*=n} (x_2^2 - x_3^2)(x_5 x_8^3 + x_7 x_8^3 - 6x_5^2 x_7 x_8) \\
& + \frac{41871730633632}{144547171} \sum_{\Phi_2 \oplus F_2^*=n} (2x_1 x_3 - x_4^2)(x_5 x_8^3 + x_7 x_8^3 - 6x_5^2 x_7 x_8)
\end{aligned}$$



$$-\frac{7694378154216}{144547171} \sum_{\substack{\Phi_2 \oplus Q_6^{(13)} = n \\ y \equiv h^{(13)} \pmod{13}}} 2x_1x_3 - x_4^2.$$

## REFERENCES

1. N. KACHAKHIDZE, On the representation of numbers by the direct sums of some binary quadratic forms. *Georgian Math. J.* **5**(1998), No. 1, 55–70.
2. K. GERMANN, Tabellen reduzierter, positiver quaternärer quadratischer formen. *Studien zur Theorie der quadratischen Formen. Birkhäuser Verlag, Basel und Stuttgart*, 1968.
3. N. D. KACHAKHIDZE, On the representation of numbers by a direct sum of quadratic forms of the kind  $x_1^2 + x_2^2 + x_3^2 + x_4^2 + x_1x_2 + x_1x_3 + x_2x_4$ . (Russian) *Proc. Tbilisi Univ. Math. Mekh. Astron.* **299**(1990), 60–85.
4. E. HECKE, *Mathematische Werke. Zweite Aufl. Vandenhoeck und Ruprecht, Göttingen*, 1970.
5. G. A. LOMADZE, On the cusp forms with prime level and of adjacent type. (Russian) *Trudy Tbiliss. Mat. Inst. Razmadze* **63**(1980), 36–53.
6. K. SH. SHAVGULIDZE, On cusp and quadratic forms of type  $(-4, q, 1)$ . (Russian) *Trudy Tbiliss. Univ. Math. Mekh. Astron.* **214**(1980), 194–219.

(Received 20.11.2000)

Author's Address:  
 Georgian Technical University  
 Department of Mathematics No. 3  
 77, Kostava St., Tbilisi 380075  
 Georgia