

## GEOMETRY OF MODULUS SPACES

R. KHALIL, D. HUSSEIN, AND W. AMIN

**Abstract.** Let  $\phi$  be a modulus function, i.e., continuous strictly increasing function on  $[0, \infty)$ , such that  $\phi(0) = 0$ ,  $\phi(1) = 1$ , and  $\phi(x + y) \leq \phi(x) + \phi(y)$  for all  $x, y$  in  $[0, \infty)$ . It is the object of this paper to characterize, for any Banach space  $X$ , extreme points, exposed points, and smooth points of the unit ball of the metric linear space  $\ell^\phi(X)$ , the space of all sequences  $(x_n)$ ,  $x_n \in X$ ,  $n = 1, 2, \dots$ , for which  $\sum \phi(\|x_n\|) < \infty$ . Further, extreme, exposed, and smooth points of the unit ball of the space of bounded linear operators on  $\ell^p$ ,  $0 < p < 1$ , are characterized.

**2000 Mathematics Subject Classification:** Primary: 47B38; Secondary: 48A65.

**Key words and phrases:** Extreme point, exposed point, smooth point, modulus function.

**0. Introduction.** Let  $\phi : [0, \infty) \rightarrow [0, \infty)$  be a continuous function. We call  $\phi$  a modulus function if:

- (i)  $\phi(x) = 0$  if and only if  $x = 0$ ;
- (ii)  $\phi$  is increasing;
- (iii)  $\phi(x + y) \leq \phi(x) + \phi(y)$ .

The functions  $\phi(x) = x^p$ ,  $p \in (0, 1)$ , and  $\phi(x) = \ln(1 + x)$  are modulus functions.

For a modulus function  $\phi$ , we let  $\ell^\phi$  denote the space of all real-valued sequences  $(x_n)$  for which  $\sum \phi(|x_n|) < \infty$ . For  $x, y \in \ell^\phi$ ,  $d(x, y) = \sum \phi(|x_n - y_n|)$  is a metric on  $\ell^\phi$ . For  $x \in \ell^\phi$  we let  $\|x\|_\phi$  denote  $d(x, 0)$ . The space  $(\ell^\phi, \|\cdot\|_\phi)$  is a metric linear space. These spaces were initiated by Ruckle [4].

Throughout this paper,  $R$  denotes the set of real numbers. If  $X$  is a Banach space,  $X^*$  will denote the dual of  $X$ . If  $x^* \in X^*$  and  $x \in X$ , we let  $\langle x^*, x \rangle$  denote the value of  $x^*$  at  $x$ . We let  $\ell^p$  denote the space of all (real) sequences  $(x_n)$  for which  $\sum |x_n|^p < \infty$ ,  $0 < p < \infty$ . For  $x \in \ell^p$ , we let

$$\|x\|_p = \begin{cases} (\sum |x_i|^p)^{\frac{1}{p}} & \text{if } 1 \leq p < \infty, \\ \sum |x_i|^p & \text{if } 0 < p < 1. \end{cases}$$

So  $\|x\|_1$  is the 1-norm of  $x$  in  $\ell^1$ . For  $p = \infty$ ,  $\ell^\infty$  is the space of all bounded (real) sequences. If  $x \in \ell^\infty$ , we let  $\|x\|_\infty = \sup_i |x_i|$ .

Let us summarize the basic properties of  $(\ell^\phi, \|\cdot\|_\phi)$  in

**Theorem A.** *Let  $\phi$  be any modulus function. Then:*

- (1)  $(\ell^\phi, \|\cdot\|_\phi)$  is a complete metric linear space.
- (2) If  $\|x\|_\phi \leq \phi(a)$ , then  $\|x\|_1 \leq a$ .
- (3)  $\ell^\phi \subseteq \ell^1$ , and the inclusion map  $I : \ell^\phi \rightarrow \ell^1$  is continuous.
- (4) If  $\phi(1) = 1$ , then for every  $x \in \ell^\phi$  there exists  $r > 0$  such that  $\|rx\|_\phi = 1$ .
- (5) There exist  $\alpha$  and  $a$  in  $[0, \infty)$  such that  $\phi(x) > \alpha x$  for all  $x \in [0, a)$ .

*Proof.* The proof of (1) is in [4]. Statements (2) and (3) are easy to handle. Statement (5) is in [5]. So we prove only (4).

There are two cases: either  $\|x\|_\phi < 1$  or  $\|x\|_\phi > 1$ . If  $\|x\|_\phi > 1$ , define  $F : [0, 1] \rightarrow [0, \infty)$  by  $F(t) = \|tx\|_\phi$ . Then  $F$  is continuous with  $F(0) = 0$  and  $F(1) > 1$ . By the intermediate value theorem there is  $r \in (0, 1)$  such that  $F(r) = 1$ . Hence  $\|rx\|_\phi = 1$ . The other case follows from statement (2) and the assumption  $\phi(1) = 1$ .  $\square$

Let  $X$  be a Banach space. A linear mapping  $T : \ell^\phi \rightarrow X$  is called bounded if there exists  $\lambda > 0$  such that  $\|Tx\| \leq \lambda$  for all  $x$  in  $\ell^\phi$  for which  $\|x\|_\phi \leq 1$ . We let  $L(\ell^\phi, X)$  denote the space of all bounded linear operators on  $\ell^\phi$  with values in  $X$ . We let  $(\ell^\phi)^*$  denote  $L(\ell^\phi, R)$ . For  $T \in L(\ell^\phi, X)$  we set  $\|T\| = \sup\{\|Tx\| : \|x\|_\phi \leq 1\}$ . For the case  $0 < p < 1$  we let  $B(\ell^p, \ell^p)$  denote the space of linear operators on  $\ell^p$  for which  $\|Tx\|_p \leq \lambda \|x\|_p$  for all  $x \in \ell^p$  with some  $\lambda$  depending on  $T$ . Since  $a|b|^p = |a^{\frac{1}{p}}b|^p$  for  $a > 0$ , it follows that  $\sup\{\|Tx\|_p : \|x\|_p \leq 1\} = \inf\{\lambda : \|Tx\|_p \leq \lambda \|x\|_p \text{ for all } x \in \ell^p\}$ . Hence  $B(\ell^p, \ell^p) = L(\ell^p, \ell^p)$ .

For a modulus function  $\phi$  and a Banach space  $X$ , we set  $\ell^\phi(X) = \{(x_n) : x_n \in X \text{ and } \sum \phi(\|x_n\|) < \infty\}$ . If  $x = (x_n) \in \ell^\phi(X)$ , then we define  $\|x\|_\phi = \sum \phi(\|x_n\|)$ . It is easy to check that  $(\ell^\phi(X), \|\cdot\|_\phi)$  is a complete metric linear space.

Extreme points of the unit ball of  $L(\ell^p, \ell^p)$ ,  $1 < p < \infty$ , have been studied extensively by many authors ([6]–[10] and others). A full characterization of extreme points of the unit ball of  $L(\ell^p, \ell^p)$ ,  $1 < p < \infty$ , is still an open problem.

In this paper we characterize extreme, exposed, and smooth points of the unit balls of  $\ell^\phi$ ,  $\ell^\phi(X)$  and  $L(\ell^p, \ell^p)$ ,  $0 < p < 1$ .

**1. Basic Structure of Spaces  $\ell^\phi(X)$ .** Throughout this paper we will assume that:

- (i)  $\phi$  is strictly increasing;
- (ii)  $\phi(1) = 1$ .

Let  $M$  denote the class of all modulus functions satisfying (i) and (ii). We set  $(\ell^\phi(X))^* = L(\ell^\phi, R)$ , where  $R$  is the set of real numbers.

**Theorem 1.1.** *Let  $\phi \in M$  and  $X$  be any Banach space. Then  $[\ell^\phi(X)]^*$  is isometrically isomorphic to  $\ell^\infty(X^*)$ .*

*Proof.* Let  $F \in \ell^\infty(X^*)$ . So  $F = (x_1^*, x_2^*, \dots)$  with  $x_i^* \in X^*$  and  $\sup_i \|x_i^*\| < \infty$ . Define  $\tilde{F} : \ell^\phi(X) \rightarrow R$  such that for  $x = (x_i) \in \ell^\phi(X)$ ,  $\tilde{F}(x) = \sum \langle x_i, x_i^* \rangle$ .

Hence  $|\tilde{F}(x)| \leq \sum \|x_i\| \|x_i^*\| \leq \|F\| \sum \|x_i\|$ . Now for any function  $\phi$  in  $M$  one can easily show that  $\ell^\phi(X) \subseteq \ell^1(X)$ . Further, if  $\|f\|_\phi = 1$ , then  $\|f\|_1 \leq 1$ . Thus

$$\|\tilde{F}\| \leq \|F\| \tag{*}$$

On the other hand, if  $\tilde{F} \in [\ell^\phi(X)]^*$ , then we define  $x_i^*$  in  $X^*$  as  $x_i^*(x) = \tilde{F}(0, 0, \dots, 0, x, 0, \dots)$  where  $x$  appears in the  $i$ th coordinate. Set  $F = (x_1^*, x_2^*, \dots)$ . Then since  $\sup_i \|x_i^*\| \leq \|\tilde{F}\|$ , we obtain  $F \in \ell^\infty(X^*)$  and  $\|F\|_\infty \leq \|\tilde{F}\|$ . This together with (\*) gives  $\|F\|_\infty = \|\tilde{F}\|$ . Thus the mapping  $J : \ell^\infty(X^*) \rightarrow [\ell^\phi(X)]^*$ ,  $J(F) = \tilde{F}$  is linear onto and an isometry. This ends the proof.  $\square$

As a consequence we get

**Corollary 1.2.**  $(\ell^\phi)^* = \ell^\infty$ .

*Remark 1.* If  $\phi(x+y) < \phi(x) + \phi(y)$  for any  $x > 0, y > 0$ , then there are some elements  $x$  of  $\ell^\phi$  such that there is no  $x^*$  in  $\ell^\infty$  for which  $\langle x, x^* \rangle = \|x\| \|x^*\|$ . Indeed, if  $\|x\|_\phi = 1$ , then the continuity of  $\phi$ , being strictly increasing and  $\phi(1) = 1$ , implies that  $\|x\|_1 = 1$  unless  $x$  has only one nonzero coordinate. So for  $x$  with more than one nonzero terms there cannot exist  $x^*$  in  $\ell^\infty$  which attains its norm at  $x$ . However, if  $x$  has only one nonzero coordinate, then  $\|x\|_1 = \|x\|_\phi$ , if  $\|x\|_\phi = 1$  and such  $x^*$  exists.

**2. Geometry of  $B_1(\ell^\phi(X))$ .** A point  $x$  of a set  $K$  of a metric linear space  $E$  is called extreme if there exist no  $y$  and  $z$  in  $K$  such that  $y \neq z$  and  $x = \frac{1}{2}(y + z)$ . The point  $x$  in  $B_1(E)$  is called exposed if there exists  $f \in B_1(E^*)$  such that  $f(x) = d(x, 0)$ , and  $f(y) < d(y, 0)$  for all  $y$  in  $B_1(E)$ ,  $y \neq x$ . We call  $x$  a smooth point of  $B_1(E)$  if there exists a unique  $f \in B_1(E^*)$  such that  $f(x) = d(x, 0)$ .

In this section we will characterize extreme, exposed, and smooth points of  $B_1(\ell^\phi(X))$  for any Banach space  $X$ .

**Theorem 2.1.** *Let  $\phi \in M$ . The following statements are equivalent:*

- (i)  $f$  is an extreme point of  $B_1(\ell^\phi(X))$ .
- (ii)  $f(n) = 0$  for all  $n$  except for one coordinate, say,  $f(n_0)$ , and  $f(n_0)$  is an extreme point of  $B_1(X)$ .

*Proof.* (i)  $\rightarrow$  (ii). Let  $f$  be extreme and, if possible, assume that  $f$  does not vanish at  $n_1$  and  $n_2$ . Define

$$g(n) = \begin{cases} f(n), & n \neq n_1, n_2, \\ \frac{\|f(n_1)\| + \|f(n_2)\|}{\|f(n_1)\|} f(n_1), & n = n_1, \\ 0, & n = n_2, \end{cases}$$

$$h(n) = \begin{cases} f(n), & n \neq n_1, n_2, \\ \frac{\|f(n_1)\| + \|f(n_2)\|}{\|f(n_2)\|} f(n_2), & n = n_2, \\ 0, & n = n_1. \end{cases}$$

Then  $g \neq h$ . Further,

$$\|g\|_\phi = \sum \phi(\|g(n)\|) \leq \sum \phi\|f(n)\| \leq 1.$$

Similarly,  $\|h\|_\phi \leq 1$ . Now

$$f = \frac{\|f(n_1)\|}{\|f(n_1)\| + \|f(n_2)\|} g + \frac{\|f(n_2)\|}{\|f(n_1)\| + \|f(n_2)\|} h = tg + (1-t)h, \quad 0 < t < 1,$$

where  $t = \frac{\|f(n_2)\|}{\|f(n_1)\| + \|f(n_2)\|}$ .

Hence  $f$  is not an extreme point. Thus  $f$  must be of the form

$$f(n) = \delta_{nn_0} \cdot x_0,$$

where  $\delta_{ij}$  stands for the Kronecker's delta.

Now we claim that  $x_0$  is an extreme point of  $B_1(X)$ . Indeed,  $\|f\|_\phi = 1 = \phi(\|x_0\|)$ . Since  $\phi$  is strictly increasing, we have  $\|x_0\| = 1$ . If  $x_0$  is not an extreme point, then  $x_0 = \frac{1}{2}(y+z)$  for some  $y$  and  $z$  in  $B_1(X)$ . Then one can construct  $f_1$  and  $f_2$  in  $B_1(\ell^\phi(X))$  such that  $f = \frac{1}{2}(f_1 + f_2)$ . Hence  $x_0$  must be extreme.

Conversely: **(ii)**  $\longrightarrow$  **(i)**. Let  $f(n) = \delta_{nn_0} \cdot x$  with  $x$  an extreme point of  $B_1(X)$ . If  $f$  is not extreme, then there exist  $g$  and  $h$  in  $B_1(\ell^\phi(X))$  such that  $f = \frac{1}{2}(g+h)$ . But then  $g(n_0) = h(n_0) = x$  since  $x$  is an extreme point. Since  $\|x\| = 1$  and  $\phi$  is strictly increasing and  $\phi(1) = 1$ , we have  $g(n) = h(n) = 0$  for all  $n \neq n_0$ . But this implies that  $f = g = h$ , and  $f$  is extreme. This ends the proof of the theorem.  $\square$

As a corollary, we get

**Theorem 2.2.** *A point  $x$  is an extreme point of  $B_1(\ell^\phi)$  if and only if  $x_n = 0$  for all  $n$  except for one  $n$ , say,  $n_0$ , and  $|x_{n_0}| = 1$ .*

*Proof.* Take  $R$  for  $X$ .  $\square$

As for the exposed points we have

**Theorem 2.3.** *Let  $f \in B_1(\ell^\phi(X))$ . The following statements are equivalent:*

- (i)**  *$f$  is an exposed point.*
- (ii)**  *$f(n) = \delta_{nn_0} \cdot x$  and  $x$  is an exposed point of  $B_1(X)$ .*

*Proof.* **(i)**  $\longrightarrow$  **(ii)**. Let  $f$  be exposed. Then  $f$  is an extreme point. Hence  $f(n)\delta_{nn_0} \cdot x$  with  $x$  an extreme point of  $B_1(X)$ . If  $x$  is not exposed, then for every  $x^* \in B_1(X^*)$  with  $x^*(x) = 1$ , there exists  $z \in B_1(X)$  such that  $x^*(z) = 1$  and  $z \neq x$ . Now let  $F \in [\ell^\phi(X)]^* = \ell^\infty(X^*)$  such that  $\|F\| = 1$ , and  $F(f) = 1$ . In that case, if  $F = (x_1^*, x_2^*, \dots)$ , then  $F(f) = x_{n_0}^*(x) = 1$ . Since  $x$  is not exposed, there exists  $z \neq x$  in  $B_1(X)$  such that  $x_{n_0}^*(z) = 1$ . But then  $F(g) = 1$ , where  $g(n) = \delta_{nn_0} \cdot z$  and  $f$  is not exposed. Hence  $x$  must be exposed in  $B_1(X)$ .

Conversely: **(ii)**  $\longrightarrow$  **(i)**. Let  $f = \delta_{nn_0} \cdot x$  with  $x$  exposed in  $B_1(X)$ . If  $x^*$  is the functional that exposes  $x$ , then one can easily see that  $F(n) = \delta_{nn_0} \cdot x^*$  is the functional that exposes  $f$ . This ends the proof.  $\square$

Theorem 2.3 readily implies

**Theorem 2.4.** *An element  $f$  is an exposed point of  $B_1(\ell^\phi)$  if and only if  $f$  is extreme.*

As for smooth points we have

**Theorem 2.5.**  *$B_1(\ell^\phi(X))$  has no smooth points for any Banach space  $X$ .*

*Proof.* Let  $f \in B_1(\ell^\phi(X))$ . If there exists  $F \in B_1(\ell^\infty(X^*))$  such that  $F(f) = 1$ , then by Remark 1  $f$  must have only one nonzero coordinate, say,  $f(n_0) = x_{n_0}$ . Since  $\phi(1) = 1$ , it follows that  $\|x_{n_0}\| = 1$ . Consider the functionals:

$$\begin{aligned} F_1(n) &= \delta_{nn_0} \cdot x^* \text{ with } x^*(x_{n_0}) = 1, \\ F_2(n) &= \delta_{nn_0} \cdot x^* + \delta_{n,n_0+1} \cdot z^* \text{ with } \|z^*\| = 1. \end{aligned}$$

Then,  $F_1$  and  $F_2$  are two different elements in  $B_2(\ell^\phi(X))$  such that  $F_1(f) = F_2(f) = 1$ . Thus  $f$  is not smooth. This ends the proof.  $\square$

It follows that  $B_1(\ell^\phi)$  has no smooth points.

**3. Geometry of  $B_1(L(\ell^p))$ ,  $0 < p < 1$ .** The characterization of the extreme points of  $B_1(L(\ell^p))$ ,  $1 < p < \infty$ , is still an open difficult problem [1], [3]. In this section we give a complete description of the extreme points and the exposed points of the unit ball of  $L(\ell^p)$  for  $0 < p < 1$ . We remark that Kalton, [2], studied isomorphisms of and some classes of operators on  $\ell^p$ ,  $0 < p < 1$ .

**Theorem 3.1.** *Let  $T \in B_1(L(\ell^p))$ ,  $0 < p < 1$ . The following statements are equivalent:*

- (i)  $T$  is an extreme point.
- (ii)  $T$  is a permutation on the basis elements.

*Proof.* (ii)  $\rightarrow$  (i). Let  $T$  be a permutation of the basis elements  $e_1, e_2, \dots$ . If  $T$  is not extreme, then there exists  $S \in B_1(L(\ell^p))$  such that  $S \neq 0$  and  $\|S \pm T\| \leq 1$ . Thus  $\|(S \pm T)x\| \leq 1$  for all  $x$  in  $B_1(\ell^p)$ . Thus, in particular,  $\|Se_n \pm Te_n\| \leq 1$  for all  $n$ . Since  $\|S\| \leq 1$ , it follows that  $Te_n$  is not extreme for those  $n$  for which  $Se_n \neq 0$ . Since  $S \neq 0$ , we get a contradiction, noting that  $\pm e_n$  are the extreme points of  $\ell^p$ . Thus  $T$  must be extreme.

Conversely: (i)  $\rightarrow$  (ii). Let  $T$  be an extreme element of  $B_1(L(\ell^p))$ , but, if it is possible, assume there exists  $k_0$  such that  $Te_{k_0}$  is not a basis element and hence not an extreme element of  $B_1(\ell^p)$ . Thus there exists  $z$  in  $B_1(\ell^p)$  such that  $\|Te_{k_0} \pm z\| \leq 1$ . Define the operator  $S$  on  $\ell^p$  as  $S = e_{k_0} \otimes z$ , so  $Sx = x_{k_0}z$ . Then

$$\begin{aligned} \|(S \pm T)x\|_p &= \|(S \pm T)(\sum x_i e_i)\|_p = \|\sum x_i (S \pm T)e_i\|_p \\ &\leq \sum |x_i|^p \|(S \pm T)e_i\|_p. \end{aligned}$$

But

$$(S \pm T)e_i = \begin{cases} Te_0, & i \neq k_0, \\ z \pm Te_{k_0}, & i = k_0. \end{cases}$$

Thus in either case we have  $\|(S \pm T)e_i\| \leq 1$  for all  $i$ . So  $\|(S \pm T)x\| \leq \sum |x_i|^p$ . It follows that  $\|S \pm T\| \leq 1$ , and  $T$  is not extreme, which contradicts the assumption. So  $T$  must be a permutation. This ends the proof.  $\square$

To characterize the exposed points, we need

**Theorem 3.2.**  $L(\ell^p)$  is isometrically isomorphic to  $\ell^\infty(\ell^p)$ .

*Proof.* Let  $f \in \ell^\infty(\ell^p)$ . Then  $f : N \rightarrow \ell^p$  with  $\sup_n \|f(n)\|_p < \infty$ . Define  $T : \ell^p \rightarrow \ell^p$ , by  $Tx = \sum x_k f(k)$ . Then  $\|Tx\|_p \leq \sum \|x_p f(k)\|_p \leq \sum |x_k|^p \|f(k)\|_p \leq \|f\|_\infty \|x\|_p$ . Thus  $\|T\| \leq \|f\|_\infty$ . But  $Te_k = f(k)$ . So  $\|f(k)\|_p = \|Te_k\|_p \leq \|T\|$ . It follows that  $\|f\|_\infty \leq \|T\|$ . Hence  $\|f\|_\infty = \|T\|$ .

On the other hand, let  $T \in L(\ell^p)$ . Define  $f(n) = Te_p$ . Then one can easily show that  $f \in \ell^\infty(\ell^p)$  and  $\|f\|_\infty = \|T\|$ . This ends the proof.  $\square$

Now for the exposed points we have

**Theorem 3.3.** Let  $T \in B_1(L(\ell^p))$ . The following statements are equivalent:

- (i)  $T$  is exposed.
- (ii)  $T$  is extreme.

*Proof.* That (i)  $\rightarrow$  (ii) is immediate.

For the converse, let  $T$  be an extreme point. By Theorem 3.1,  $T$  is a permutation of the basis elements. Let  $f$  be the function corresponding to  $T$  as in Theorem 3.2. Thus  $f(n) = \pm e_{k(n)}$ . Define  $G : L(\ell^p) \rightarrow R$ ,  $G(S) = \sum t_n \langle f(n), g(n) \rangle$ , where  $0 < t_n$ ,  $\sum t_n = 1$ , and  $g$  is the element in  $\ell^\infty(\ell^p)$  that represents  $S$  as in Theorem 3.2. Then,  $G$  is bounded and  $\|G\| \leq 1$ . Further  $G(T) = 1$ . Now, if it is possible, assume there exists some  $S$  in  $B_1(L(\ell^p))$  such that  $G(S) = 1$ . Then  $\sum t_n \langle f(n), g(n) \rangle = 1$ . This implies that  $\langle f(n), g(n) \rangle = 1$ . Since  $f(n) = e_{k(n)}$ , it follows that  $g(n) = f(n)$ , and so  $S = T$ . Hence  $T$  is exposed. This ends the proof.  $\square$

#### ACKNOWLEDGMENT

The authors would like to thank the referee for a very careful reading of the manuscript. His suggestions and comments helped us to write the final version of the paper.

#### REFERENCES

1. W. DEEB and R. KHALIL, Exposed and smooth points of some classes of operators in  $L(\ell^p)$ ,  $1 \leq p < \infty$ . *J. Funct. Anal.* **103**(1992), 217–228.
2. N. J. KALTON, Isomorphisms between  $L_p$ -functions. *J. Funct. Anal.* **42**(1981), 299–337.
3. R. KHALIL, Smooth points of unit balls of operator and function spaces. *Demonstr. Math.* **29**(1996), 723–732.
4. W. RUCKLE, FK spaces in which the sequence of coordinate vector is bounded. *Canadian J. Math.* **25**(1973), 973–978.
5. W. DEEB and R. YOUNIS, Extreme points of a class of non-locally convex topological vector spaces. *Math. Rep. Toyama Univ.* **6**(1983), 95–103.

6. W. DEEB and R. KHALIL, Exposed and smooth points of some classes of operators in  $L(\ell^p)$ . *J. Funct. Anal.* **103**(1992), 217–228.
7. R. GRZASLEWICZ, Extreme operators on 2-dimensional  $\ell^p$ -spaces. *Coll. Math.* **44**(1981), 209–215.
8. R. GRZASLEWICZ, A note on extreme contractions on  $\ell^p$ -spaces. *Portugal. Math.* **40**(1981), 413–419.
9. R. KHALIL, A class of extreme contractions in  $L(\ell^p)$ . *Annali di Mat. Pura ed Applic.* **145**(1988), 1–5.
10. C.-H. KAN, A class of extreme  $L^p$  contractions,  $p \neq 1, 2, \infty$ . *Illinois J. Math.* **30**(1986), 612–635.

(Received 9.02.2001; revised 4.10.2001)

Authors' address:

Department of Mathematics  
University of Jordan, Amman  
Jordan

E-mail: roshdi@ju.edu.jo  
husein@ju.edu.jo  
wamin@ju.edu.jo