

## SOLVABILITY AND THE UNIQUE SOLVABILITY OF A PERIODIC TYPE BOUNDARY VALUE PROBLEM FOR FIRST ORDER SCALAR FUNCTIONAL DIFFERENTIAL EQUATIONS

R. HAKL, A. LOMTATIDZE, AND J. ŠREMR

**Abstract.** Nonimprovable in a certain sense, sufficient conditions for the solvability and unique solvability of the problem

$$u'(t) = F(u)(t), \quad u(a) - \lambda u(b) = h(u)$$

are established, where  $F : C([a, b]; R) \rightarrow L([a, b]; R)$  is a continuous operator satisfying the Carathéodory condition,  $h : C([a, b]; R) \rightarrow R$  is a continuous functional, and  $\lambda \in R_+$ .

**2000 Mathematics Subject Classification:** 34K10.

**Key words and phrases:** Functional differential equations, periodic type boundary value problem, existence and uniqueness.

### INTRODUCTION

The following notation is used throughout.

$R$  is the set of all real numbers,  $R_+ = [0, +\infty[$ .

$C([a, b]; R)$  is the Banach space of continuous functions  $u : [a, b] \rightarrow R$  with the norm  $\|u\|_C = \max\{|u(t)| : a \leq t \leq b\}$ .

$C([a, b]; R_+) = \{u \in C([a, b]; R) : u(t) \geq 0 \text{ for } t \in [a, b]\}$ .

$B_{\lambda c}^i([a, b]; R)$  is the set of functions  $u \in C([a, b]; R)$  satisfying the condition  $(-1)^{i+1}(u(a) - \lambda u(b)) \operatorname{sgn}((2-i)u(a) + (i-1)u(b)) \leq c$ , where  $c \in R$ ,  $i = 1, 2$ .

$\tilde{C}([a, b]; D)$ , where  $D \subseteq R$ , is the set of absolutely continuous functions  $u : [a, b] \rightarrow D$ .

$L([a, b]; R)$  is the Banach space of Lebesgue integrable functions  $p : [a, b] \rightarrow R$  with the norm  $\|p\|_L = \int_a^b |p(s)| ds$ .

$L([a, b]; R_+) = \{p \in L([a, b]; R) : p(t) \geq 0 \text{ for almost all } t \in [a, b]\}$ .

$\mathcal{M}_{ab}$  is the set of measurable functions  $\tau : [a, b] \rightarrow [a, b]$ .

$\mathcal{L}_{ab}$  is the set of linear operators  $\ell : C([a, b]; R) \rightarrow L([a, b]; R)$  for which there is a function  $\eta \in L([a, b]; R_+)$  such that

$$|\ell(v)(t)| \leq \eta(t) \|v\|_C \quad \text{for } t \in [a, b], \quad v \in C([a, b]; R).$$

$\mathcal{P}_{ab}$  is the set of linear operators  $\ell \in \mathcal{L}_{ab}$  mapping the set  $C([a, b]; R_+)$  into the set  $L([a, b]; R_+)$ .

$K_{ab}$  is the set of continuous operators  $F : C([a, b]; R) \rightarrow L([a, b]; R)$  satisfying the Carathéodory condition, i.e., for each  $r > 0$  there exists  $q_r \in L([a, b]; R_+)$  such that

$$|F(v)(t)| \leq q_r(t) \quad \text{for } t \in [a, b], \quad \|v\|_C \leq r.$$

$K([a, b] \times A; B)$ , where  $A \subseteq R^2$ ,  $B \subseteq R$ , is the set of functions  $f : [a, b] \times A \rightarrow B$  satisfying the Carathéodory conditions, i.e.,  $f(\cdot, x) : [a, b] \rightarrow B$  is a measurable function for all  $x \in A$ ,  $f(t, \cdot) : A \rightarrow B$  is a continuous function for almost all  $t \in [a, b]$ , and for each  $r > 0$  there exists  $q_r \in L([a, b]; R_+)$  such that

$$|f(t, x)| \leq q_r(t) \quad \text{for } t \in [a, b], \quad x \in A, \quad \|x\| \leq r.$$

$$[x]_+ = \frac{1}{2}(|x| + x), \quad [x]_- = \frac{1}{2}(|x| - x).$$

By a solution of the equation

$$u'(t) = F(u)(t), \tag{0.1}$$

where  $F \in K_{ab}$ , we understand a function  $u \in \tilde{C}([a, b]; R)$  satisfying the equation (0.1) almost everywhere in  $[a, b]$ .

Consider the problem on the existence and uniqueness of a solution of (0.1) satisfying the boundary condition

$$u(a) - \lambda u(b) = h(u), \tag{0.2}$$

where  $h : C([a, b]; R) \rightarrow R$  is a continuous functional such that for each  $r > 0$  there exists  $M_r \in R_+$  such that

$$|h(v)| \leq M_r \quad \text{for } \|v\|_C \leq r,$$

and  $\lambda \in R_+$ .

In the case where  $F$  is the so-called Nemitsky operator, the problem (0.1), (0.2) and analogous problems for systems of linear and nonlinear ordinary differential equations have been studied in detail (see [7, 15–18] and the references therein). The foundation of the theory of general boundary value problems for functional differential equations were laid in the monographs [1] and [30] (see also [2, 3, 8, 14, 19–27, 29]). In spite of a large number of papers devoted to boundary value problems for functional differential equations, at present only a few efficient sufficient solvability conditions are known even for the linear problem

$$u'(t) = \ell(u)(t) + q_0(t), \tag{0.3}$$

$$u(a) - \lambda u(b) = c_0, \tag{0.4}$$

where  $\ell \in \mathcal{L}_{ab}$ ,  $q_0 \in L([a, b]; R)$ ,  $\lambda \in R_+$ , and  $c_0 \in R$  (see [5, 6, 8–13, 19–26, 29]). Here we attempt to fill this gap in a certain way. More precisely, in Sections 1 and 2, nonimprovable efficient sufficient conditions are established for the solvability and unique solvability of the problems (0.3), (0.4) and (0.1), (0.2), respectively. Section 3 is devoted to the examples showing the optimality of our main results.

Finally, all results are concretized for the differential equations with deviating arguments of the forms

$$u'(t) = p(t)u(\tau(t)) - g(t)u(\mu(t)) + f(t, u(t), u(\nu(t))) \quad (0.5)$$

and

$$u'(t) = p(t)u(\tau(t)) - g(t)u(\mu(t)) + q_0(t), \quad (0.6)$$

where  $p, g \in L([a, b]; R_+)$ ,  $q_0 \in L([a, b]; R)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and  $f \in K([a, b] \times R^2; R)$ .

## 1. LINEAR PROBLEM

We need the following well-known result from the general theory of linear boundary value problems for functional differential equations (see, e.g., [4, 21, 30]).

**Theorem 1.1.** *The problem (0.3), (0.4) is uniquely solvable iff the corresponding homogeneous problem*

$$u'(t) = \ell(u)(t), \quad (0.3_0)$$

$$u(a) - \lambda u(b) = 0 \quad (0.4_0)$$

has only a trivial solution.

*Remark 1.1.* It follows from the Riesz–Schauder theory that if  $\ell \in \mathcal{L}_{ab}$  and the problem (0, 3<sub>0</sub>), (0, 4<sub>0</sub>) has a nontrivial solution, then there exist  $q_0 \in L([a, b]; R)$  and  $c_0 \in R$  such that the problem (0.3), (0.4) has no solution.

### 1.1. Main Results.

**Theorem 1.2.** *Assume that  $\lambda \in [0, 1[$ , the operator  $\ell$  admits the representation  $\ell = \ell_0 - \ell_1$ , where*

$$\ell_0, \ell_1 \in \mathcal{P}_{ab}, \quad (1.1)$$

and let there exist a function  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfying the inequalities

$$\gamma'(t) \geq \ell_0(\gamma)(t) + \ell_1(1)(t) \quad \text{for } t \in [a, b], \quad (1.2)$$

$$\gamma(a) > \lambda \gamma(b), \quad (1.3)$$

$$\gamma(b) - \gamma(a) < 3 + \lambda. \quad (1.4)$$

Then the problem (0.3), (0.4) has a unique solution.

*Remark 1.2.* Theorem 1.2 is nonimprovable in a certain sense. More precisely, the strict inequality (1.4) cannot be replaced by nonstrict one (see On Remark 1.2 and Example 3.1).

*Remark 1.3.* Let  $\lambda \in [1, +\infty[$  and  $\ell = \ell_0 - \ell_1$ , where  $\ell_0, \ell_1 \in \mathcal{P}_{ab}$ . Introduce the operator  $\psi : L([a, b]; R) \rightarrow L([a, b]; R)$  by

$$\psi(w)(t) \stackrel{\text{def}}{=} w(a + b - t) \quad \text{for } t \in [a, b].$$

Let  $\varphi$  be a restriction of  $\psi$  to the space  $C([a, b]; R)$ . Put  $\vartheta = \frac{1}{\lambda}$ , and

$$\widehat{\ell}_0(w)(t) \stackrel{\text{def}}{=} \psi(\ell_0(\varphi(w)))(t), \quad \widehat{\ell}_1(w)(t) \stackrel{\text{def}}{=} \psi(\ell_1(\varphi(w)))(t) \quad \text{for } t \in [a, b].$$

It is clear that if  $u$  is a solution of the problem  $(0, 3_0)$ ,  $(0, 4_0)$ , then the function  $v \stackrel{\text{def}}{=} \varphi(u)$  is a solution of the problem

$$v'(t) = \widehat{\ell}_1(v)(t) - \widehat{\ell}_0(v)(t), \quad v(a) - \vartheta v(b) = 0, \quad (1.5)$$

and, conversely, if  $v$  is a solution of the problem (1.5), then the function  $u \stackrel{\text{def}}{=} \varphi(v)$  is a solution of the problem  $(0, 3_0)$ ,  $(0, 4_0)$ .

It is also obvious that if a function  $\gamma \in \widetilde{C}([a, b]; ]0, +\infty[)$  satisfies the inequality (1.2), then  $\beta \stackrel{\text{def}}{=} \varphi(\gamma)$  satisfies the inequality

$$\beta'(t) \leq -\widehat{\ell}_0(\beta)(t) - \widehat{\ell}_1(1)(t) \quad \text{for } t \in [a, b], \quad (1.6)$$

and, conversely, if a function  $\beta \in \widetilde{C}([a, b]; ]0, +\infty[)$  satisfies the inequality (1.6), then  $\gamma \stackrel{\text{def}}{=} \varphi(\beta)$  satisfies the inequality (1.2).

In view of Remark 1.3 the following statement is an immediate consequence of Theorem 1.2.

**Theorem 1.3.** *Let  $\lambda \in ]1, +\infty[$ , the operator  $\ell$  admit the representation  $\ell = \ell_0 - \ell_1$ , where  $\ell_0, \ell_1 \in \mathcal{P}_{ab}$ , and let there exist a function  $\gamma \in \widetilde{C}([a, b]; ]0, +\infty[)$  satisfying the inequalities*

$$\gamma'(t) \leq -\ell_1(\gamma)(t) - \ell_0(1)(t) \quad \text{for } t \in [a, b], \quad (1.7)$$

$$\gamma(a) < \lambda\gamma(b), \quad (1.8)$$

$$\gamma(a) - \gamma(b) < 3 + \frac{1}{\lambda}. \quad (1.9)$$

*Then the problem (0.3), (0.4) has a unique solution.*

*Remark 1.4.* According to Remarks 1.2 and 1.3, Theorem 1.3 is nonimprovable in that sense that the strict inequality (1.9) cannot be replaced by nonstrict one.

Theorem 1.2 implies the following assertions for the problem (0.6), (0.4).

**Corollary 1.1.** *Let  $\lambda \in [0, 1[$ ,  $p, g \in L([a, b]; R_+)$ , and  $\tau \in \mathcal{M}_{ab}$  be such that*

$$\lambda \exp \left( \int_a^b p(s) ds \right) < 1, \quad (1.10)$$

$$(t - \tau(t))p(t) \geq 0 \quad \text{for } t \in [a, b], \quad (1.11)$$

and

$$\frac{1 - \lambda}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \int_a^b g(s) \exp\left(\int_s^b p(\xi) d\xi\right) ds < 3 + \lambda. \quad (1.12)$$

Then the problem (0.6), (0.4) has a unique solution.

**Corollary 1.2.** Let  $\lambda \in [0, 1[$ ,  $p, g \in L([a, b]; R_+)$ , and  $\tau \in \mathcal{M}_{ab}$  be such that

$$(1 - \lambda) \left( \int_a^b g(s) ds + \alpha_1 \right) + (3 + \lambda) \beta_1 < 3 + \lambda, \quad (1.13)$$

where

$$\alpha_1 = \int_a^b p(s) \left( \int_a^{\tau(s)} g(\xi) d\xi \right) \exp\left(\int_s^b p(\xi) d\xi\right) ds, \quad (1.14)$$

$$\beta_1 = \lambda \exp\left(\int_a^b p(s) ds\right) + \int_a^b p(s) \sigma(s) \left( \int_s^{\tau(s)} p(\xi) d\xi \right) \exp\left(\int_s^b p(\xi) d\xi\right) ds, \quad (1.15)$$

$$\sigma(t) = \frac{1}{2} (1 + \operatorname{sgn}(\tau(t) - t)) \quad \text{for } t \in [a, b]. \quad (1.16)$$

Then the problem (0.6), (0.4) has a unique solution.

*Remark 1.5.* Corollaries 1.1 and 1.2 are nonimprovable in a certain sense. More precisely, the strict inequalities (1.12) in Corollary 1.1 and (1.13) in Corollary 1.2 cannot be replaced by nonstrict ones (see On Remark 1.2 and Example 3.1).

Theorem 1.3 implies the following statements.

**Corollary 1.3.** Let  $\lambda \in ]1, +\infty[$ ,  $p, g \in L([a, b]; R_+)$ , and  $\mu \in \mathcal{M}_{ab}$  be such that

$$\exp\left(\int_a^b g(s) ds\right) < \lambda, \quad (1.17)$$

$$(\mu(t) - t)g(t) \geq 0 \quad \text{for } t \in [a, b], \quad (1.18)$$

and

$$\frac{\lambda - 1}{\lambda - \exp\left(\int_a^b g(\xi) d\xi\right)} \int_a^b p(s) \exp\left(\int_a^s g(\xi) d\xi\right) ds < 3 + \frac{1}{\lambda}. \quad (1.19)$$

Then the problem (0.6), (0.4) has a unique solution.

**Corollary 1.4.** *Let  $\lambda \in ]1, +\infty[$ ,  $p, g \in L([a, b]; R_+)$ , and  $\mu \in \mathcal{M}_{ab}$  be such that*

$$\frac{\lambda - 1}{\lambda} \left( \int_a^b p(s) ds + \alpha_2 \right) + \left( 3 + \frac{1}{\lambda} \right) \beta_2 < 3 + \frac{1}{\lambda}, \quad (1.20)$$

where

$$\alpha_2 = \int_a^b g(s) \left( \int_{\mu(s)}^b p(\xi) d\xi \right) \exp \left( \int_a^s g(\xi) d\xi \right) ds, \quad (1.21)$$

$$\beta_2 = \frac{1}{\lambda} \exp \left( \int_a^b g(s) ds \right) + \int_a^b g(s) \sigma(s) \left( \int_{\mu(s)}^s g(\xi) d\xi \right) \exp \left( \int_a^s g(\xi) d\xi \right) ds, \quad (1.22)$$

$$\sigma(t) = \frac{1}{2} (1 + \operatorname{sgn}(t - \mu(t))) \quad \text{for } t \in [a, b]. \quad (1.23)$$

Then the problem (0.6), (0.4) has a unique solution.

*Remark 1.6.* Corollaries 1.3 and 1.4 are nonimprovable in a certain sense. More precisely, the strict inequalities (1.19) in Corollary 1.3 and (1.20) in Corollary 1.4 cannot be replaced by nonstrict ones.

**1.2. Proofs of Main Results.** To prove Theorem 1.2, we need a result from [14]. Let us first introduce the following definition.

**Definition 1.1.** We will say that an operator  $\ell \in \mathcal{L}_{ab}$  belongs to the set  $V^+(\lambda)$  (resp.  $V^-(\lambda)$ ) if the homogeneous problem  $(0, 3_0)$ ,  $(0, 4_0)$  has only a trivial solution and, for arbitrary  $q_0 \in L([a, b]; R_+)$  and  $c_0 \in R_+$ , the solution of the problem (0.3), (0.4) is nonnegative (resp. nonpositive).

*Remark 1.7.* It follows immediately from Definition 1.1 that  $\ell \in V^+(\lambda)$  (resp.  $\ell \in V^-(\lambda)$ ) iff a certain theorem on differential inequalities holds for the equation (0.3), i.e., if  $u, v \in \tilde{C}([a, b]; R)$  satisfy the inequalities

$$\begin{aligned} u'(t) &\leq \ell(u)(t) + q_0(t), & v'(t) &\geq \ell(v)(t) + q_0(t) \quad \text{for } t \in [a, b], \\ u(a) - \lambda u(b) &\leq v(a) - \lambda v(b), \end{aligned}$$

then  $u(t) \leq v(t)$  (resp.  $u(t) \geq v(t)$ ) for  $t \in [a, b]$ .

**Lemma 1.1** ([14]). *Let  $\lambda \in [0, 1[$  and  $\ell \in \mathcal{P}_{ab}$ . Then  $\ell \in V^+(\lambda)$  iff there exist  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfying*

$$\begin{aligned} \gamma'(t) &\geq \ell(\gamma)(t) \quad \text{for } t \in [a, b], \\ \gamma(a) &> \lambda \gamma(b). \end{aligned}$$

*Proof of Theorem 1.2.* According to Theorem 1.1, it is sufficient to show that the homogeneous problem  $(0, 3_0)$ ,  $(0, 4_0)$  has only a trivial solution.

Assume that, on the contrary, the problem  $(0, 3_0)$ ,  $(0, 4_0)$  has a nontrivial solution  $u$ . By virtue of (1.1) and Lemma 1.1, it follows from (1.2) and (1.3) that  $\ell_0 \in V^+(\lambda)$ . Consequently, according to Definition 1.1,  $u$  must change its sign. Put

$$M = \max\{u(t) : t \in [a, b]\}, \quad m = -\min\{u(t) : t \in [a, b]\}, \quad (1.24)$$

and choose  $t_M, t_m \in [a, b]$  so that

$$u(t_M) = M, \quad u(t_m) = -m. \quad (1.25)$$

Obviously,

$$M > 0, \quad m > 0 \quad (1.26)$$

and without loss of generality we can assume that  $t_M < t_m$ . In view of (1.26), the relations  $(0, 3_0)$ ,  $(0, 4_0)$ , (1.2) and (1.3) yield

$$\begin{aligned} (M\gamma(t) + u(t))' &\geq \ell_0(M\gamma + u)(t) + \ell_1(M - u)(t) \quad \text{for } t \in [a, b], \\ M\gamma(a) + u(a) - \lambda(M\gamma(b) + u(b)) &\geq 0, \end{aligned} \quad (1.27)$$

and

$$\begin{aligned} (m\gamma(t) - u(t))' &\geq \ell_0(m\gamma - u)(t) + \ell_1(m + u)(t) \quad \text{for } t \in [a, b], \\ m\gamma(a) - u(a) - \lambda(m\gamma(b) - u(b)) &\geq 0. \end{aligned} \quad (1.28)$$

Hence, according to (1.1), (1.24), the condition  $\ell_0 \in V^+(\lambda)$ , and Remark 1.7, it follows that

$$M\gamma(t) + u(t) \geq 0, \quad m\gamma(t) - u(t) \geq 0 \quad \text{for } t \in [a, b].$$

By virtue of the last two inequalities, (1.1), and (1.24), from (1.27) and (1.28) we get

$$(M\gamma(t) + u(t))' \geq 0, \quad (m\gamma(t) - u(t))' \geq 0 \quad \text{for } t \in [a, b]. \quad (1.29)$$

In view of (1.25) and (1.26), the integration of the first inequality in (1.29) from  $t_M$  to  $t_m$  results in

$$M\gamma(t_m) - m - M\gamma(t_M) - M \geq 0,$$

i.e.,

$$\gamma(t_m) - \gamma(t_M) \geq 1 + \frac{m}{M}. \quad (1.30)$$

On the other hand, on account of (1.25) and (1.26), the integration of the second inequality in (1.29) from  $a$  to  $t_M$  and from  $t_m$  to  $b$  yields

$$\begin{aligned} m\gamma(t_M) - M - m\gamma(a) + u(a) &\geq 0, \\ m\gamma(b) - u(b) - m\gamma(t_m) - m &\geq 0. \end{aligned}$$

Summing these two inequalities and taking into account that

$$u(b) - u(a) = u(b)(1 - \lambda) \geq -m(1 - \lambda),$$

we get

$$\gamma(t_M) - \gamma(t_m) + \gamma(b) - \gamma(a) \geq \lambda + \frac{M}{m}. \quad (1.31)$$

Now from (1.30) and (1.31) we have

$$\gamma(b) - \gamma(a) \geq 1 + \lambda + \frac{m}{M} + \frac{M}{m} \geq 3 + \lambda,$$

which contradicts (1.4).  $\square$

*Proof of Corollary 1.1.* According to (1.12), there exists  $\varepsilon > 0$  such that

$$\begin{aligned} & \frac{\varepsilon}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \left( \exp\left(\int_a^b p(s) ds\right) - 1 \right) \\ & + \frac{1 - \lambda}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \int_a^b g(s) \exp\left(\int_s^b p(\xi) d\xi\right) ds < 3 + \lambda. \end{aligned}$$

Put

$$\begin{aligned} \gamma(t) = & \frac{\varepsilon}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \exp\left(\int_a^t p(s) ds\right) \\ & + \frac{1}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \int_a^t g(s) \exp\left(\int_s^t p(\xi) d\xi\right) ds \\ & + \frac{\lambda \exp\left(\int_a^b p(s) ds\right)}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \int_t^b g(s) \exp\left(\int_s^t p(\xi) d\xi\right) ds \quad \text{for } t \in [a, b]. \end{aligned}$$

Then  $\gamma$  is a solution of the problem

$$\gamma'(t) = p(t)\gamma(t) + g(t), \quad \gamma(a) - \lambda\gamma(b) = \varepsilon. \quad (1.32)$$

Since  $\varepsilon > 0$ , in view of (1.10), we have  $\gamma(t) > 0$  for  $t \in [a, b]$ . Consequently, (1.32) implies  $\gamma'(t) \geq 0$  for  $t \in [a, b]$ , and (1.11) yields

$$p(t)\gamma(t) \geq p(t)\gamma(\tau(t)) \quad \text{for } t \in [a, b]. \quad (1.33)$$



Therefore, by virtue of (1.32) and (1.33), the function  $\gamma$  satisfies the inequalities (1.2), (1.3), and (1.4) with

$$\begin{aligned} \ell_0(w)(t) \stackrel{\text{def}}{=} p(t)w(\tau(t)), \quad \ell_1(w)(t) \stackrel{\text{def}}{=} g(t)w(\mu(t)) \\ \text{for } t \in [a, b]. \quad \square \end{aligned} \quad (1.34)$$

*Proof of Corollary 1.2.* Let the operators  $\ell_0$  and  $\ell_1$  be defined by (1.34). From (1.13) it follows that  $\beta_1 < 1$ . Consequently, by [14, Theorem 2.1 c)], we have  $\ell_0 \in V^+(\lambda)$ . Choose  $\delta > 0$  and  $\varepsilon > 0$  such that

$$(1 - \lambda)(1 - \beta_1)^{-1} \left( \alpha_1 + \int_a^b g(s) ds \right) < 3 + \lambda - \delta, \quad (1.35)$$

$$\varepsilon < \frac{\delta(1 - \beta_1)}{1 - \lambda} \exp \left( - \int_a^b p(s) ds \right). \quad (1.36)$$

According to the condition  $\ell_0 \in V^+(\lambda)$  and Theorem 1.1, the problem

$$\gamma'(t) = p(t)\gamma(\tau(t)) + g(t), \quad (1.37)$$

$$\gamma(a) - \lambda\gamma(b) = \varepsilon \quad (1.38)$$

has a unique solution  $\gamma$ . It is clear that the conditions (1.2) and (1.3) are fulfilled. Due to the conditions  $\ell_0 \in V^+(\lambda)$ ,  $g \in L([a, b]; R_+)$ , and  $\varepsilon > 0$ , we get  $\gamma(t) \geq 0$  for  $t \in [a, b]$ . Hence, by (1.37), we find that  $\gamma(t) > 0$  for  $t \in [a, b]$ . On the other hand,  $\gamma$  is a solution of the equation

$$\gamma'(t) = p(t)\gamma(t) + p(t) \int_t^{\tau(t)} p(s)\gamma(\tau(s)) ds + p(t) \int_t^{\tau(t)} g(s) ds + g(t).$$

Hence, the Cauchy formula implies

$$\gamma(b) \leq \beta_1\gamma(b) + \alpha_1 + \int_a^b g(s) ds + \varepsilon \exp \left( \int_a^b p(s) ds \right).$$

The last inequality results in

$$\gamma(b) \leq (1 - \beta_1)^{-1} \left( \alpha_1 + \int_a^b g(s) ds \right) + \varepsilon(1 - \beta_1)^{-1} \exp \left( \int_a^b p(s) ds \right)$$

and thus, in view of (1.35), (1.36), and (1.38), we have

$$\gamma(b) - \gamma(a) \leq (1 - \lambda)\gamma(b) < 3 + \lambda.$$

Therefore the assumptions of Theorem 1.2 are fulfilled.  $\square$

Corollaries 1.3 and 1.4 can be proved analogously.

## 2. NONLINEAR PROBLEM

In what follows we will assume that  $q \in K([a, b] \times R; R_+)$  is nondecreasing in the second argument and satisfies

$$\lim_{x \rightarrow +\infty} \frac{1}{x} \int_a^b q(s, x) ds = 0. \quad (2.1)$$

## 2.1. Main Results.

**Theorem 2.1.** *Let  $\lambda \in [0, 1[$ ,  $c \in R_+$ ,*

$$h(v) \operatorname{sgn} v(a) \leq c \quad \text{for } v \in C([a, b]; R), \quad (2.2)$$

and let there exist  $\ell_0, \ell_1 \in \mathcal{P}_{ab}$  such that the inequality

$$[F(v)(t) - \ell_0(v)(t) + \ell_1(v)(t)] \operatorname{sgn} v(t) \leq q(t, \|v\|_C) \quad \text{for } t \in [a, b] \quad (2.3)$$

holds on the set  $B_{\lambda c}^1([a, b]; R)$ . If, moreover, there exists  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfying inequalities (1.2), (1.3), and

$$\gamma(b) - \gamma(a) < 2, \quad (2.4)$$

then the problem (0.1), (0.2) has at least one solution.

*Remark 2.1.* Theorem 2.1 is nonimprovable in a certain sense. More precisely, the inequality (2.4) cannot be replaced by the inequality

$$\gamma(b) - \gamma(a) \leq 2 + \varepsilon \quad (2.5)$$

no matter how small  $\varepsilon > 0$  is (see On Remark 2.1 and Example 3.2).

**Theorem 2.2.** *Let  $\lambda \in [0, 1[$ ,*

$$[h(v) - h(w)] \operatorname{sgn}(v(a) - w(a)) \leq 0 \quad \text{for } v, w \in C([a, b]; R), \quad (2.6)$$

and let there exist  $\ell_0, \ell_1 \in \mathcal{P}_{ab}$  such that the inequality

$$\begin{aligned} & [F(v)(t) - F(w)(t) - \ell_0(v - w)(t) \\ & + \ell_1(v - w)(t)] \operatorname{sgn}(v(t) - w(t)) \leq 0 \quad \text{for } t \in [a, b] \end{aligned} \quad (2.7)$$

is fulfilled on the set  $B_{\lambda c}^1([a, b]; R)$ , where  $c = |h(0)|$ . If, moreover, there exists a function  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfying the inequalities (1.2), (1.3), and (2.4), then the problem (0.1), (0.2) is uniquely solvable.

*Remark 2.2.* Theorem 2.2 is nonimprovable in the sense that the inequality (2.4) cannot be replaced by the inequality (2.5) no matter how small  $\varepsilon > 0$  is (see On Remark 2.2).

*Remark 2.3.* Let  $\lambda \in [1, +\infty[$ , and  $\varphi, \psi$  be the operators defined in Remark 1.3. Put  $\vartheta = \frac{1}{\lambda}$ , and

$$\widehat{F}(w)(t) \stackrel{\text{def}}{=} -\psi(F(\varphi(w)))(t) \quad \text{for } t \in [a, b], \quad \widehat{h}(w) \stackrel{\text{def}}{=} -\vartheta h(\varphi(w)).$$

It is clear that if  $u$  is a solution of the problem (0.1), (0.2), then the function  $v \stackrel{\text{def}}{=} \varphi(u)$  is a solution of the problem

$$v'(t) = \widehat{F}(v)(t), \quad v(a) - \vartheta v(b) = \widehat{h}(v), \quad (2.8)$$

and vice versa, if  $v$  is a solution of the problem (2.8), then the function  $u \stackrel{\text{def}}{=} \varphi(v)$  is a solution of the problem (0.1), (0.2).

Therefore, according to Remarks 1.3 and 2.3, Theorems 2.1 and 2.2 imply

**Theorem 2.3.** *Let  $\lambda \in ]1, +\infty[$ ,  $c \in R_+$ ,*

$$h(v) \operatorname{sgn} v(b) \geq -c \quad \text{for } v \in C([a, b]; R), \quad (2.9)$$

and let there exist  $\ell_0, \ell_1 \in \mathcal{P}_{ab}$  such that the inequality

$$[F(v)(t) - \ell_0(v)(t) + \ell_1(v)(t)] \operatorname{sgn} v(t) \geq -q(t, \|v\|_C) \quad \text{for } t \in [a, b]$$

is fulfilled on the set  $B_{\lambda c}^2([a, b]; R)$ . If, moreover, there exists  $\gamma \in \widetilde{C}([a, b]; ]0, +\infty[)$  satisfying inequalities (1.7), (1.8), and

$$\gamma(a) - \gamma(b) < 2, \quad (2.10)$$

then the problem (0.1), (0.2) has at least one solution.

**Theorem 2.4.** *Let  $\lambda \in ]1, +\infty[$ ,*

$$[h(v) - h(w)] \operatorname{sgn}(v(b) - w(b)) \geq 0 \quad \text{for } v, w \in C([a, b]; R), \quad (2.11)$$

and let there exist  $\ell_0, \ell_1 \in \mathcal{P}_{ab}$  such that the inequality

$$\begin{aligned} & [F(v)(t) - F(w)(t) - \ell_0(v - w)(t) \\ & + \ell_1(v - w)(t)] \operatorname{sgn}(v(t) - w(t)) \geq 0 \quad \text{for } t \in [a, b] \end{aligned}$$

is fulfilled on the set  $B_{\lambda c}^2([a, b]; R)$ , where  $c = |h(0)|$ . If, moreover, there exists a function  $\gamma \in \widetilde{C}([a, b]; ]0, +\infty[)$  satisfying inequalities (1.7), (1.8), and (2.10), then the problem (0.1), (0.2) is uniquely solvable.

*Remark 2.4.* According to Remarks 1.3 and 2.1–2.3, Theorems 2.3 and 2.4 are nonimprovable in the sense that the strict inequality (2.10) cannot be replaced by the inequality

$$\gamma(a) - \gamma(b) \leq 2 + \varepsilon$$

no matter how small  $\varepsilon > 0$  is.

Theorems 2.1 and 2.2 imply the following results for the problem (0.5), (0.2).

**Corollary 2.1.** *Let  $\lambda \in [0, 1[$ ,  $c \in R_+$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and let (2.2) and the condition*

$$f(t, x, y) \operatorname{sgn} x \leq q(t, |x|) \quad \text{for } t \in [a, b], \quad x, y \in R \tag{2.12}$$

*be fulfilled. If, moreover, the inequalities (1.10), (1.11), and*

$$\frac{1 - \lambda}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \int_a^b g(s) \exp\left(\int_s^b p(\xi) d\xi\right) ds < 2 \tag{2.13}$$

*hold, then the problem (0.5), (0.2) has at least one solution.*

**Corollary 2.2.** *Let  $\lambda \in [0, 1[$ ,  $c \in R_+$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and let conditions (2.2) and (2.12) be fulfilled. If, moreover,*

$$(1 - \lambda) \left( \int_a^b g(s) ds + \alpha_1 \right) + 2\beta_1 < 2 \tag{2.14}$$

*holds, where  $\alpha_1$  and  $\beta_1$  are defined by (1.14) and (1.15) with  $\sigma$  given by (1.16), then the problem (0.5), (0.2) has at least one solution.*

*Remark 2.5.* Corollaries 2.1 and 2.2 are nonimprovable in a certain sense. More precisely, the strict inequalities (2.13) in Corollary 2.1 and (2.14) in Corollary 2.2 cannot be replaced by the inequalities

$$\frac{1 - \lambda}{1 - \lambda \exp\left(\int_a^b p(s) ds\right)} \int_a^b g(s) \exp\left(\int_s^b p(\xi) d\xi\right) ds \leq 2 + \varepsilon \tag{2.15}$$

and

$$(1 - \lambda) \left( \int_a^b g(s) ds + \alpha_1 \right) + 2\beta_1 \leq 2 + \varepsilon, \tag{2.16}$$

no matter how small  $\varepsilon > 0$  is (see On Remark 2.1 and Example 3.2).

**Corollary 2.3.** *Let  $\lambda \in [0, 1[$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and let (2.6) and the condition*

$$\begin{aligned} [f(t, x_1, y_1) - f(t, x_2, y_2)] \operatorname{sgn}(x_1 - x_2) &\leq 0 \\ \text{for } t \in [a, b], \quad x_1, x_2, y_1, y_2 \in R \end{aligned} \tag{2.17}$$

*be fulfilled. If, moreover, inequalities (1.10), (1.11), and (2.13) hold, then the problem (0.5), (0.2) is uniquely solvable.*

**Corollary 2.4.** *Let  $\lambda \in [0, 1[$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and let the conditions (2.6) and (2.17) be fulfilled. If, moreover, the inequality (2.14) holds, where  $\alpha_1$  and  $\beta_1$  are defined by (1.14) and (1.15) with  $\sigma$  given by (1.16), then the problem (0.5), (0.2) is uniquely solvable.*

*Remark 2.6.* Corollaries 2.3 and 2.4 are nonimprovable in that sense that the strict inequalities (2.13) in Corollary 2.3 and (2.14) in Corollary 2.4 cannot be replaced by inequalities (2.15) and (2.16), no matter how small  $\varepsilon > 0$  is (see On Remark 2.2).

The following corollaries of Theorems 2.3 and 2.4 hold true.

**Corollary 2.5.** *Let  $\lambda \in ]1, +\infty[$ ,  $c \in R_+$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and let (2.9) and the condition*

$$f(t, x, y) \operatorname{sgn} x \geq -q(t, |x|) \quad \text{for } t \in [a, b], x, y \in R \quad (2.18)$$

*be fulfilled. If, moreover, inequalities (1.17), (1.18), and*

$$\frac{\lambda - 1}{\lambda - \exp\left(\int_a^b g(\xi) d\xi\right)} \int_a^b p(s) \exp\left(\int_a^s g(\xi) d\xi\right) ds < 2 \quad (2.19)$$

*hold, then the problem (0.5), (0.2) has at least one solution.*

**Corollary 2.6.** *Let  $\lambda \in ]1, +\infty[$ ,  $c \in R_+$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and the conditions (2.9) and (2.18) be fulfilled. If, moreover,*

$$\frac{\lambda - 1}{\lambda} \left( \int_a^b p(s) ds + \alpha_2 \right) + 2\beta_2 < 2, \quad (2.20)$$

*where  $\alpha_2$  and  $\beta_2$  are defined by (1.21) and (1.22) with  $\sigma$  given by (1.23), then the problem (0.5), (0.2) has at least one solution.*

**Corollary 2.7.** *Let  $\lambda \in ]1, +\infty[$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu \in \mathcal{M}_{ab}$  and let (2.11) and the condition*

$$\begin{aligned} [f(t, x_1, y_1) - f(t, x_2, y_2)] \operatorname{sgn}(x_1 - x_2) &\geq 0 \\ \text{for } t \in [a, b], x_1, x_2, y_1, y_2 \in R, \end{aligned} \quad (2.21)$$

*be fulfilled. If, moreover, inequalities (1.17), (1.18), and (2.19) hold, then the problem (0.5), (0.2) is uniquely solvable.*

**Corollary 2.8.** *Let  $\lambda \in ]1, +\infty[$ ,  $p, g \in L([a, b]; R_+)$ ,  $\tau, \mu, \nu \in \mathcal{M}_{ab}$ , and let the conditions (2.11) and (2.21) be fulfilled. If, moreover, the inequality (2.20) holds, where  $\alpha_2$  and  $\beta_2$  are defined by (1.21) and (1.22) with  $\sigma$  given by (1.23), then the problem (0.5), (0.2) is uniquely solvable.*

*Remark 2.7.* Corollaries 2.5–2.8 are nonimprovable in the sense that the strict inequality (2.19) in Corollaries 2.5 and (2.7), and the strict inequality (2.20) in Corollaries 2.6 and (2.8) cannot be replaced by the inequalities

$$\frac{\lambda - 1}{\lambda - \exp\left(\int_a^b g(\xi) d\xi\right)} \int_a^b p(s) \exp\left(\int_a^s g(\xi) d\xi\right) ds \leq 2 + \varepsilon$$

and

$$\frac{\lambda - 1}{\lambda} \left( \int_a^b p(s) ds + \alpha_2 \right) + 2\beta_2 \leq 2 + \varepsilon,$$

no matter how small  $\varepsilon > 0$  is.

**2.2. Auxiliary Propositions.** First we formulate the result from [24, Theorem 1] in the form suitable for us.

**Lemma 2.1.** *Let there exist a positive number  $\rho$  and an operator  $\ell \in \mathcal{L}_{ab}$  such that the homogeneous problem  $(0, 3_0)$ ,  $(0, 4_0)$  has only a trivial solution, and let for every  $\delta \in ]0, 1[$ , an arbitrary function  $u \in \tilde{C}([a, b]; R)$  satisfying the relations*

$$u'(t) = \ell(u)(t) + \delta[F(u)(t) - \ell(u)(t)], \quad u(a) - \lambda u(b) = \delta h(u), \quad (2.22)$$

admits the estimate

$$\|u\|_C \leq \rho. \quad (2.23)$$

Then the problem (0.1), (0.2) has at least one solution.

**Definition 2.1.** We say that an operator  $\ell \in \mathcal{L}_{ab}$  belongs to the set  $U(\lambda)$  if there exists a positive number  $r$  such that, for arbitrary  $q^* \in L([a, b]; R_+)$  and  $c \in R_+$ , every function  $u \in \tilde{C}([a, b]; R)$  satisfying the inequalities

$$[u(a) - \lambda u(b)] \operatorname{sgn} u(a) \leq c, \quad (2.24)$$

$$[u'(t) - \ell(u)(t)] \operatorname{sgn} u(t) \leq q^*(t) \quad \text{for } t \in [a, b], \quad (2.25)$$

admits the estimate

$$\|u\|_C \leq r(c + \|q^*\|_L). \quad (2.26)$$

**Lemma 2.2.** *Let  $c \in R_+$ ,*

$$h(v) \operatorname{sgn} v(a) \leq c \quad \text{for } v \in C([a, b]; R), \quad (2.27)$$

and let there exist  $\ell \in U(\lambda)$  such that the inequality

$$[F(v)(t) - \ell(v)(t)] \operatorname{sgn} v(t) \leq q(t, \|v\|_C) \quad \text{for } t \in [a, b] \quad (2.28)$$

is fulfilled on the set  $B_{\lambda c}^1([a, b]; R)$ . Then the problem (0.1), (0.2) has at least one solution.

*Proof.* First of all we note that, due to the condition  $\ell \in U(\lambda)$ , the homogeneous problem  $(0, 3_0)$ ,  $(0, 4_0)$  has only a trivial solution.

Let  $r$  be the number appearing in Definition 2.1. According to (2.1), there exists  $\rho > 2rc$  such that

$$\frac{1}{x} \int_a^b q(s, x) ds < \frac{1}{2r} \quad \text{for } x > \rho.$$

Now assume that  $u \in \tilde{C}([a, b]; R)$  satisfies (2.22) for some  $\delta \in ]0, 1[$ . Then, according to (2.27),  $u$  satisfies the inequality (2.24), i.e.,  $u \in B_{\lambda c}^1([a, b]; R)$ . By (2.28) we obtain that the inequality (2.25) is fulfilled with  $q^*(t) = q(t, \|u\|_C)$  for  $t \in [a, b]$ . Hence, according to the condition  $\ell \in U(\lambda)$  and the definition of the number  $\rho$ , we arrive at the estimate (2.23).

Since  $\rho$  depends neither on  $u$  nor on  $\delta$ , it follows from Lemma 2.1 that the problem (0.1), (0.2) has at least one solution.  $\square$

**Lemma 2.3.** *Let*

$$[h(v) - h(w)] \operatorname{sgn}(v(a) - w(a)) \leq 0 \quad \text{for } v, w \in C([a, b]; R) \quad (2.29)$$

and let there exist  $\ell \in U(\lambda)$  such that the inequality

$$[F(v)(t) - F(w)(t) - \ell(v - w)(t)] \operatorname{sgn}(v(t) - w(t)) \leq 0 \quad \text{for } t \in [a, b] \quad (2.30)$$

holds on the set  $B_{\lambda c}^1([a, b]; R)$ , where  $c = |h(0)|$ . Then the problem (0.1), (0.2) is uniquely solvable.

*Proof.* It follows from (2.29) that the condition (2.27) is fulfilled with  $c = |h(0)|$ . By (2.30) we see that on the set  $B_{\lambda c}^1([a, b]; R)$  the inequality (2.28) holds with  $q \equiv |F(0)|$ . Consequently, all the assumptions of Lemma 2.2 are satisfied and therefore the problem (0.1), (0.2) has at least one solution. It remains to show that the problem (0.1), (0.2) has at most one solution.

Let  $u_1, u_2$  be arbitrary solutions of the problem (0.1), (0.2). Put  $u(t) = u_1(t) - u_2(t)$  for  $t \in [a, b]$ . Then (2.29) and (2.30) yield

$$\begin{aligned} [u(a) - \lambda u(b)] \operatorname{sgn} u(a) &\leq 0, \\ [u'(t) - \ell(u)(t)] \operatorname{sgn} u(t) &\leq 0 \quad \text{for } t \in [a, b]. \end{aligned}$$

These two inequalities, together with the assumption  $\ell \in U(\lambda)$ , result in  $u \equiv 0$ . Consequently,  $u_1 \equiv u_2$ .  $\square$

**Lemma 2.4.** *Let  $\ell_0 \in \mathcal{L}_{ab}$  and the homogeneous problem*

$$v'(t) = \ell_0(v)(t), \quad v(a) - \lambda v(b) = 0$$

have only the trivial solution. Then there exists a positive number  $r_0$  such that for any  $q^* \in L([a, b]; R)$  and  $c \in R$ , the solution  $v$  of the problem

$$v'(t) = \ell_0(v)(t) + q^*(t), \quad v(a) - \lambda v(b) = c \quad (2.31)$$

admits the estimate

$$\|v\|_C \leq r_0 (|c| + \|q^*\|_L). \quad (2.32)$$

*Proof.* Let

$$R \times L([a, b]; R) = \{(c, q^*) : c \in R, q^* \in L([a, b]; R)\}$$

denote the Banach space with the norm

$$\|(c, q^*)\|_{R \times L} = |c| + \|q^*\|_L,$$

and  $\Omega$  be the operator which to every  $(c, q^*) \in R \times L([a, b]; R)$  assigns the solution  $v$  of the problem (2.31). According to Theorem 1.4 from [21],  $\Omega : R \times L([a, b]; R) \rightarrow C([a, b]; R)$  is a linear bounded operator. Let  $r_0$  be the norm of  $\Omega$ . Then, clearly, the inequality

$$\|\Omega(c, q^*)\|_C \leq r_0(|c| + \|q^*\|_L)$$

holds for arbitrary  $(c, q^*) \in R \times L([a, b]; R)$ . Consequently, the solution  $v = \Omega(c, q^*)$  of the problem (2.31) admits the estimate (2.32).  $\square$

**Lemma 2.5.** *Let  $\lambda \in [0, 1[$  and the operator  $\ell$  admit the representation  $\ell = \ell_0 - \ell_1$ , where  $\ell_0$  and  $\ell_1$  satisfy the condition (1.1). Let, moreover, there exist a function  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfying the inequalities (1.2), (1.3) and (2.4). Then  $\ell \in U(\lambda)$ .*

*Proof.* Let  $q^* \in L([a, b]; R_+)$ ,  $c \in R_+$  and  $u \in \tilde{C}([a, b]; R)$  satisfy the inequalities (2.24) and (2.25). It is clear that

$$u'(t) = \ell_0(u)(t) - \ell_1(u)(t) + \tilde{q}(t), \quad (2.33)$$

where

$$\tilde{q}(t) = u'(t) - \ell(u)(t) \quad \text{for } t \in [a, b].$$

Obviously,

$$\tilde{q}(t) \operatorname{sgn} u(t) \leq q^*(t) \quad \text{for } t \in [a, b]. \quad (2.34)$$

According to (1.1), (1.2), (1.3), and Lemma 1.1, we see that  $\ell_0 \in V^+(\lambda)$ . Therefore, the assumptions of Lemma 2.4 are fulfilled. Let  $r_0$  be the number appearing in Lemma 2.4 and put

$$r = r_0 \left( 1 + 4 \left( 1 + \gamma(b) - \gamma(a) \right) \left( 4 - (\gamma(b) - \gamma(a))^2 \right)^{-1} \right). \quad (2.35)$$

We will show that (2.26) holds with  $r$  defined by (2.35).

Let us first suppose that  $u$  does not change its sign. Then, in view of (1.1) and (2.34), the equality (2.33) yields

$$|u(t)|' \leq \ell_0(|u|)(t) + q^*(t) \quad \text{for } t \in [a, b],$$

and from (2.24) we get

$$|u(a)| - \lambda |u(b)| \leq c.$$

Therefore, by Remark 1.7, the condition  $\ell_0 \in V^+(\lambda)$  implies

$$|u(t)| \leq v(t) \quad \text{for } t \in [a, b],$$

where  $v$  is a solution of the problem (2.31). Due to Lemma 2.4, the function  $v$  admits the estimate (2.32), and so the estimate (2.26) holds.

Now assume that  $u$  changes its sign. Define numbers  $M$  and  $m$  by (1.24) and choose  $t_M, t_m \in [a, b]$  such that (1.25) holds. It is clear that (1.26) is fulfilled and either

$$t_M < t_m \quad (2.36)$$



or

$$t_m < t_M. \quad (2.37)$$

According to (1.1), (1.2), (1.3), (1.24), (1.26), and (2.31), we have

$$\begin{aligned} (M\gamma(t) + v(t))' &\geq \ell_0(M\gamma + v)(t) + M\ell_1(1)(t) + q^*(t) \\ &\geq \ell_0(M\gamma + v)(t) + \ell_1([u]_+)(t) + q^*(t) \quad \text{for } t \in [a, b], \\ M\gamma(a) + v(a) - \lambda(M\gamma(b) + v(b)) &\geq c, \end{aligned} \quad (2.38)$$

and

$$\begin{aligned} (m\gamma(t) + v(t))' &\geq \ell_0(m\gamma + v)(t) + m\ell_1(1)(t) + q^*(t) \\ &\geq \ell_0(m\gamma + v)(t) + \ell_1([u]_-)(t) + q^*(t) \quad \text{for } t \in [a, b], \\ m\gamma(a) + v(a) - \lambda(m\gamma(b) + v(b)) &\geq c. \end{aligned} \quad (2.39)$$

On the other hand, in view of (2.34) and (2.24), the equality (2.33) yields

$$\begin{aligned} [u(t)]'_+ &\leq \ell_0([u]_+)(t) + \ell_1([u]_-)(t) + q^*(t) \quad \text{for } t \in [a, b], \\ [u(a)]_+ - \lambda[u(b)]_+ &\leq c, \end{aligned} \quad (2.40)$$

and

$$\begin{aligned} [u(t)]'_- &\leq \ell_0([u]_-)(t) + \ell_1([u]_+)(t) + q^*(t) \quad \text{for } t \in [a, b], \\ [u(a)]_- - \lambda[u(b)]_- &\leq c. \end{aligned} \quad (2.41)$$

In view of the condition  $\ell_0 \in V^+(\lambda)$  and Remark 1.7, it follows from (2.38) and (2.41), and from (2.39) and (2.40), that

$$M\gamma(t) + v(t) \geq [u(t)]_- \quad \text{and} \quad m\gamma(t) + v(t) \geq [u(t)]_+ \quad \text{for } t \in [a, b]. \quad (2.42)$$

Inequalities (2.38)–(2.41), by virtue of (2.42) and the assumption  $\ell_0 \in \mathcal{P}_{ab}$ , imply

$$[u(t)]'_- \leq (M\gamma(t) + v(t))', \quad [u(t)]'_+ \leq (m\gamma(t) + v(t))' \quad \text{for } t \in [a, b]. \quad (2.43)$$

Note also that in view of the condition  $\ell_0 \in V^+(\lambda)$ , we have

$$v(t) \geq 0 \quad \text{for } t \in [a, b]. \quad (2.44)$$

Let us first suppose that (2.36) is fulfilled. On account of (1.25), (1.26), and (2.44), the integration of the first inequality in (2.43) from  $t_M$  to  $t_m$  results in

$$m \leq M\gamma(t_m) + v(t_m) - M\gamma(t_M) - v(t_M) \leq M(\gamma(t_m) - \gamma(t_M)) + \|v\|_C. \quad (2.45)$$

On the other hand, in view of (1.25), (1.26), and (2.44), the integration of the second inequality in (2.43) from  $a$  to  $t_M$  and from  $t_m$  to  $b$  yields

$$\begin{aligned} M - [u(a)]_+ &\leq m\gamma(t_M) + v(t_M) - m\gamma(a) - v(a) \\ &\leq m(\gamma(t_M) - \gamma(a)) - v(a) + \|v\|_C, \end{aligned} \quad (2.46)$$

$$[u(b)]_+ \leq m\gamma(b) + v(b) - m\gamma(t_m) - v(t_m) \leq m(\gamma(b) - \gamma(t_m)) + v(b). \quad (2.47)$$

Multiplying both sides of (2.47) by  $\lambda$  and taking into account that  $m > 0$ ,  $\lambda < 1$  and  $\gamma$  is a nondecreasing function, we obtain

$$\lambda[u(b)]_+ \leq m(\gamma(b) - \gamma(t_m)) + \lambda v(b).$$

Summing the last inequality and (2.46) and taking into account (2.31) and (2.40), we get

$$M \leq m(\gamma(t_M) - \gamma(t_m) + \gamma(b) - \gamma(a)) + \|v\|_C. \quad (2.48)$$

In view of (1.24), (2.36), and the condition  $\gamma'(t) \geq 0$  for  $t \in [a, b]$ , it follows from (2.45) and (2.48) that

$$\begin{aligned} \|u\|_C &\leq \|u\|_C(\gamma(t_m) - \gamma(t_M))(\gamma(t_M) - \gamma(t_m) + \gamma(b) - \gamma(a)) \\ &\quad + (1 + \gamma(b) - \gamma(a))\|v\|_C. \end{aligned}$$

Consequently, by virtue of the inequality

$$AB \leq \frac{1}{4}(A + B)^2, \quad (2.49)$$

$$\|u\|_C \leq \frac{\|u\|_C}{4}(\gamma(b) - \gamma(a))^2 + (1 + \gamma(b) - \gamma(a))\|v\|_C.$$

Hence, by (2.4),

$$\|u\|_C \leq 4(1 + \gamma(b) - \gamma(a))(4 - (\gamma(b) - \gamma(a))^2)^{-1}\|v\|_C. \quad (2.50)$$

Therefore, according to (2.32) and (2.35), the estimate (2.26) holds.

Now suppose that (2.37) is fulfilled. On account of (1.25), (1.26), and (2.44), the integration of the second inequality in (2.43) from  $t_m$  to  $t_M$  results in

$$\begin{aligned} M &\leq m\gamma(t_M) + v(t_M) - m\gamma(t_m) - v(t_m) \\ &\leq m(\gamma(t_M) - \gamma(t_m)) + \|v\|_C. \end{aligned} \quad (2.51)$$

On the other hand, in view of (1.25), (1.26), and (2.44), the integration of the first inequality in (2.43) from  $a$  to  $t_m$  and from  $t_M$  to  $b$  yields

$$\begin{aligned} m - [u(a)]_- &\leq M\gamma(t_m) + v(t_m) - M\gamma(a) - v(a) \\ &\leq M(\gamma(t_m) - \gamma(a)) - v(a) + \|v\|_C, \end{aligned} \quad (2.52)$$

$$[u(b)]_- \leq M\gamma(b) + v(b) - M\gamma(t_M) - v(t_M) \leq M(\gamma(b) - \gamma(t_M)) + v(b). \quad (2.53)$$

Multiplying both sides of (2.53) by  $\lambda$  and taking into account that  $M > 0$ ,  $\lambda < 1$  and  $\gamma$  is a nondecreasing function, we get

$$\lambda[u(b)]_- \leq M(\gamma(b) - \gamma(t_M)) + \lambda v(b).$$

Summing the last inequality and (2.52) and taking into account (2.31) and (2.41), we show that

$$m \leq M(\gamma(t_m) - \gamma(t_M) + \gamma(b) - \gamma(a)) + \|v\|_C. \quad (2.54)$$

Accordingg to (1.24), (2.37), and the condition  $\gamma'(t) \geq 0$  for  $t \in [a, b]$ , it follows from (2.51) and (2.54) that

$$\|u\|_{C^{1eq}} \|u\|_C (\gamma(t_M) - \gamma(t_m)) (\gamma(t_m) - \gamma(t_M) + \gamma(b) - \gamma(a)) + (1 + \gamma(b) - \gamma(a)) \|v\|_C.$$

Consequently, by virtue of (2.4) and (2.49), the inequality (2.50) is fulfilled. Therefore, according to (2.32) and (2.35), the estimate (2.26) holds.  $\square$

**2.3. Proofs of Main Results.** Theorem 2.1 follows from Lemmas 2.2 and 2.5, whereas Theorem 2.2 is a consequence Lemmas 2.3 and 2.5.

*Proof of Corollary 2.1.* Obviously, the condition (2.12) yields (2.3), where

$$\begin{aligned} F(v)(t) &\stackrel{def}{=} p(t)v(\tau(t)) - g(t)v(\mu(t)) + f(t, v(t), v(\nu(t))), \\ \ell_0(v)(t) &\stackrel{def}{=} p(t)v(\tau(t)), \quad \ell_1(v)(t) \stackrel{def}{=} g(t)v(\mu(t)) \quad \text{for } t \in [a, b]. \end{aligned} \quad (2.55)$$

Moreover, similarly to the proof of Corollary 1.1 one can show that, according to the conditions (1.10), (1.11), and (2.13), there exists a function  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfying the inequalities (1.2), (1.3), and (2.4). Therefore, the assumptions of Theorem 2.1 are satisfied.  $\square$

*Proof of Corollary 2.4.* Obviously, the condition (2.17) yields the condition (2.7) with  $F$ ,  $\ell_0$  and  $\ell_1$  defined by (2.55). Moreover, analogously to the proof of Corollary 1.2, according to the condition (2.14), where  $\alpha_1$  and  $\beta_1$  are defined by (1.14) and (1.15) with  $\sigma$  given by (1.16), one can show that there exists a function  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfying the inequalities (1.2), (1.3), and (2.4). Therefore, the assumptions of Theorem 2.2 are fulfilled.  $\square$

Corollaries 2.2, 2.3 and 2.5–2.8 can be proved in a similar manner.

### 3. ON REMARKS 1.2, 2.1 AND 2.2

**On Remark 1.2.** In Example 3.1, we have constructed an operator  $\ell \in \mathcal{L}_{ab}$  such that the homogeneous problem  $(0, 3_0)$ ,  $(0, 4_0)$  has a nontrivial solution. Then, according to Remark 1.1, there exist  $q_0 \in L([a, b]; R)$  and  $c_0 \in R$  such that the problem  $(0.3)$ ,  $(0.4)$  has no solution.

**Example 3.1.** Let  $\lambda \in [0, 1[$ ,  $a = 0$ ,  $b = 4$ ,  $\varepsilon \geq 0$ , and let

$$\ell_0 \equiv 0, \quad \ell_1(v)(t) \stackrel{def}{=} g(t)v(\mu(t)) \quad \text{for } t \in [a, b], \quad (3.1)$$

where

$$g(t) = \begin{cases} 1 + \lambda & \text{for } t \in [0, 1[ \\ 1 & \text{for } t \in [1, 3[ \\ \varepsilon & \text{for } t \in [3, 4] \end{cases}, \quad \mu(t) = \begin{cases} 3 & \text{for } t \in [0, 1[ \\ 1 & \text{for } t \in [1, 3[ \\ 2 & \text{for } t \in [3, 4] \end{cases}.$$

Obviously,  $\|g\|_L = 3 + \lambda + \varepsilon$ . Choose  $\delta > 0$  such that  $\delta > \frac{\lambda\|g\|_L}{1-\lambda}$  and define the function  $\gamma$  by

$$\gamma(t) = \delta + \int_a^t g(s)ds \quad \text{for } t \in [a, b]. \tag{3.2}$$

It is clear that  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfies the conditions (1.2), (1.3), and

$$\gamma(b) - \gamma(a) = 3 + \lambda + \varepsilon.$$

On the other hand, the problem  $(0, 3_0)$ ,  $(0, 4_0)$  has a nontrivial solution

$$u(t) = \begin{cases} \lambda - (1 + \lambda)t & \text{for } t \in [0, 1[ \\ t - 2 & \text{for } t \in [1, 3[ \\ 1 & \text{for } t \in [3, 4] \end{cases} .$$

Example 3.1 shows that the strict inequality (1.4) in Theorem 1.2 cannot be replaced by nonstrict one. This example also shows that the strict inequalities (1.12) in Corollary 1.1 and (1.13) in Corollary 1.2 cannot be replaced by nonstrict ones.

**On Remark 2.1.** In Example 3.2 functions  $g, z \in L([a, b]; R_+)$  and  $\mu \in \mathcal{M}_{ab}$  are constructed such that the problem

$$u'(t) = -g(t)u(\mu(t)) - z(t)u(t), \quad u(a) - \lambda u(b) = 0 \tag{3.3}$$

has a nontrivial solution. Then, by Remark 1.1, there exist  $q_0 \in L([a, b]; R)$  and  $c_0 \in R$  such that the problem (0.1), (0.2), with

$$F(v)(t) \stackrel{def}{=} -g(t)v(\mu(t)) - z(t)v(t) + q_0(t) \quad \text{for } t \in [a, b], \quad h(v) \stackrel{def}{=} c_0,$$

has no solution, while the conditions (2.2) and (2.3) are fulfilled, where  $\ell_0, \ell_1$  are defined by (3.1),  $q \equiv |q_0|$ , and  $c = |c_0|$ .

**Example 3.2.** Let  $\lambda \in [0, 1[$ ,  $\varepsilon > 0$ , and choose  $\eta_1 \in [0, \lambda]$  such that  $0 < \eta_1 < \lambda$  if  $\lambda \neq 0$ , and  $\eta_2 \in ]0, 1[$  such that  $\eta_1 + \eta_2 \leq \varepsilon$ . Put  $a = 0$ ,  $b = 5$ ,  $t_0 = \frac{\eta_1}{1+\eta_1} + 1$ ,

$$g(t) = \begin{cases} 0 & \text{for } t \in [0, 1[ \cup [2, 3[ \\ 1 + \eta_1 & \text{for } t \in [1, 2[ \\ 1 + \eta_2 & \text{for } t \in [3, 4[ \\ \varepsilon - \eta_1 - \eta_2 & \text{for } t \in [4, 5] \end{cases}, \quad \mu(t) = \begin{cases} 5 & \text{for } t \in [0, 2[ \\ 2 & \text{for } t \in [2, 4[ \\ t_0 & \text{for } t \in [4, 5] \end{cases},$$

$$z(t) = \begin{cases} z_0(t) & \text{for } t \in [0, 1[ \\ 0 & \text{for } t \in [1, 2[ \cup [3, 5] \\ \frac{(1 - \eta_2)}{(1 - \eta_2)(2 - t) + 1} & \text{for } t \in [2, 3[ \end{cases},$$

where

$$z_0(t) = \begin{cases} 0 & \text{if } \lambda = 0 \\ \frac{\lambda - \eta_1}{\lambda - (\lambda - \eta_1)t} & \text{if } \lambda \neq 0 \end{cases} \quad \text{for } t \in [0, 1[.$$

Obviously,  $\|g\|_L = 2 + \varepsilon$ . Choose  $\delta > 0$  such that  $\delta > \frac{\lambda \|g\|_L}{1 - \lambda}$  and define the function  $\gamma$  by (3.2). It is clear that  $\gamma \in \tilde{C}([a, b]; ]0, +\infty[)$  satisfies the conditions (1.2), (1.3), and

$$\gamma(b) - \gamma(a) = 2 + \varepsilon.$$

On the other hand, the problem (3.3) has a nontrivial solution

$$u(t) = \begin{cases} (\lambda - \eta_1)t - \lambda & \text{for } t \in [0, 1[ \\ (\eta_1 + 1)(t - 1) - \eta_1 & \text{for } t \in [1, 2[ \\ (1 - \eta_2)(2 - t) + 1 & \text{for } t \in [2, 3[ \\ (1 + \eta_2)(3 - t) + \eta_2 & \text{for } t \in [3, 4[ \\ -1 & \text{for } t \in [4, 5] \end{cases}.$$

Example 3.2 shows that the strict inequality (2.4) in Theorem 2.1 cannot be replaced by the inequality (2.5), no matter how small  $\varepsilon > 0$  is. Example 3.2 also shows that the strict inequalities (2.13) in Corollary 2.1 and (2.14) in Corollary 2.2 cannot be replaced by the inequalities (2.15) and (2.16), no matter how small  $\varepsilon > 0$  is.

**On Remark 2.2.** Example 3.2 shows that the strict inequality (2.4) in Theorem 2.2, resp. (2.13) in Corollary 2.3, resp. (2.14) Corollary 2.4, cannot be replaced by the inequality (2.5), resp. (2.15), resp. (2.16), no matter how small  $\varepsilon > 0$  is.

#### ACKNOWLEDGEMENT

For the first author this work was supported by Grant No. 201/00/D058 of the Grant Agency of the Czech Republic, for the second author by the RI No. J07/98:143100001 and for the third author by Grant No. 201/99/0295 of the Grant Agency of the Czech Republic.

#### REFERENCES

1. N. V. AZBELEV, V. P. MAKSIMOV, and L. F. RAKHMATULLINA, Introduction to the theory of functional differential equations. (Russian) *Nauka, Moscow*, 1991.
2. N. V. AZBELEV and L. F. RAKHMATULLINA, Theory of linear abstract functional differential equations and applications. *Mem. Differential Equations Math. Phys.* **8**(1996), 1–102.
3. S. R. BERNFELD and V. LAKSHMIKANTHAM, An introduction to nonlinear boundary value problems. *Academic Press, Inc., New York and London*, 1974.
4. E. BRAVYI, A note on the Fredholm property of boundary value problems for linear functional differential equations. *Mem. Differential Equations Math. Phys.* **20**(2000), 133–135.

5. E. BRAVYI, R. HAKL, and A. LOMTATIDZE, Optimal conditions for unique solvability of the Cauchy problem for first order linear functional differential equations. *Czechoslovak Math. J.* (to appear).
6. E. BRAVYI, R. HAKL, and A. LOMTATIDZE, On Cauchy problem for the first order nonlinear functional differential equations of non-Volterra's type. *Czechoslovak Math. J.* (to appear).
7. E. A. Coddington and N. LEVINSON, Theory of ordinary differential equations. *Mc-Graw-Hill Book Company, Inc., New York-Toronto-London*, 1955.
8. S. M. GELASHVILI, On a boundary value problem for systems of functional differential equations. (Russian) *Arch. Math. (Brno)* **20**(1964), 157–168.
9. R. HAKL, On some boundary value problems for systems of linear functional-differential equations. *Electron. J. Qual. Theory Differ. Equ.* **1999**, No. 10, 1–16.
10. R. HAKL, I. KIGURADZE, and B. PŮŽA, Upper and lower solutions of boundary value problems for functional differential equations and theorems on functional differential inequalities. *Georgian Math. J.* **7**(2000), No. 3, 489–512.
11. R. HAKL and A. LOMTATIDZE, A note on the Cauchy problem for first order linear differential equations with a deviating argument. *Arch. Math. (Brno)* **38**(2002), No. 1, 61–71.
12. R. HAKL, A. LOMTATIDZE, and B. PŮŽA, On nonnegative solutions of first order scalar functional differential equations. *Mem. Differential Equations Math. Phys.* **23**(2001), 51–84.
13. R. HAKL, A. LOMTATIDZE, and B. PŮŽA, New optimal conditions for unique solvability of the Cauchy problem for first order linear functional differential equations. *Math. Bohemica* (to appear).
14. R. HAKL, A. LOMTATIDZE, and J. ŠREMR, On nonnegative solutions of a periodic type boundary value problem for first order scalar functional differential equations. *Adv. Differential Equations* (submitted).
15. R. HAKL, A. LOMTATIDZE, and J. ŠREMR, On constant sign solutions of a periodic type boundary value problem for first order scalar functional differential equations. *Mem. Differential Equations Math. Phys.* **26**(2002), 65–92.
16. J. HALE, Theory of functional differential equations. *Springer-Verlag, New York-Heidelberg-Berlin*, 1977.
17. P. HARTMAN, Ordinary differential equations. *John Wiley & Sons, Inc., New York-London-Sydney*, 1964.
18. I. KIGURADZE, Some singular boundary value problems for ordinary differential equations. (Russian) *Tbilisi Univ. Press, Tbilisi*, 1975.
19. I. KIGURADZE, Boundary value problems for systems of ordinary differential equations. Current problems in mathematics. Newest results. (Russian) *Itogi Nauki i Tekhniki, Current problems in mathematics. Newest results*, 30 (Russian), 3–103, 204, *Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow*, 1987; Translated in *J. Soviet Math.* **43**(1988), No. 2, 2259–2339.
20. I. KIGURADZE, Initial and boundary value problems for systems of ordinary differential equations, I. (Russian) *Metsniereba, Tbilisi*, 1997.
21. I. KIGURADZE and B. PŮŽA, On boundary value problems for systems of linear functional differential equations. *Czechoslovak Math. J.* **47**(1997), No. 2, 341–373.
22. I. KIGURADZE and B. PŮŽA, On a certain singular boundary value problem for linear differential equations with deviating arguments. *Czechoslovak Math. J.* **47**(1997), 233–244.

23. I. KIGURADZE and B. PŮŽA, Conti–Opial type theorems for systems of functional differential equations. (Russian) *Differentsial'nye Uravneniya* **33**(1997), No. 2, 185–194.
24. I. KIGURADZE and B. PŮŽA, On boundary value problems for functional differential equations. *Mem. Differential Equations Math. Phys.* **12**(1997), 106–113.
25. I. KIGURADZE and B. PŮŽA, On periodic solutions of systems of linear functional differential equations. *Arch. Math. (Brno)* **33**(1997), No. 3, 197–212.
26. I. KIGURADZE and B. PŮŽA, On the solvability of boundary value problems for systems of nonlinear differential equations with deviating arguments. *Mem. Differential Equations Math. Phys.* **10**(1997), 157–161.
27. I. KIGURADZE and B. PŮŽA, On a singular two–point boundary value problem for the nonlinear  $m$ -th order differential equation with deviating arguments. *Georgian Math. J.* **4**(1997), No. 6, 557–566.
28. I. KIGURADZE and B. PŮŽA, On the solvability of nonlinear boundary value problems for functional differential equations. *Georgian Math. J.* **5**(1998), No. 3, 251–262.
29. V. KOLMANOVSKII and A. MYSHKIS, Introduction to the theory and applications of functional differential equations. *Mathematics and its Applications*, 463, *Kluwer Academic Publishers, Dordrecht*, 1999.
30. Š. SCHWABIK, M. TVRDÝ, and O. VEJVODA, Differential and integral equations: boundary value problems and adjoints. *Academia, Praha*, 1979.

(Received 30.01.2002)

Authors' addresses:

R. Hakl  
Mathematical Institute  
Czech Academy of Sciences  
Žižkova 22, 616 62 Brno  
Czech Republic  
E-mail: hakl@ipm.cz

A. Lomtadze and J. Šremr  
Department of Mathematical Analysis  
Faculty of Science, Masaryk University  
Janáčkovo nám. 2a, 662 95 Brno  
Czech Republic  
E-mail: bacho@math.muni.cz  
sremr@math.muni.cz