

ON THE EQUIVALENCE OF QUILLEN'S AND SWAN'S *K*-THEORIES

PAUL EBELING AND FRANS KEUNE

Abstract. The *K*-theory of rings can be defined in terms of nonabelian derived functors as described in [9]; see also the books [7] and [8] of Inassaridze for a similar approach. In fact both Swan's theory and Quillen's theory can be described this way. The equivalence of both *K*-theories is proved by Gersten [5]. In this paper we give a proof using these descriptions that involve nonabelian derived functors.

2000 Mathematics Subject Classification: 19D50.

Key words and phrases: Higher algebraic *K*-theory, simplicial resolutions.

INTRODUCTION

Quillen's higher algebraic *K*-groups of a unital ring *R* are defined as the homotopy groups of the space that is obtained from the classifying space of the elementary group of the ring *R*:

$$K_n(R) = \pi_n(BE(R)^+) \quad (\text{for } n \geq 2).$$

In Section 4 we replace this space by a simplicial set $\mathbb{Z}_\infty \bar{W}E(R)$ also depending functorially on *R*, having a geometric realization which is homotopy equivalent to $BE(R)^+$. This description depends on the notion of integral completion in the sense of Bousfield and Kan [3].

Swan's *K*-groups of a (nonunital) ring *R* are defined by means of a free simplicial resolution of *R*. In Section 8 we consider a simplicial group *H*(*R*) depending functorially on *R*, having Swan's *K*-groups of *R* as homotopy groups. Writing $K'_n(R)$ for these groups the formula becomes

$$K'_n(R) = \pi_{n-2}(H(R)).$$

The main result is the theorem in Section 8, which says that Swan's *K*-groups coincide with Quillen's when extended to the category of nonunital rings in the standard way: $K'_n(R) \cong K_n(R^+, R)$. Here R^+ stands for the ring obtained from *R* by formally adjoining a unity element. The proof uses Gersten's result [5]: free associative nonunital rings have trivial *K*-theory which historically was the missing part of the earliest proof of the equivalence of these *K*-theories by Anderson [1].

It should be noted that the proof in Section 8 is a corrected and improved version of the proof in the unpublished paper [11].

1. THE PLUS CONSTRUCTION

The so-called plus construction is used in Quillen's definition of the higher K -groups of a unital ring. Its defining properties are described in the theorem below; see also Property P1. Good references for the plus construction are Loday's thesis [13] and the book by Berrick [2].

Theorem and definition ([13], Théorème 1.1.1). *Let X be a connected CW-complex with basepoint $*$, and let N be a perfect normal subgroup of π ($= \pi_1(X, *)$). Then there exists a connected CW-complex X^+ and a map $j: X \rightarrow X^+$ such that*

- (i) $\pi_1(j): \pi_1(X, *) \rightarrow \pi_1(X^+, j(*))$ identifies with the canonical projection $\pi \rightarrow \pi/N$.
- (ii) The map j induces an isomorphism on integral homology.

In the sequel the following properties will be used.

Property P1 ([13], Proposition 1.1.2). *The pair (X^+, j) is universal (in the homotopy category) among pairs (Y, f) , where Y is a connected space and $f: X \rightarrow Y$ a continuous map satisfying $\pi_1(f)(N) = 0$.*

A consequence is that the plus construction is functorial up to homotopy ([13], Corollaire 1.1.3).

Property P2 ([13], Proposition 1.1.7). *Let G be a group with a perfect commutator subgroup $[G, G]$. Then $B[G, G]^+$ is up to homotopy the universal covering of BG^+ , where in both cases the plus construction is relative to the subgroup $[G, G]$. (As usual BG is the classifying space of the group G .)*

The plus construction is used in the definition of higher K -groups as follows. Let R be a unital ring. The general linear group $GL(R)$ of R has the elementary subgroup $E(R)$ as the commutator subgroup (the 'Whitehead Lemma'). Apply the plus construction to the classifying space $BGL(R)$ of $GL(R)$ relative to the subgroup $E(R)$ (which is perfect), and finally take homotopy groups.

Definition. $K_n(R) = \pi_n(BGL(R)^+)$ for $n \geq 1$.

From Property P2 one deduces

$$K_n(R) = \pi_n(BE(R)^+) \quad \text{for } n \geq 2$$

(and of course $\pi_1(BE(R)^+) = 0$). In this paper this identity is taken as definition of K_n for $n \geq 2$. An important property of the space $BE(R)^+$ is the following.

Property P3. $BE(R)^+$ is an H -space. ([13], §1.3.4.)

2. THE INTEGRAL COMPLETION OF A GROUP

In Section 4 the space $BE(R)^+$ will be replaced by the integral completion (in the sense of Bousfield and Kan [3]) of $BE(R)$. The definition and properties of this completion are given in the next section. We will need the notion of integral completion of a group.

Let G be a group. The subgroups $\Gamma_i G$ (for $i \geq 1$) are defined inductively by

$$\begin{aligned} \Gamma_1 G &= G, \\ \Gamma_{i+1} G &= [\Gamma_i G, G]. \end{aligned}$$

As usual $[H_1, H_2]$ denotes the subgroup generated by the commutators $[h_1, h_2] = h_1 h_2 h_1^{-1} h_2^{-1}$ with $h_1 \in H_1$ and $h_2 \in H_2$. Clearly, the subgroups $\Gamma_i G$ are normal subgroups of G . The series of subgroups

$$G \supseteq \Gamma_2 G \supseteq \Gamma_3 G \cdots$$

is known as the *lower central series* of the group G . It induces a tower of groups

$$\cdots \rightarrow G/\Gamma_3 G \rightarrow G/\Gamma_2 G \rightarrow 1$$


in which every group homomorphism $G/\Gamma_{i+1} G \rightarrow G/\Gamma_i G$ is a central extension.

The *integral completion* CG ($= G_{\mathbb{Z}}^{\wedge}$ in the terminology of [3]) of G is defined as the inverse limit of the tower of groups:

$$CG = \varprojlim_i G/\Gamma_i G.$$

It was pointed out to the authors that this construction also appears in the literature under the name “pronilpotent completion”.

In an obvious way C is a functor $\mathbf{Gr} \rightarrow \mathbf{Gr}$, where \mathbf{Gr} denotes the category of groups. This functor can be viewed as the inverse limit of the functor which assigns to every homomorphism $G \rightarrow N$, with N a nilpotent group, the group N , and to every commutative triangle



with N and N' nilpotent, the map $N \rightarrow N'$. The existence of this inverse limit follows from the existence of small cofinal diagrams, e.g. given by the tower of groups above.

3. THE INTEGRAL COMPLETION OF A SPACE

We will use here the simplicial terminology. For a category \mathbf{C} the category of simplicial \mathbf{C} -objects is denoted by \mathbf{sC} . The category of reduced simplicial sets is denoted by \mathbf{rsSet} . It is the full subcategory of \mathbf{sSet} , the category of simplicial sets, consisting of those $X \in \mathbf{sSet}$ which have only one vertex, i.e. X_0 is a one-element set.

The functor $G: \mathbf{rsSet} \rightarrow \mathbf{sGr}$ assigns to a reduced simplicial set its loop group GX , which is a simplicial group satisfying $\pi_i(GX) \cong \pi_{i+1}(X)$ for all $i \geq 0$. This functor G has a right adjoint $\bar{W}: \mathbf{sGr} \rightarrow \mathbf{rsSet}$, which is the simplicial analogue of the classifying space functor.

The reduced simplicial set $\bar{W}H$ is called the *classifying complex* of the simplicial group H . The adjunction of G and \bar{W} induces a natural simplicial map

$X \rightarrow \bar{W}GX$, which induces isomorphisms on the homotopy groups when X is a Kan complex. A good reference for this is May [14].

A functor $T: \mathbf{Gr} \rightarrow \mathbf{Gr}$ determines a functor $\tilde{T}: \mathbf{rsSet} \rightarrow \mathbf{rsSet}$ in the following way: let X be a reduced simplicial set; first form its loop group GX , next apply T dimension-wise to obtain a simplicial group TGX , and finally take the classifying complex. In a formula: $\tilde{T} = \bar{W}TG$, where T stands for T applied dimension-wise. In particular the integral completion $C: \mathbf{Gr} \rightarrow \mathbf{Gr}$ as defined in Section 2 determines a functor

$$\mathbb{Z}_\infty = \tilde{C}: \mathbf{rsSet} \rightarrow \mathbf{rsSet}.$$

This functor assigns to a reduced simplicial set its so-called *integral completion*. This integral completion functor was introduced by Bousfield and Kan [3]. The definition given here is in fact one of various possible definitions: it is the definition they give in Chapter III of [3]. Some of the main properties of the integral completion functor are:

Property I1 ([3], Ch. I, Lemma 5.5, p. 25). *Let $f: X \rightarrow Y$ be a map in \mathbf{rsSet} . Then $\mathbb{Z}_\infty f: \mathbb{Z}_\infty X \rightarrow \mathbb{Z}_\infty Y$ is a homotopy equivalence if and only if f induces an isomorphism on integral homology.*

Property I2 ([3], Ch. V, Proposition 3.4, p. 134). *For $X \in \mathbf{rsSet}$ there is a natural map $i: X \rightarrow \mathbb{Z}_\infty X$ which is a weak homotopy equivalence if X is nilpotent.*

Property I3 ([3], Ch. II, Lemma 5.4, p. 63). *Let $p: E \rightarrow B$ (in \mathbf{rsSet}) be a fibration with connected fibre F such that the Serre action of $\pi_1(B)$ on $H_i(F; \mathbb{Z})$ is nilpotent for all $i \geq 0$. Then $\mathbb{Z}_\infty(p): \mathbb{Z}_\infty E \rightarrow \mathbb{Z}_\infty B$ is a fibration and the inclusion $\mathbb{Z}_\infty F \rightarrow \mathbb{Z}_\infty(p)^{-1}(*)$ is a homotopy equivalence ([3], Ch. II, Lemma 5.1, p. 62). The action of $\pi_1(B)$ on $H_*(F; \mathbb{Z})$ is in particular nilpotent if $\pi_1(E)$ acts nilpotently on $\pi_i(F)$ for all $i \geq 1$.*

4. THE INTEGRAL COMPLETION OF $\bar{W}E(R)$

For any group H , there is a simplicial group which is H in every dimension, having the identity as degeneracy and boundary maps. This object is a *constant simplicial group* and we denote it by H again.

Proposition 1. *Let R be a unital ring. Then the geometric realization $|\mathbb{Z}_\infty \bar{W}E(R)|$ of $\mathbb{Z}_\infty \bar{W}E(R)$ is homotopy equivalent to $BE(R)^+$, the equivalence being functorial in R .*

Proof. The plus construction has its simplicial analogue in \mathbf{rsSet} , the category of reduced simplicial sets. There exists a map $j: \bar{W}E(R) \rightarrow \bar{W}E(R)^+$ in \mathbf{rsSet} such that its geometric realization $|j|: BE(R) \rightarrow |\bar{W}E(R)^+|$ is the map $j: BE(R) \rightarrow BE(R)^+$ of Section 1. Consider the commutative square (which

is functorial in R)

$$\begin{CD} \bar{W}E(R) @>j>> \bar{W}E(R)^+ \\ @V i VV @VV i V \\ \mathbb{Z}_\infty \bar{W}E(R) @>\mathbb{Z}_\infty(j)>> \mathbb{Z}_\infty(\bar{W}E(R)^+) \end{CD}$$

It suffices to prove that $i: \bar{W}E(R)^+ \rightarrow \mathbb{Z}_\infty(\bar{W}E(R)^+)$ and $\mathbb{Z}_\infty(j)$ are homotopy equivalences. The first map is a homotopy equivalence because of Property I2 and Property P3: the space $BE(R)^+$ is an H -space, so it is nilpotent. The map $\mathbb{Z}_\infty(j)$ is a homotopy equivalence because of Property I1. \square

The proof above can also be found in [5], where it is attributed to E. Dror. In exactly the same way one proves the homotopy equivalence of the spaces $|\mathbb{Z}_\infty \bar{W}GL(R)|$ and $BGL(R)^+$.

Corollary. For $n \geq 2$ we have $K_n \cong \pi_n \mathbb{Z}_\infty \bar{W}E$.

5. DERIVED FUNCTORS

Let \mathbf{Gr} be the category of groups. In this section we will review the theory of (left) derived functors of a given functor $T: \mathbf{Gr} \rightarrow \mathbf{Gr}$ as introduced in [9]. The situation is analogous to the Abelian case where projective resolutions are used.

Let G be a simplicial group. For each $n \geq 1$ we define

$$Z_n(G) = \{ (x_0, \dots, x_{n+1}) \in G_n^{n+2} \mid d_i x_j = d_{j-1} x_i \text{ for all } 0 \leq i < j \leq n + 1 \},$$

a subgroup of G_n^{n+2} ($= G_n \times \dots \times G_n$, $n + 2$ times). The elements of $Z_n(G)$ are those $n + 2$ -tuples of elements of G_n that fit together in exactly the same way as the $n + 2$ faces of an $n + 1$ -simplex do. The group $Z_n(G)$ will be called the *group of n -spheres* in the simplicial group G . There is an obvious homomorphism $d: G_{n+1} \rightarrow Z_n(G)$ which assigns to an $n + 1$ -simplex x the n -sphere $(d_0 x, \dots, d_{n+1} x)$ of its faces. A simplicial group is called *aspherical* if for each $n \geq 1$ the map $d: G_{n+1} \rightarrow Z_n(G)$ is surjective, that is if every n -sphere is the boundary of an $n + 1$ -simplex.

For $n > 0$ there is an isomorphism

$$\bar{\alpha}: \pi_n(G) \rightarrow Z_n(G)/dG_{n+1},$$

which is induced by the homomorphism

$$\alpha: \tilde{G}_n \rightarrow Z_n(G), \quad g \mapsto (1, \dots, 1, g),$$

where $\tilde{G}_n = \bigcap_i \text{Ker } d_i$.

The isomorphism $\bar{\alpha}$ has the useful property that it respects the natural action of G_0 on G , given by conjugating G dimension-wise by the images of elements of G_0 under the degeneracy maps. The action obviously induces actions on \tilde{G}_n and $Z_n(G)$ by restricting the actions on G_n and the $n + 2$ -fold product $G_n \times \dots \times G_n$ respectively.

For any set X let FX be the free group on the elements of X . A simplicial group G is called *free* if there is a subset X_n of G_n for each $n \geq 0$ such that $G \cong FX_n$ and moreover the degeneracy maps $s_i: G_n \rightarrow G_{n+1}$ ($i = 0, \dots, n$) map X_n into X_{n+1} for each $n \geq 0$. An example of a free simplicial group is the loop group GX of a reduced simplicial set X .

Let H be a group. A *free resolution* (G, ε) , or simply G , of H consists of:

- (1) a free aspherical simplicial group G ;
- (2) a group homomorphism $\varepsilon: G_0 \rightarrow H$, which induces an isomorphism $\pi_0(G) \rightarrow H$.

Free resolutions do exist. What is more, there are functorial free resolutions $G(H)$. One example is the cotriple resolution $G_n(H) = F^{n+1}(H)$. Another example is $G\bar{W}(H)$, the loop group on the classifying complex of H .

Let G be a free resolution of a group H and G' a free resolution of a group H' . In [9] it is proved that a homomorphism $h: H \rightarrow H'$ can be covered by a simplicial homomorphism $g: G \rightarrow G'$, i.e. $\pi_0(g): \pi_0(G) \rightarrow \pi_0(G')$ induces f via the isomorphisms $\pi_0(G) \rightarrow H$ and $\pi_0(G') \rightarrow H'$. Moreover, two such simplicial homomorphisms are **Gr**-homotopic. As a consequence one can define derived functors of a functor $T: \mathbf{Gr} \rightarrow \mathbf{Gr}$. On objects the n -th derived functor $L_n T: \mathbf{Gr} \rightarrow \mathbf{Gr}$ is defined as follows. Let $H \in \mathbf{Gr}$; take a free resolution G of H ; then put

$$L_n T = \pi_n(TG),$$

where TG means: T applied dimension-wise to G . On morphisms $L_n T$ is defined by

$$(L_n T)(h) = \pi_n(Tg),$$

where $g: G \rightarrow G'$ covers $h: H \rightarrow H'$, G and G' being free resolutions of H and H' respectively.

Example. Let $T: \mathbf{Gr} \rightarrow \mathbf{Gr}$ be the Abelianization functor. What are its derived functors? $G\bar{W}H$ is a free resolution of $H \in \mathbf{Gr}$. Therefore

$$\begin{aligned} (L_n T)(H) &= \pi_n(TG\bar{W}H) = \pi_n(G\bar{W}H/[G\bar{W}H, G\bar{W}H]) \\ &= H_{n+1}(\bar{W}H; \mathbb{Z}) = H_{n+1}(H; \mathbb{Z}) \end{aligned}$$

(cf. [14], p.121). So $L_n T$ is the homology functor $H_{n+1}(-; \mathbb{Z})$.

One can also consider functors $T: \mathbf{Rg} \rightarrow \mathbf{Gr}$ on the category **Rg** of nonunital rings instead of **Gr**. The role of the free group is then taken over by the free ring: for X a set FX is the ring of polynomials without constant term in the non-commuting variables $x \in X$, with coefficients in \mathbb{Z} . Analogously one then considers: free simplicial rings, rings of n -spheres in a simplicial ring, etc. Then too one has a theory of derived functors for functors from **Rg** to **Gr**. More generally, the procedure is applicable to functors $T: \mathbf{A} \rightarrow \mathbf{Set}_*$, where **A** is a category of triple algebras and \mathbf{Set}_* the category of pointed sets. In fact this is how the theory is presented in [9].

6. DERIVED FUNCTORS OF THE INTEGRAL COMPLETION

We will consider the derived functor $L_n C$ of $C: \mathbf{Gr} \rightarrow \mathbf{Gr}$, the integral completion functor as described in Section 2. Let H be a group. Note that for any simplicial set X the simplicial group GX is free. It follows that the simplicial group $G\bar{W}H$ is a free resolution of H . The groups $(L_n C)(H)$ are therefore the homotopy groups of $CG\bar{W}H$, a simplicial group which has as classifying complex the integral completion of $\bar{W}H$ (cf. Section 3):

$$\mathbb{Z}_\infty \bar{W}H = \bar{W}CG\bar{W}H.$$

Hence

$$(L_n C)(H) = \pi_n(CG\bar{W}H) \cong \pi_{n+1}(\bar{W}CG\bar{W}H) = \pi_{n+1}(\mathbb{Z}_\infty \bar{W}H).$$

In the special case $H = GL(R)$ with R a unital ring, we obtain

Lemma. *For each $n \geq 0$ there is a canonical isomorphism*

$$(L_n C)(GL(R)) \cong \pi_{n+1}(\mathbb{Z}_\infty \bar{W}GL(R)) \cong K_{n+1}(R).$$

7. DERIVED FUNCTORS OF GL

In [9] higher K -functors were defined as derived functors of $GL: \mathbf{Rg} \rightarrow \mathbf{Gr}$ by the formula

$$K'_n(R) = L_{n-2}GL \quad (n \geq 3)$$

and K'_1 and K'_2 are then defined by the exactness of

$$1 \rightarrow K'_2 \rightarrow L_0GL \rightarrow GL \rightarrow K'_1 \rightarrow 1.$$

Since $\text{St}(FX) \cong GL(FX)$ for free rings FX , we have $L_0GL = L_0\text{St}$, where St denotes the Steinberg group. It is easily seen that $L_0\text{St} = \text{St}$, i.e. St is a right exact functor, see [10] for details. Hence the exact sequence above becomes

$$1 \rightarrow K'_2 \rightarrow \text{St} \rightarrow GL \rightarrow K'_1 \rightarrow 1,$$

which shows that the functors K'_1 and K'_2 coincide with the classical ones. It will be proven in Section 8 that the functors as defined above are isomorphic to the functors K_n as defined in Section 1.

Remark. The groups $K'_n(R)$ defined in this section coincide with the groups $K_n(R)$ as defined by Gersten [4], since Gersten uses the cotriple resolution of R for their definition, which is simply one of possible resolutions of R . In [17] Swan proved that his functor K_n , which he defined in [16], coincides with Gersten's.

8. COMPARISON OF BOTH K -THEORIES

In this section we prove the main theorem.

Theorem. *Let $R \in \mathbf{Rg}$. Then for all $n \geq 2$,*

$$K'_n(R) \cong K_n(R^+, R).$$

Let R be a nonunital ring. Form the simplicial ring FR by applying the free resolution functor. Adjoining a unit in every dimension of FR we get a split homomorphism of simplicial unital rings,

$$(FR)^+ \begin{array}{c} \longrightarrow \\ \longleftarrow \end{array} \mathbb{Z},$$

where the right-hand side is interpreted as a constant simplicial ring.

The adjunction of G and \bar{W} induces a cotriple on \mathbf{sGr} . By first applying this cotriple resolution to the diagram $E((FR)^+) \rightarrow E(\mathbb{Z})$, and then the integral completion functor C , we define the two following bisimplicial groups together with a split homomorphism.

$$Q_{pq}^+ = (C(G\bar{W})^{q+1}E((FR)^+))_p \longrightarrow Z_{pq} = (C(G\bar{W})^{q+1}E(\mathbb{Z}))_p$$

Let Q denote the fibre (or kernel) of this map. Taking homotopy in each row of these bisimplicial groups, we can consider the usual long exact sequence of a fibration. Since the homomorphism splits, this sequence of simplicial groups degenerates into split short exact sequences

$$1 \longrightarrow \pi_q^h Q \longrightarrow \pi_q^h Q^+ \begin{array}{c} \longrightarrow \\ \longleftarrow \end{array} \pi_q^h Z \longrightarrow 1.$$

Again computing homotopy, each of these fibrations induces a long exact sequence, which too is split. Hence for every p and q we have a split short exact sequence

$$1 \longrightarrow \pi_p^v \pi_q^h Q \longrightarrow \pi_p^v \pi_q^h Q^+ \begin{array}{c} \longrightarrow \\ \longleftarrow \end{array} \pi_p^v \pi_q^h Z \longrightarrow 1.$$

Repeating this process, but now taking vertical homotopy groups first, we have

$$1 \longrightarrow \pi_q^h \pi_p^v Q \longrightarrow \pi_q^h \pi_p^v Q^+ \begin{array}{c} \longrightarrow \\ \longleftarrow \end{array} \pi_q^h \pi_p^v Z \longrightarrow 1.$$

Note that for each p , the simplicial group Q_{p*}^+ is the result of an application of the functor C to a free \mathbf{Gr} -resolution of $E((F_p R)^+)$. Hence,

$$\pi_q^h \pi_p^v Q^+ = (L_p C)(E((F_q R)^+)) = \pi_{p+1} \mathbb{Z}_\infty \bar{W} E((F_q R)^+).$$

Using the description of the Quillen K -groups from Section 6, it follows that for all p and q we have a short exact sequence

$$0 \longrightarrow \pi_q^h \pi_p^v Q \longrightarrow K_{p+1}((F_q R)^+) \longrightarrow K_{p+1}(\mathbb{Z}) \longrightarrow 0.$$

Using Gersten’s theorem on the K -theory of free rings [5], we find that for all p and q

$$\pi_q^h \pi_p^v Q = 0,$$

and hence, e.g. by Quillen’s spectral sequence [15] we also have

$$\pi_p^v \pi_q^h Q = 0.$$

By letting $p = 0$ and applying this formula to the relevant split exact sequence above, we obtain an isomorphism

$$\pi_0^v \pi_q^h Q^+ \xrightarrow{\sim} \pi_0^v \pi_q^h Z.$$

We will now determine these homotopy groups. Note that the functorial homomorphisms $G\bar{W}(H) \rightarrow H$ are homotopy equivalences for any simplicial group H . From this it follows that all the maps $d_i^v: Q_{p,q+1}^+ \rightarrow Q_{p,q}^+$ are homotopy equivalences too, since the functor C is applied dimension-wise. Hence,

$$\pi_p^v \pi_q^h Q^+ = 0 \quad \text{for } p > 0$$

and

$$\pi_0^v \pi_q^h Q^+ = \pi_q CG\bar{W}E((FR)^+) = \pi_{q+1} \mathbb{Z}_\infty \bar{W}E((FR)^+).$$

Performing the same calculation for Z and substituting this into the isomorphism above, we find the following proposition, which can be seen as a generalization of Gersten's theorem to include some types of free simplicial rings:

Proposition 2. *For each $q \geq 0$ we have*

$$\pi_q \mathbb{Z}_\infty \bar{W}E((FR)^+) \cong \pi_q \mathbb{Z}_\infty \bar{W}E(\mathbb{Z}).$$

Let FR once again be the cotriple resolution of R in \mathbf{Rg} . Then $F_0R \rightarrow R$ induces a surjective homomorphism $E(FR) \rightarrow E(R)$. Its kernel is denoted by HR . From the long exact sequence of the fibration $HR \rightarrow E(FR) \rightarrow E(R)$ it follows that $\pi_n HR = K'_{n+2}(R)$ for all $n > 0$ and $\pi_0 HR = \text{Ker}(\text{St}(R) \rightarrow E(R))$. Hence we have

$$\pi_n HR = K'_{n+2}(R) \quad \text{for all } n \geq 0.$$

Note that the simplicial group HR is also the kernel of the homomorphism $E((FR)^+) \rightarrow E(R^+)$. To see this, apply the snake lemma to the following diagram having split exact rows:

$$\begin{array}{ccccccc} & & HR & & & & \\ & & \downarrow & & & & \\ 1 & \longrightarrow & E(FR) & \longrightarrow & E(((FR)^+)) & \longrightarrow & E(\mathbb{Z}) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 1 & \longrightarrow & E(R) & \longrightarrow & E(R^+) & \longrightarrow & E(\mathbb{Z}) \longrightarrow 1 \end{array}$$

This gives the desired identification of the simplicial group HR with the kernel of $E((FR)^+) \rightarrow E(R^+)$.

The point of introducing this simplicial group HR is that the fibration

$$1 \rightarrow HR \rightarrow E((FR)^+) \rightarrow E(R^+) \rightarrow 1$$

has good behavior under application of the composite functor $\mathbb{Z}_\infty \bar{W}$. We need to verify the requirements of Property I3. Taking classifying complexes we obtain the following fibration in \mathbf{sSet} :

$$\bar{W}HR \rightarrow \bar{W}E((FR)^+) \rightarrow \bar{W}E(R^+).$$

Recall that for any simplicial group G the usual action of $\pi_1(\bar{W}G) = \pi_0(G)$ on $\pi_i(\bar{W}G)$ is the action induced by dimension-wise conjugation of G by degenerate elements originating from G_0 .

Lemma. *There is a natural isomorphism $GL(Z_i(FR)) \rightarrow Z_i(GL(FR))$ which is induced by the projection maps $Z_i(FR) \rightarrow F_iR$.*

Proof. The simplicial kernel $Z_i(FR)$ is an inverse limit of a suitable system of rings and the functor GL preserves inverse limits. By inspecting the pullback diagrams, it is clear that $Z_i(FR)$ is completely determined by the projection maps $p_*: Z_i(FR) \rightarrow F_iR$. We have a homomorphism $(GL(p_1), \dots, GL(p_{i+2})) : GL(Z_i(FR)) \rightarrow (GL(F_iR))^{i+2}$. Its image is precisely $Z_i(GL(FR))$. \square

Proposition 3. *The usual action of $\pi_1(\bar{W}E((FR)^+))$ on $\pi_i(\bar{W}E((FR)^+))$ is trivial for $i > 1$.*

Proof. For $i > 0$ we have the isomorphism

$$\pi_i(E(FR)^+) \rightarrow Z_i(E((FR)^+)/dE((FR)^+)_{i+1})$$

which commutes with the action of $E((FR)^+)$ (see Section 5.) Using this we have that

$$\begin{aligned} Z_i(E((FR)^+)) &\cong Z_i(E(FR) \rtimes E(\mathbb{Z})) = Z_i(GL(FR) \rtimes E(\mathbb{Z})) \\ &\cong Z_i(GL(FR)) \rtimes E(\mathbb{Z}) \cong GL(Z_i(FR) \rtimes E(\mathbb{Z})). \end{aligned}$$

The image of the group $dE((FR)^+)_{i+1}$ under this composition is the group $E(Z_i(FR)) \rtimes E(\mathbb{Z})$. To see this, note that $d: (FR)_{i+1}^+ \rightarrow Z_i((FR)^+)$ is onto and that the functor E preserves such maps.

The images of the elements of $E((F_0R)^+) \subseteq Z_i(E((FR)^+))$ are contained in $E(Z_i(FR)) \rtimes E(\mathbb{Z})$. Hence the action on $Z_i(E((FR)^+))$ corresponds to an action which becomes trivial when passing to quotients. It follows that the action of $E((FR)^+)$ on $\pi_{i+1}(\bar{W}E((FR)^+))$ is trivial. \square

Corollary. *The action of $\pi_1(\bar{W}E((FR)^+))$ on $\pi_i(\bar{W}HR)$ is trivial for all i .*

Proof. For $i > 1$ this is a consequence of the previous lemma. For $i = 1$ the action is also trivial because $\pi_1(\bar{W}HR)$ maps isomorphically onto the kernel of $\pi_1(\bar{W}E((FR)^+) \rightarrow E(R^+))$, which identifies with the central extension $\text{St}(R) \rtimes E(\mathbb{Z}) \rightarrow E(R) \rtimes E(\mathbb{Z})$. \square

Corollary. *The action of $\pi_1(\bar{W}HR)$ on $\pi_i(\bar{W}HR)$ is trivial.*

Proof. The homomorphism $\pi_1(\bar{W}HR) \rightarrow \pi_1(\bar{W}E((FR)^+))$ is injective by the long exact sequence of a fibration. Now use the previous corollary. \square

Proposition 4. *The induced map $\mathbb{Z}_\infty \bar{W}E((FR)^+) \rightarrow \mathbb{Z}_\infty \bar{W}E(R^+)$ is a fibration and its fibre is homotopy equivalent to $\bar{W}HR$.*

Proof. From Property I3 and the above corollary it follows that the map $\mathbb{Z}_\infty \bar{W}E((FR)^+) \rightarrow \mathbb{Z}_\infty \bar{W}E(R^+)$ is a fibration and that the canonical map from $\mathbb{Z}_\infty \bar{W}HR$ to the fibre is a homotopy equivalence. From Property I2 and this

corollary it follows that the natural map $i: \bar{W}HR \rightarrow \mathbb{Z}_\infty \bar{W}HR$ is a weak homotopy equivalence. Since all simplicial sets involved are Kan complexes, this map is a fortiori a homotopy equivalence. \square

Now we can finish the proof of the theorem.

Proof. Let X be the fibre of the map $\mathbb{Z}_\infty \bar{W}E(R^+) \rightarrow \mathbb{Z}_\infty \bar{W}E(\mathbb{Z})$. We have the following diagram which consists horizontally and vertically of long exact sequences of suitable fibrations:

$$\begin{array}{ccccccc}
 \pi_{p+1}(X) & \xrightarrow{\sim} & \pi_p(\cdots) & \longrightarrow & 0 & \longrightarrow & \pi_p(X) \\
 \downarrow & & \downarrow \wr & & \downarrow & & \downarrow \\
 \pi_{p+1}\mathbb{Z}_\infty \bar{W}E(R^+) & \longrightarrow & \pi_p \bar{W}HR & \longrightarrow & \pi_p \mathbb{Z}_\infty \bar{W}E((FR)^+) & \longrightarrow & \pi_p \mathbb{Z}_\infty \bar{W}E(R^+) \\
 \downarrow & & \downarrow & & \downarrow \wr & & \downarrow \\
 \pi_{p+1}\mathbb{Z}_\infty \bar{W}E(\mathbb{Z}) & \longrightarrow & 0 & \longrightarrow & \pi_p \mathbb{Z}_\infty \bar{W}E(\mathbb{Z}) & \xrightarrow{=} & \pi_p \mathbb{Z}_\infty \bar{W}E(\mathbb{Z})
 \end{array}$$

The map in the third column is an isomorphism by Proposition 2. The other relations are evident from the diagram. The group $K_{p+1}(R^+, R)$ equals $\pi_{p+1}(X)$ for $p \geq 1$, which is in turn isomorphic to $\pi_p(\bar{W}H(R))$ by the above diagram. The latter group equals $K'_{p+1}(R)$ for $p \geq 1$. \square

Final remark. An alternative way to prove the equivalence of both algebraic K -theories is by showing that the Quillen K -theory satisfies the axioms for multirelative K -theory given in [12]. To do so, the Quillen K -theory has to be extended to include multirelative groups. The main concern is then to extend long exact sequences in such a way that they include K_0 -groups as well.

REFERENCES

1. D. W. ANDERSON, Relationship among K -theories. *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, 57–72. *Lecture Notes in Math.* 341, Springer, Berlin, 1973.
2. A. J. BERRICK, An approach to algebraic K -theory. *Research Notes in Mathematics*, 56. Pitman, Boston, 1972.
3. A. K. BOUSFIELD and D. M. KAN, Homotopy limits, completions and localizations. *Lecture Notes in Mathematics*. 04. Springer-Verlag, Berlin, 1972.
4. S. M. GERSTEN, On the functor K_2 . *J. Algebra* **17**(1971), 212–237.
5. S. M. GERSTEN, K -theory of free rings. *Comm. Algebra* **1**(1974), 39–64.
6. S. M. GERSTEN, Higher K -theory of rings. *Algebraic K-theory, I: Higher K-theories (Proc. Conf. Seattle Res. Center, Battelle Memorial Inst., 1972)*, 3–42. *Lecture Notes in Math.*, 341. Springer, Berlin, 1973.
7. H. INASSARIDZE, Algebraic K -theory. *Mathematics and its Applications*, 311. Kluwer Academic Publishers Group, Dordrecht, 1995.
8. H. INASSARIDZE, Non-abelian homological algebra and its applications. *Mathematics and Its Applications*, 421. Kluwer Academic Publishers, Dordrecht, 1997.

9. F. KEUNE, Nonabelian derived functors and algebraic K -theory. *Algebraic K-theory, I: Higher K-theories (Proc. Conf. Seattle Res. Center, Battelle Memorial Inst., 1972)*, 166–176. *Lecture Notes in Math.*, 341, Springer, Berlin, 1973.
10. F. KEUNE, The relativization of K_2 . *J. Algebra* **54**(1978), 159–177.
11. F. KEUNE, On the equivalence of two higher algebraic K -theories. *Report 7913, July 1979, Mathematisch Instituut K.U. Nijmegen*.
12. F. KEUNE, Multirelative K -theory and axioms for the K -theory of rings. *Doc. Math.* **1**(1996), No. 14, 277–291 (electronic).
13. J. -L. LODAY, K -théorie algébrique et représentations de groupes. *Ann. Sci. École Norm. Sup. (4)* **9**(1976), No. 3, 309–377.
14. J. P. MAY, Simplicial objects in algebraic topology. *Van Nostrand Mathematical Studies*, No. 11. *D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London*, 1967.
15. D. QUILLEN, Spectral sequences of a double simplicial group. *Topology* **5**(1966), 155–158.
16. R. G. SWAN, Nonabelian homological algebra and K -theory. *Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968)*, 88–123. *Amer. Math. Soc., Providence, R.I.*, 1970.
17. R. G. SWAN, Some relations between higher K -functors. *J. Algebra* **21**(1972), 113–136.

(Received 30.04.2002)

Authors' address:

Department of Mathematics

Katholieke Universiteit Nijmegen

Postbus 9010, 6500 GL Nijmegen

The Netherlands

E-mail: ebeling@sci.kun.nl

keune@sci.kun.nl