

Research Article

Subharmonic Solutions of Nonautonomous Second Order Differential Equations with Singular Nonlinearities

N. Daoudi-Merzagui,¹ F. Derrab,² and A. Boucherif³

¹ *Department of Mathematics, University of Tlemcen, Tlemcen 13000, Algeria*

² *Department of Mathematics, University of Sidi Bel Abbes, Sidi Bel Abbes 22000, Algeria*

³ *Department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, P.O. Box 5046, Dhahran 31261, Saudi Arabia*

Correspondence should be addressed to A. Boucherif, aboucher@kfupm.edu.sa

Received 4 November 2011; Revised 14 January 2012; Accepted 19 January 2012

Academic Editor: Juan J. Nieto

Copyright © 2012 N. Daoudi-Merzagui et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We discuss the existence of subharmonic solutions for nonautonomous second order differential equations with singular nonlinearities. Simple sufficient conditions are provided enable us to obtain infinitely many distinct subharmonic solutions. Our approach is based on a variational method, in particular the saddle point theorem.

1. Introduction and Main Result

In this paper we discuss the problem of the existence of infinitely many subharmonic solutions for nonautonomous second order differential equations with singular nonlinearities of the form

$$u''(t) + f(t, u(t)) = e(t), \quad (1.1)$$

where $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is continuous, is T -periodic, in its first argument with $T > 0$, and presents a singularity with respect to its second argument. Here by a subharmonic solution we mean a kT -periodic solution for any integer k if $T > 0$ is the minimal period. When the solution is not T -periodic we call it a true subharmonic. It was pointed out in [1] that singular differential equations of the form (1.1) appear in the description of many phenomena in the applied sciences, such as the Brillouin focusing system and nonlinear elasticity. Several authors have

investigated the problem of existence of periodic solutions for second order differential equations with singular nonlinearities (see [2–4] and the references therein). Topological and variational methods are the two main techniques that have been developed for the study of (1.1). We refer the interested reader to the paper [1] for details and references on the topological methods. In this work we shall rely on the saddle point theorem, see [5, 6], to prove our main result. We use the truncation techniques introduced in [7] to modify our problem to one without singularities. We assume that the nonlinearity f is monotone with respect to its first variable t . When f is increasing, our result generalizes the result in [8]. We can obtain the same result by considering the monotonicity of the potential function instead of the field f . For more results on the subject and different techniques one can consult the papers [9–12]. We should point out some related recent articles, for instance [13, 14].

Throughout this paper we shall use the following notations. Let $I = [0, T]$. $L^p(I)$ is the classical Lebesgue space of functions $u : I \rightarrow \mathbb{R}$ such that $|u(\cdot)|^p$ is integrable, and for $u \in L^p(I)$ we define its norm by

$$\|u\|_{L^p} = \left(\int_0^T |u(t)|^p dt \right)^{1/p}. \quad (1.2)$$

Let $\|u\|_\infty = \sup\{|u(t)|; t \in [0, kT]\}$. For $T > 0$ and $k \in \mathbb{N}$ we let $H_{kT}^1 = \{u \in W^{1,2}([0, kT], \mathbb{R}); u(0) = u(kT)\}$ and for $u \in H_{kT}^1$ we define its norm by

$$\|u\|_{H_{kT}^1} = (\|u\|_{L^2} + \|u'\|_{L^2})^{1/2}. \quad (1.3)$$

H_{kT}^1 endowed with the norm $\|\cdot\|_{H_{kT}^1}$ is a reflexive Banach space. Also $H_{kT}^1 = H^+ \oplus H^-$, orthogonal decomposition, where H^+ is the subspace of constant functions in H_{kT}^1 and H^- denotes the subspace of functions in H_{kT}^1 with mean value zero; so that $u \in H_{kT}^1$ can be written as $u = \bar{u} + \tilde{u}$ with $\bar{u} \in H^+$ and $\tilde{u} \in H^-$.

We shall assume that $e : \mathbb{R} \rightarrow \mathbb{R}$ is a locally integrable T -periodic function. We denote the mean value of e by \bar{e} , that is, $\bar{e} = 1/T \int_0^T e(t) dt$. It follows that \bar{e} , $\|e\|_{L^1}$ and $\|e\|_{L^2}$ are bounded. Moreover, since e is T -periodic, we have $\int_0^{kT} e(t) dt = \sum_{j=0}^{k-1} \int_0^T e(t + jT) dt = \sum_{j=0}^{k-1} \int_0^T e(t) dt = k \int_0^T e(t) dt = kT\bar{e}$.

Let $F(t, u) = \int_1^u f(t, s) ds$ be an antiderivative of f defined for all $u \in \mathbb{R}$ and for all $t \in I$. We introduce the following assumptions on the nonlinearity.

(H1) $f : \mathbb{R} \times (0, +\infty) \rightarrow \mathbb{R}$ is continuous, $f(t+T, u) = f(t, u)$ for all $(t, u) \in \mathbb{R} \times (0, +\infty)$, and such that

- (i) $t \rightarrow f(t, u)$ is monotone for each fixed u in $(0, +\infty)$,
- (ii) $\lim_{u \rightarrow 0^+} f(t, u) = -\infty$, uniformly in $t \in I$,
- (iii) $f(0, u) = f(T, u) > \bar{e}$ for all $u > 1$.

(H2)

- (i) $\lim_{u \rightarrow +\infty} 2F(t, u)/u^2 = 0$, uniformly in $t \in I$.
- (ii) $\lim_{u \rightarrow +\infty} \int_0^T [F(t, u) - \bar{e}u] dt = +\infty$.

Theorem 1.1. *Assume that e is a locally integrable T -periodic function. If (H1) and (H2) are satisfied then (1.1) has a sequence $(u_k)_{k \geq 1}$ of kT -periodic solutions whose amplitudes and minimal periods tend to infinity. In particular, if T is the minimal period of e and of f with respect to t , (1.1) admits solutions with minimal periods kT for every sufficiently Large Integer k .*

2. Proof of Theorem 1.1

The proof of this result will be based on several auxiliary results.

2.1. Modification of the Problem

Define the truncation function $f_r : \mathbb{R}^2 \rightarrow \mathbb{R}, 0 < r \leq 1$, by

$$f_r(t, u) = \begin{cases} f(t, u), & u \geq r, \\ f(t, r), & u < r. \end{cases} \quad (2.1)$$

Note that condition (H1) implies that f_r is continuous with respect to $(t, u) \in I \times \mathbb{R}$ and T -periodic with respect to its first variable t .

Lemma 2.1. *Assume (H1) and (H2) (ii) are satisfied. Then there exists $d > 1$ such that for every $u \in (0, 1/d) \cup (d, +\infty)$*

$$(f(t, u) - \bar{e})(u - 1) > 0, \quad \text{uniformly in } t \in I. \quad (2.2)$$

Proof. First, it follows from (H1)(ii) that for any $A > 0$, there is $\delta_A > 0$ such that for every $u \in (0, \delta_A)$ we have $f(t, u) < -A$, uniformly in $t \in I$.

In particular for $A > 2|\bar{e}| + 1$, there is $\delta_A > 0$ such that for every $u \in (0, \delta_A)$ it holds

$$f(t, u) - \bar{e} < -|\bar{e}| - 1 < 0. \quad (2.3)$$

Choose $d_1 > 1$ such that $1/d_1 < \delta_A$. Then for every $u \in (0, 1/d_1)$, we have $f(t, u) - \bar{e} < 0$.

Therefore, condition (H1)(ii) implies that there exists $d_1 > 1$ such that for every $u \in (0, 1/d_1)$ it holds $(f(t, u) - \bar{e})(u - 1) > 0$, uniformly in $t \in I$.

Next, condition (H2)(ii) implies that for any $B > 1$, sufficiently large, there is $\chi_B > 0$, large enough, such that for every $u > \chi_B$ we have

$$\int_0^T [F(t, u) - \bar{e}u] dt > B. \quad (2.4)$$

Hence, there exists d_2 , sufficiently large such that $d_2 > \max(1, \chi_B)$ and for every $u > d_2$ it holds

$$\int_0^T [F(t, u) - \bar{e}u] dt > B. \quad (2.5)$$

We show that for $u > d_2$ we have $f(t, u) - \bar{e} > 0$, uniformly in $t \in I$. Assume, on the contrary, that there exists $y > d_2$ for which $f(t_0, y) - \bar{e} \leq 0$, for some $t_0 \in (0, T)$. By the continuity of

f on I and ((H1)(iii)), there exists $\sigma \in (0, T)$ such that $f(\sigma, u) = \max_{t \in I} f(t, u)$. If $t_0 \leq \sigma$ then $0 < f(0, y) - \bar{e} < f(t_0, y) - \bar{e} \leq 0$, and this is a contradiction. Similarly, if $t_0 \geq \sigma$ then $0 < f(0, y) - \bar{e} = f(T, y) - \bar{e} < f(t_0, y) - \bar{e} \leq 0$, and we again arrive at a contradiction.

Hence we deduce that if (H1) and (H2)(ii) hold, then there exists $d > \max(d_1, d_2)$ such that for any $u \in (0, 1/d) \cup (d, +\infty)$ $(f(t, u) - \bar{e})(u - 1) > 0$, uniformly in $t \in I$. This completes the proof of Lemma 2.1. \square

Lemma 2.2. *For every positive integer k , there exist r_k and R_k with $0 < r_k < 1/d < d < R_k$ such that for any $\mu \in (0, r_k]$ each kT -periodic solution u of*

$$u''(t) + f_\mu(t, u(t)) = e(t), \quad (2.6)$$

satisfies

$$r_k \leq u(t) \leq R_k, \quad \forall t \in \mathbb{R}. \quad (2.7)$$

In particular, any kT -periodic solution of (2.6), with $\mu = r_k$ is a solution of (1.1).

Proof. This is essentially Proposition 2.1 in [8]. We shall use some ideas from [8] (see also [15]). Fix $k \in \mathbb{N}$, and suppose, on the contrary, that for each integer n , there exist $\mu_n \in (0, 1/n)$ and a kT -periodic solution u_n satisfying

$$u_n''(t) + f_{\mu_n}(t, u_n(t)) = e(t) \quad (2.8)$$

and $\{u_n(t); t \in \mathbb{R}\} \not\subseteq [1/n, n]$.

Claim 1. Let d be as in Lemma 2.1 and let u_n be as above. Then for every n there exists $\tau_n \in [0, kT]$ such that $u_n(\tau_n) \in [1/d, d]$.

Indeed, it follows from (2.8) that

$$\int_0^{kT} f_{\mu_n}(t, u_n(t)) dt = kT\bar{e}. \quad (2.9)$$

Now, if $u_n(t) > d$ for all $t \in [0, kT]$, then Lemma 2.1 implies that $f_{\mu_n}(t, u_n(t)) - \bar{e} > 0$, which in turn yields $\int_0^{kT} f_{\mu_n}(t, u_n(t)) dt > kT\bar{e}$.

This contradicts (2.9). On the other hand, if $u_n(t) < 1/d$ for all $t \in [0, kT]$, then $f_{\mu_n}(t, u_n(t)) - \bar{e} < 0$, so that $\int_0^{kT} f_{\mu_n}(t, u_n(t)) dt < kT\bar{e}$. This is again a contradiction to (2.9).

Claim 2. There exists $R > 0$ such that $M_n = \max_{t \in [0, kT]} u_n(t) \leq R$ for each integer n . To prove the claim notice, there exists $t_n^1 \in [0, kT]$ such that $u_n(t_n^1) = M_n$. If $u_n(t_n^1) = M_n \in [1/d, d]$ then $u_n(t) \leq R$ for any $R > d$. So, assume that there exists a subsequence of $(u_n)_n$, which we label

the same, for which $M_n \rightarrow +\infty$ when $n \rightarrow +\infty$. So that $M_n > d$ for n large enough. Since $u_n(\tau_n) < d$, there exists an interval $[\alpha_n, \beta_n]$, containing t_n^1 , such that

$$\begin{aligned}\beta_n - \alpha_n &\leq kT, \\ u_n(\alpha_n) &= d = u_n(\beta_n), \\ d \leq u_n(t) &\leq u_n(t_n^1), \text{ for all } t \in [\alpha_n, \beta_n].\end{aligned}\tag{2.10}$$

Equation (2.8) can be written as

$$\begin{aligned}u_n'(t) &= v_n(t) + \int_{\alpha_n}^t [e(s) - \bar{e}] ds, \\ v_n'(t) &= -f_{\mu_n}(t, u_n(t)) + \bar{e}.\end{aligned}\tag{2.11}$$

Since for all $t \in (\alpha_n, \beta_n)$, $u_n(t) > d$ and $\mu_n < 1/n$, then the second equation in (2.11) is equivalent to

$$v_n'(t) = -f(t, u_n(t)) + \bar{e}.\tag{2.12}$$

Lemma 2.1 implies that $f(t, u_n(t)) - \bar{e} > 0$ for all $t \in [\alpha_n, \beta_n]$. Then $v_n'(t) < 0$ for all $t \in [\alpha_n, \beta_n]$ and hence v_n is decreasing on $[\alpha_n, \beta_n]$. The first equation in (2.11) implies

$$u_n'(t) \leq v_n(\alpha_n) + \int_{\alpha_n}^t [e(s) - \bar{e}] ds \quad \forall t \in [\alpha_n, \beta_n].\tag{2.13}$$

This yields

$$u_n'(t) \leq v_n(\alpha_n) + \|e\|_{L^1} + kT|\bar{e}| \quad \forall t \in [\alpha_n, \beta_n].\tag{2.14}$$

Integrating the above inequality over $[\alpha_n, t_n^1] \subset [\alpha_n, \beta_n]$ we obtain

$$u_n(t_n^1) - u_n(\alpha_n) = M_n - d \leq kT[v_n(\alpha_n) + \|e\|_{L^1} + kT|\bar{e}|].\tag{2.15}$$

Equation (2.15) leads to

$$\lim_{n \rightarrow +\infty} \left(1 - \frac{d}{M_n}\right) \leq kT \lim_{n \rightarrow +\infty} \left(\frac{v_n(\alpha_n) + \|e\|_{L^1} + kT|\bar{e}|}{M_n}\right).\tag{2.16}$$

Hence

$$1 \leq kT \lim_{n \rightarrow +\infty} \frac{v_n(\alpha_n)}{M_n}.\tag{2.17}$$

It is clear from (2.17) that $v_n(\alpha_n) \rightarrow +\infty$ when $n \rightarrow +\infty$. So that, for n large enough, we have

$$v_n(\alpha_n) \geq \|e\|_{L^1} + kT|\bar{e}|.\tag{2.18}$$

Since $M_n = u_n(t_n^1)$, we have $u_n'(t_n^1) = 0$. It follows from the first equation in (2.11) that

$$\left| v_n(t_n^1) \right| = \left| \int_{\alpha_n}^{t_n^1} [e(s) - \bar{e}] ds \right| \leq \|e\|_{L^1} + kT|\bar{e}|. \quad (2.19)$$

We see from (2.18) and (2.19) that for n large enough

$$v_n(\alpha_n) \geq \|e\|_{L^1} + kT|\bar{e}| \geq v_n(t_n^1). \quad (2.20)$$

Since $v_n(\cdot)$ is continuous on $[\alpha_n, \beta_n]$, then, for n large enough, there exists at least one $t_n^* \in (\alpha_n, t_n^1)$ such that $v_n(t_n^*) = \|e\|_{L^1} + kT|\bar{e}|$. We denote by t_n^2 , the first such t_n^* . Then

$$v_n(t_n^2) = \|e\|_{L^1} + kT|\bar{e}|. \quad (2.21)$$

We distinguish two cases.

Case 1. $f(\cdot, s)$ is increasing for each fixed $s \in (0, +\infty)$.

Consider the function B_1 defined by

$$B_1(t) = F(t_n^2, u_n(t)) - \bar{e}u_n(t) + \frac{1}{2}(v_n(t) - \|e\|_{L^1} - kT|\bar{e}|)^2. \quad (2.22)$$

Then

$$B_1'(t) = \left(f(t_n^2, u_n(t)) - \bar{e} \right) u_n'(t) + (v_n(t) - \|e\|_{L^1} - kT|\bar{e}|) v_n'(t). \quad (2.23)$$

Since $v_n'(t) = -f(t, u_n(t)) + \bar{e}$, it follows from the first equation in (2.11)

$$\begin{aligned} B_1'(t) &= \left(f(t_n^2, u_n(t)) - \bar{e} \right) \left(v_n(t) + \int_{\alpha_n}^t [e(s) - \bar{e}] ds \right) \\ &\quad + (v_n(t) - \|e\|_{L^1} - kT|\bar{e}|) (-f(t, u_n(t)) + \bar{e}) \\ &= \left(f(t_n^2, u_n(t)) - \bar{e} \right) \left(v_n(t) + \int_{\alpha_n}^t [e(s) - \bar{e}] ds + \|e\|_{L^1} + kT|\bar{e}| - \|e\|_{L^1} - kT|\bar{e}| \right) \\ &\quad + (v_n(t) - \|e\|_{L^1} - kT|\bar{e}|) (-f(t, u_n(t)) + \bar{e}). \end{aligned} \quad (2.24)$$

Hence

$$\begin{aligned} B_1'(t) &= \left(f(t_n^2, u_n(t)) - \bar{e} \right) \left(\int_{\alpha_n}^t [e(s) - \bar{e}] ds + \|e\|_{L^1} + kT|\bar{e}| \right) \\ &\quad + \left(f(t_n^2, u_n(t)) - f(t, u_n(t)) \right) (v_n(t) - \|e\|_{L^1} - kT|\bar{e}|). \end{aligned} \quad (2.25)$$

Since, for all $t \in [\alpha_n, \beta_n]$

$$\left| \int_{\alpha_n}^t [e(s) - \bar{e}] ds \right| \leq \|e\|_{L^1} + kT|\bar{e}|, \quad (2.26)$$

it follows that for all $t \in [\alpha_n, \beta_n]$

$$\int_{\alpha_n}^t [e(s) - \bar{e}] ds + \|e\|_{L^1} + kT|\bar{e}| \geq 0. \quad (2.27)$$

Also, Lemma 2.1 implies that $f(t_n^2, u_n(t)) - \bar{e} > 0$ for all $t \in [\alpha_n, \beta_n]$. Furthermore, the monotonicity of f implies that $f(t_n^2, u_n(t)) - f(t, u_n(t)) \geq 0$ for all $t \in [\alpha_n, t_n^2]$. Since $v_n(\cdot)$ is decreasing on $[\alpha_n, t_n^2]$ and $v_n(t_n^2) = \|e\|_{L^1} + kT|\bar{e}|$, it follows that

$$v_n(t) - \|e\|_{L^1} - kT|\bar{e}| \geq v_n(t_n^2) - \|e\|_{L^1} - kT|\bar{e}| = 0 \quad (2.28)$$

for all $t \in [\alpha_n, t_n^2]$. Now, Lemma 2.1 combined with (2.27), (2.28), and the monotonicity of f with respect to its first variable shows that

$$B_1'(t) \geq 0, \quad \forall t \in [\alpha_n, t_n^2]. \quad (2.29)$$

Thus, the function B_1 is increasing on $[\alpha_n, t_n^2]$. Since $u_n(\alpha_n) = d$,

$$\begin{aligned} B_1(\alpha_n) &= F(t_n^2, d) - \bar{e}d + \frac{1}{2}(v_n(\alpha_n) - \|e\|_{L^1} - kT|\bar{e}|)^2 \\ &\leq B_1(t_n^2) = F(t_n^2, u_n(t_n^2)) - \bar{e}u_n(t_n^2) + \frac{1}{2}(v_n(t_n^2) - \|e\|_{L^1} - kT|\bar{e}|)^2. \end{aligned} \quad (2.30)$$

Since $v_n(t_n^2) - \|e\|_{L^1} - kT|\bar{e}| = 0$, it follows that

$$B_1(\alpha_n) = F(t_n^2, d) - \bar{e}d \leq F(t_n^2, u_n(t_n^2)) - \bar{e}u_n(t_n^2). \quad (2.31)$$

Notice that

$$F(t_n^2, u_n(t_n^2)) - \bar{e}u_n(t_n^2) = \int_1^d (f(t_n^2, s) - \bar{e}) ds + \int_d^{u_n(t_n^2)} (f(t_n^2, s) - \bar{e}) ds - \bar{e}. \quad (2.32)$$

Also, $f(t_n^2, s) - \bar{e} > 0$ if $s \in (d, u_n(t_n^2)) \subset [d, u_n(t_n^1)]$ (see Lemma 2.1). It follows that

$$\int_d^{u_n(t_n^2)} [f(t_n^2, s) - \bar{e}] ds \leq \int_d^{u_n(t_n^1)} [f(t_n^2, s) - \bar{e}] ds. \quad (2.33)$$

Hence

$$B_1(\alpha_n) \leq \int_1^d [f(t_n^2, s) - \bar{e}] ds + \int_d^{u_n(t_n^1)} [f(t_n^2, s) - \bar{e}] ds - \bar{e} = F(t_n^2, u_n(t_n^1)) - \bar{e}u_n(t_n^1). \quad (2.34)$$

Set

$$\rho_n = \frac{1}{2} \left(\frac{v_n(\alpha_n) - \|e\|_{L^1} - kT|\bar{e}|}{M_n} \right)^2. \quad (2.35)$$

Then

$$\lim_{n \rightarrow +\infty} \rho_n = 0. \quad (2.36)$$

Indeed, since $M_n = u_n(t_n^1)$, we have

$$B_1(\alpha_n) = F(t_n^2, d) - \bar{e}d + \rho_n M_n^2 \leq F(t_n^2, u_n(t_n^1)) - \bar{e}u_n(t_n^1) = F(t_n^2, M_n) - \bar{e}M_n. \quad (2.37)$$

From (2.37) we deduce

$$\lim_{n \rightarrow +\infty} \frac{F(t_n^2, d)}{(M_n)^2} + \lim_{n \rightarrow +\infty} \rho_n \leq \lim_{n \rightarrow +\infty} \frac{F(t_n^2, M_n)}{(M_n)^2}. \quad (2.38)$$

Notice that $F(t_n^2, d) = \int_1^d f(t_n^2, s) ds$ is bounded, so that

$$\lim_{n \rightarrow +\infty} \frac{F(t_n^2, d)}{(M_n)^2} = 0. \quad (2.39)$$

Also, (H2)(i) implies that

$$\lim_{n \rightarrow +\infty} \frac{F(t_n^2, M_n)}{(M_n)^2} = 0. \quad (2.40)$$

Therefore (2.36) holds; that is, $\lim_{n \rightarrow +\infty} \rho_n = 0$. Since $\|e\|_{L^1}$ and $|\bar{e}|$ are bounded, it follows that

$$\lim_{n \rightarrow +\infty} \frac{v_n(\alpha_n)}{M_n} = 0. \quad (2.41)$$

It is clear that (2.41) contradicts (2.17).

Case 2. $f(\cdot, s)$ is decreasing for each fixed $s \in (0, +\infty)$.

In this case we consider the function B_2 defined by

$$B_2(t) = F(\alpha_n, u_n(t)) - \bar{e}u_n(t) + \frac{1}{2}(v_n(t) - \|e\|_{L^1} - kT|\bar{e}|)^2. \quad (2.42)$$

Then

$$\begin{aligned}
 B_2'(t) &= (f(\alpha_n, u_n(t)) - \bar{e}) \left(\int_{\alpha_n}^t [e(s) - \bar{e}] ds + \|e\|_{L^1} + kT|\bar{e}| \right) \\
 &\quad + (f(\alpha_n, u_n(t)) - f(t, u_n(t)))(v_n(t) - \|e\|_{L^1} - kT|\bar{e}|) \\
 &\geq 0 \quad \forall t \in [\alpha_n, t_n^2].
 \end{aligned} \tag{2.43}$$

Repeating the same reasoning as in Case 1, we arrive at a contradiction.

Therefore, we deduce that there must exist $R > 0$ such that $M_n = \max_{t \in [0, kT]} u_n(t) \leq R$ for each n .

Next, we prove that there is $r > 0$ such that $u_n(t) \geq r$ for every $t \in [0, kT]$. Assuming that this is not true we will obtain a contradiction.

Consider the following sets

$$\begin{aligned}
 I_{1/n} &= \left\{ t \in [0, kT]; u_n(t) < \frac{1}{n} \right\}, \\
 I_{1/n, 1/d} &= \left\{ t \in [0, kT]; \frac{1}{n} \leq u_n(t) < \frac{1}{d} \right\}, \\
 I_{1/d, R} &= \left\{ t \in [0, kT]; \frac{1}{d} \leq u_n(t) \leq R \right\}.
 \end{aligned} \tag{2.44}$$

It is clear that for $n \geq R$, $u_n(t) \leq R \leq n$. Also, we cannot have $u_n(t) \geq 1/n$ for every $t \in [0, kT]$, for otherwise we would have

$$\frac{1}{n} \leq u_n(t) \leq n \quad \forall t \in [0, kT], \tag{2.45}$$

which contradicts the assumption $\{u_n(t); t \in \mathbb{R}\} \notin [(1/n), n]$. Hence, for $n \geq R$ there exists $t_n^3 \in [0, kT]$ such that $u_n(t_n^3) < 1/n$. This shows that $I_{1/n} \neq \emptyset$. The continuity of u_n implies that $I_{1/n}$ is open, and so $meas(I_{1/n}) \neq 0$.

Define

$$\Psi_n = \int_0^{kT} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt, \quad n \in \mathbb{N}. \tag{2.46}$$

It follows from (2.9) that $\Psi_n = 0$. On the other hand

$$\Psi_n = \int_{I_{1/n}} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt + \int_{I_{1/n, 1/d}} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt + \int_{I_{1/d, R}} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt. \tag{2.47}$$

(i) Assume we are integrating positively on all subintervals of $[0, kT]$.

If $t \in I_{1/n, 1/d}$, then $u_n(t) \in [1/n, 1/d) \subset (0, 1/d)$. So that, by Lemma 2.1,

$$\int_{I_{1/n, 1/d}} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt < 0. \tag{2.48}$$

For $t \in I_{1/d, R}$ we have $u_n(t) \in [1/d, R]$. This means that $u_n(t)$ is bounded uniformly in $t \in I_{1/d, R}$. Since f_{μ_n} is continuous it is bounded on $I_{1/d, R}$.

Let

$$c = \max \left\{ |f_{\mu_n}(t, x)|; t \in [0, kT], \frac{1}{d} \leq x \leq R \right\} = \max_{I_{1/d, R}} \{ |f_{\mu_n}(t, x)| \}. \quad (2.49)$$

Then

$$\left| \int_{I_{1/d, R}} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt \right| \leq \int_{I_{1/d, R}} [|f_{\mu_n}(t, u_n(t))| + |\bar{e}|] dt \leq kT(c + |\bar{e}|). \quad (2.50)$$

It follows from (2.47), (2.48), and (2.50) that

$$\Psi_n < \int_{I_{1/n}} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt + kT(c + |\bar{e}|). \quad (2.51)$$

Claim 3.

$$\lim_{n \rightarrow \infty} \int_{I_{1/n}} [f_{\mu_n}(t, u_n(t)) - \bar{e}] dt = -\infty. \quad (2.52)$$

Proof. Recall that $\mu_n \in (0, 1/n)$ and $u_n(t) < 1/n$ for each $t \in I_{1/n}$. Then, if $u_n(t) < \mu_n$ we have $f_{\mu_n}(t, u_n(t)) = f(t, \mu_n)$, and if $u_n(t) \in [\mu_n, 1/n]$, we have $f_{\mu_n}(t, u_n(t)) = f(t, u_n(t))$. In both cases condition (H1)(ii) and the continuity of f imply that $\lim_{n \rightarrow \infty} (f_{\mu_n}(t, u_n(t)) - \bar{e}) = -\infty$ for every $t \in I_{1/n}$.

Since \bar{e} is bounded, then (2.51) implies that

$$\lim_{n \rightarrow +\infty} \Psi_n = -\infty, \quad (2.53)$$

which is a contradiction with (2.9).

(ii) If we integrate negatively on all subintervals of $[0, kT]$ we will obtain $\lim_{n \rightarrow +\infty} \Psi_n = +\infty$, which, again, contradicts (2.9). Thus, the proof of Lemma 2.2 is complete. \square

Remark 2.3. Lemma 2.2 shows that any kT -periodic solution u of (2.6), with $\mu = r_k$ is a solution of (1.1), since it satisfies $u(t) \geq r_k$ for all $t \in \mathbb{R}$ and $f_{r_k}(t, u(t)) = f(t, u(t))$.

In the remainder of the paper we shall deal with (2.6), with $\mu = r_k$ instead of (1.1). Let $F_{r_k}(t, u) = \int_1^u f_{r_k}(t, s) ds$ be a primitive of f_{r_k} defined for all $t \in I$ and $u \in \mathbb{R}$.

Lemma 2.4. *If (H1) and (H2) hold, then f_{r_k} and F_{r_k} satisfy the following conditions.*

- (L1) f_{r_k} is defined and continuous in $(t, u) \in I \times \mathbb{R}$ and T -periodic with respect to $t \in I$.
- (L2) $\liminf_{|u| \rightarrow +\infty} 2F_{r_k}(t, u)/u^2 = 0$, uniformly in $t \in I$.
- (L3) $\exists d > 1$ such that for $u \in (-\infty, 1/d) \cup (d, +\infty)$ it holds $(f_{r_k}(t, u) - \bar{e})(u - 1) > 0$ uniformly in $t \in I$.
- (L4) $\lim_{|u| \rightarrow +\infty} \int_0^T [F_{r_k}(t, u) - \bar{e}u] dt = +\infty$.

2.2. Existence of kT -Periodic Solutions for Equation (2.6), $\mu = r_k$

Using a variational method we shall show that equation (2.6), with $\mu = r_k$ has infinitely many kT -periodic solutions. In fact, we have the following.

Lemma 2.5. *Assume that e is locally integrable T -periodic function and the conditions (L1), (L2), (L3), (L4) hold. Then (2.6), with $\mu = r_k$ admits a kT -periodic solution u_k .*

Proof. We shall rely on a variational method and more precisely on the saddle point theorem. Define for each $k \geq 1$ the action functional $J_k : H_{kT}^1 \rightarrow \mathbb{R}$ by

$$J_k(u) = \int_0^{kT} \left[\frac{1}{2} [u'(t)]^2 - F_{r_k}(t, u(t)) + e(t)u(t) \right] dt. \tag{2.54}$$

J_k is well defined on H_{kT}^1 , weakly lower semicontinuous and continuously differentiable on H_{kT}^1 . Furthermore,

$$\langle J'_k(u), v \rangle = \int_0^{kT} [u'(t)v'(t) - f_{r_k}(t, u(t))v(t) + e(t)v(t)] dt, \quad \forall u, v \in H_{kT}^1. \tag{2.55}$$

The critical points of J_k are precisely the weak solutions of equation (2.6), with $\mu = r_k$.

First, we show that the functional J_k satisfies the Palais-Smale condition.

For this, let $k \geq 1$ be fixed and let $(u_n)_{n \in \mathbb{N}}$ be a sequence in H_{kT}^1 such that $(J_k(u_n))_{n \in \mathbb{N}}$ is bounded and $\lim_{n \rightarrow +\infty} J'_k(u_n) = 0$. Then $(u_n)_{n \in \mathbb{N}}$ has a convergent subsequence.

Suppose, on the contrary, that $\lim_{n \rightarrow +\infty} \|u_n\|_{H_{kT}^1} = +\infty$. Condition (L2) implies that for any $\varepsilon > 0$, small enough, there is $C_\varepsilon \geq 0$ such that

$$F_{r_k}(t, u) \leq \varepsilon u^2 + C_\varepsilon, \quad \forall u \in \mathbb{R}. \tag{2.56}$$

Writing $u_n(t) = \overline{u}_n + \widetilde{u}_n(t)$ for $t \in [0, kT]$, we obtain

$$-\int_0^{kT} F_{r_k}(t, u_n(t)) dt \geq -\varepsilon kT |\overline{u}_n|^2 - \varepsilon \int_0^{kT} |\widetilde{u}_n(t)|^2 dt - 2\varepsilon \overline{u}_n \int_0^{kT} \widetilde{u}_n(t) dt - C_\varepsilon kT. \tag{2.57}$$

Since $\widetilde{u}_n \in H^-$, we have $\int_0^{kT} \widetilde{u}_n(t) dt = kT \overline{\widetilde{u}_n} = 0$, so that

$$-\int_0^{kT} F_{r_k}(t, u_n(t)) dt \geq -\varepsilon \|\widetilde{u}_n\|_{L^2}^2 - \varepsilon kT |\overline{u}_n|^2 - C_\varepsilon kT. \tag{2.58}$$

Now, Hölder's inequality gives

$$\int_0^{kT} e(t)u_n(t) dt = \overline{u}_n \int_0^{kT} e(t) dt + \int_0^{kT} e(t)\widetilde{u}_n(t) dt \geq -|\overline{u}_n|kT|\overline{e}| - \|e\|_{L^2} \|\widetilde{u}_n\|_{L^2}. \tag{2.59}$$

Since $u'_n(t) = \widetilde{u}'_n(t)$ for $t \in [0, kT]$, it follows from (2.54), (2.58), and (2.59)

$$J_k(u_n) \geq \frac{1}{2} \|\widetilde{u}'_n\|_{L^2}^2 - \varepsilon \|\widetilde{u}_n\|_{L^2}^2 - \varepsilon kT |\overline{u}_n|^2 - C_\varepsilon kT - kT \left[\bar{e} |\overline{u}_n| - \|e\|_{L^2} \|\widetilde{u}_n\|_{L^2} \right]. \quad (2.60)$$

Wirtinger's inequality

$$\|\widetilde{u}'_n\|_{L^2}^2 \geq \left(\frac{4\pi^2}{4\pi^2 + (kT)^2} \right) \|\widetilde{u}_n\|_{H^1_{kT}}^2 \quad (2.61)$$

combined with the inequality $\|\widetilde{u}_n\|_{L^2} \leq \|\widetilde{u}_n\|_{H^1_{kT}}$ give

$$J_k(u_n) + \left(\varepsilon kT |\overline{u}_n|^2 + kT \bar{e} |\overline{u}_n| \right) + C_\varepsilon kT \geq \|\widetilde{u}_n\|_{H^1_{kT}} \left(\left[\frac{2\pi^2}{4\pi^2 + (kT)^2} - \varepsilon \right] \|\widetilde{u}_n\|_{H^1_{kT}} - \|e\|_{L^2} \right). \quad (2.62)$$

This leads to

$$\lim_{n \rightarrow +\infty} |\overline{u}_n| = +\infty. \quad (2.63)$$

Indeed, if (2.63) does not hold then there would exist a subsequence of $(\overline{u}_n)_{n \in \mathbb{N}}$, still denoted the same, which is bounded. Since $(J_k(u_n))_{n \in \mathbb{N}}$ and \bar{e} are bounded and ε is chosen arbitrarily small, then (2.62) implies that $\|\widetilde{u}_n\|_{H^1_{kT}}$ is bounded. It follows from the inequality

$$\|u_n\|_{H^1_{kT}} \leq \sqrt{kT |\overline{u}_n|} + \|\widetilde{u}_n\|_{H^1_{kT}} \quad (2.64)$$

that $\|u_n\|_{H^1_{kT}}$ is bounded, but this contradicts our assumption $\lim_{n \rightarrow +\infty} \|u_n\|_{H^1_{kT}} = +\infty$. Therefore, (2.63) holds.

Using Wirtinger's inequality

$$\|\widetilde{u}_n\|_{L^2} \leq \frac{kT}{2\pi} \|\widetilde{u}'_n\|_{L^2} \quad (2.65)$$

in (2.60), we get

$$\frac{\|\widetilde{u}'_n\|_{L^2}}{|\overline{u}_n|} \left(\left[\frac{1}{2} - \varepsilon \left(\frac{kT}{2\pi} \right)^2 \right] \frac{\|\widetilde{u}'_n\|_{L^2}}{|\overline{u}_n|} - \left(\frac{kT}{2\pi} \right) \frac{\|e\|_{L^2}}{|\overline{u}_n|} \right) \leq \frac{J_k(u_n)}{|\overline{u}_n|^2} + \varepsilon kT + \frac{kT \bar{e}}{|\overline{u}_n|} + \frac{C_\varepsilon kT}{|\overline{u}_n|^2}. \quad (2.66)$$

It follows from (2.63) that

$$\lim_{n \rightarrow +\infty} \frac{\|\widetilde{u}'_n\|_{L^2}}{|\overline{u}_n|} = 0. \quad (2.67)$$

Using Sobolev's inequality we obtain

$$\lim_{n \rightarrow +\infty} \frac{\|\widetilde{u}_n\|_\infty}{|\overline{u}_n|} \leq \sqrt{\frac{kT}{12}} \lim_{n \rightarrow +\infty} \frac{\|\widetilde{u}_n'\|_{L^2}}{|\overline{u}_n|} = 0. \quad (2.68)$$

The identity $u_n(t) = \overline{u}_n(1 + (\widetilde{u}_n(t)/\overline{u}_n))$ for all $t \in [0, kT]$ and (2.63) imply that

$$\lim_{n \rightarrow +\infty} \min_{t \in [0, kT]} |u_n(t)| = +\infty. \quad (2.69)$$

Assume that $\lim_{n \rightarrow +\infty} \min_{t \in [0, kT]} u_n(t) = +\infty$ (the other case can be treated similarly). Then for n large enough, $u_n(t) > d$ uniformly in $t \in [0, kT]$. By (L3) we have for all $t \in [0, kT]$

$$f_{r_k}(t, u_n(t)) - \bar{e} > 0. \quad (2.70)$$

Consequently, for n large enough

$$\int_0^{kT} |f_{r_k}(t, u_n(t)) - \bar{e}| dt = \int_0^{kT} [f_{r_k}(t, u_n(t)) - \bar{e}] dt = \left| \int_0^{kT} [f_{r_k}(t, u_n(t)) - \bar{e}] dt \right|. \quad (2.71)$$

Since $\lim_{n \rightarrow +\infty} J'_k(u_n) = 0$, then for all $v \in H^1_{kT}$ and for n large enough

$$\left| \int_0^{kT} [u'_n(t)v'(t) - f_{r_k}(t, u_n(t))v(t) + e(t)v(t)] dt \right| \leq \varepsilon_n \|v\|_{H^1_{kT}}, \quad (2.72)$$

where $\varepsilon_n > 0$ for every n , and $\lim_{n \rightarrow +\infty} \varepsilon_n = 0$. In particular, if we take $v(t) = -1$ in the above inequality we obtain for every $n \in \mathbb{N}$

$$\left| \int_0^{kT} [f_{r_k}(t, u_n(t)) - \bar{e}] dt \right| \leq \varepsilon_n \sqrt{kT}, \quad (2.73)$$

which infer

$$\begin{aligned} \int_0^{kT} |f_{r_k}(t, u_n(t))| dt &\leq \int_0^{kT} |f_{r_k}(t, u_n(t)) - \bar{e}| dt + \int_0^{kT} |\bar{e}| dt \\ &= \left| \int_0^{kT} [f_{r_k}(t, u_n(t)) - \bar{e}] dt \right| + \int_0^{kT} |\bar{e}| dt \leq \varepsilon_n \sqrt{kT} + kT|\bar{e}|. \end{aligned} \quad (2.74)$$

Now, taking $v = \widetilde{u}_n$ in (2.72) we obtain

$$\varepsilon_n \|\widetilde{u}_n\|_{H^1_{kT}} \geq \|\widetilde{u}_n'\|_{L^2}^2 - \left| \int_0^{kT} [f_{r_k}(t, u_n(t)) - e(t)] \widetilde{u}_n(t) dt \right|. \quad (2.75)$$

Obviously, we have for n large enough

$$\left| \int_0^{kT} [f_{r_k}(t, u_n(t)) - e(t)] \widetilde{u}_n(t) dt \right| \leq \sup_{t \in [0, kT]} |\widetilde{u}_n(t)| \left(\int_0^{kT} |f_{r_k}(t, u_n(t))| dt + \int_0^{kT} |e(t)| dt \right) \tag{2.76}$$

$$\leq \|\widetilde{u}_n\|_\infty \left(\varepsilon_n \sqrt{kT} + kT|\bar{e}| + \|e\|_{L^1} \right).$$

Thus, for n large enough, (2.75) implies that

$$\varepsilon_n \|\widetilde{u}_n\|_{H_{kT}^1} \geq \|\widetilde{u}_n'\|_{L^2}^2 - \|\widetilde{u}_n\|_\infty \left(\varepsilon_n \sqrt{kT} + kT|\bar{e}| + \|e\|_{L^1} \right). \tag{2.77}$$

Sobolev's inequality

$$\|\widetilde{u}_n\|_\infty \leq \sqrt{\frac{kT}{12}} \|\widetilde{u}_n'\|_{L^2} \leq \sqrt{\frac{kT}{12}} \|\widetilde{u}_n\|_{H_{kT}^1} \tag{2.78}$$

and Wirtinger's inequality combined with (2.77) give, for n large enough,

$$\varepsilon_n \|\widetilde{u}_n\|_{H_{kT}^1} \geq \left(\frac{4\pi^2}{4\pi^2 + (kT)^2} \right) \|\widetilde{u}_n\|_{H_{kT}^1}^2 - \sqrt{\frac{kT}{12}} \|\widetilde{u}_n\|_{H_{kT}^1} \left(\varepsilon_n \sqrt{kT} + kT|\bar{e}| + \|e\|_{L^1} \right). \tag{2.79}$$

So, for n large enough, we deduce that

$$\|\widetilde{u}_n\|_{H_{kT}^1} \leq \ell := \left(\frac{4\pi^2 + (kT)^2}{4\pi^2} \right) \left(1 + \sqrt{\frac{kT}{12}} \left[\sqrt{kT} + kT|\bar{e}| + \|e\|_{L^1} \right] \right). \tag{2.80}$$

Hence $(\widetilde{u}_n)_n$ is bounded in H_{kT}^1 . Consequently $\|u_n'\|_{L^2} = \|\widetilde{u}_n'\|_{L^2} \leq \|\widetilde{u}_n\|_{H_{kT}^1} \leq \ell$. Since $(J_k(u_n))_{n \in \mathbb{N}}$ is bounded, it follows that

$$\int_0^{kT} [F_{r_k}(t, u_n(t)) - e(t)u_n(t)] dt \tag{2.81}$$

is bounded. Hölder's inequality gives

$$\left| \int_0^{kT} e(t) \widetilde{u}_n(t) dt \right| \leq \|e\|_{L^2} \|\widetilde{u}_n\|_{L^2} \leq \|e\|_{L^2} \|\widetilde{u}_n\|_{H_{kT}^1} \leq \ell \|e\|_{L^2}. \tag{2.82}$$

Since

$$\int_0^{kT} F_{r_k}(t, u_n(t)) dt - \bar{u}_n \int_0^{kT} e(t) dt = \int_0^{kT} [F_{r_k}(t, u_n(t)) - e(t)u_n(t)] dt + \int_0^{kT} e(t) \widetilde{u}_n(t) dt, \tag{2.83}$$

it follows that

$$\int_0^{kT} F_{r_k}(t, u_n(t)) dt - \bar{u}_n \int_0^{kT} e(t) dt, \tag{2.84}$$

is bounded. But,

$$\bar{u}_n \int_0^{kT} e(t) dt = kT \bar{e} \bar{u}_n = kT \bar{e} \frac{1}{kT} \int_0^{kT} u_n(t) dt = \int_0^{kT} \bar{e} u_n(t) dt. \tag{2.85}$$

Consequently, there exists $C > 0$ such that

$$\int_0^{kT} [F_{r_k}(t, u_n(t)) - \bar{e} u_n(t)] dt \leq C. \tag{2.86}$$

On the other hand, extending F_{r_k} by T -periodicity we obtain

$$\begin{aligned} \int_0^{kT} [F_{r_k}(t, u_n(t)) - \bar{e} u_n(t)] dt &= \sum_{j=0}^{k-1} \int_0^T [F_{r_k}(t + jT, u_n(t + jT)) - \bar{e} u_n(t + jT)] dt \\ &= \sum_{j=0}^{k-1} \int_0^T [F_{r_k}(t, u_n(t + jT)) - \bar{e} u_n(t + jT)] dt. \end{aligned} \tag{2.87}$$

Setting $x_n = u_n(t + jT)$ for $t \in [0, T]$, we get

$$\int_0^{kT} [F_{r_k}(t, u_n(t)) - \bar{e} u_n(t)] dt = \sum_{j=0}^{k-1} \int_0^T [F_{r_k}(t, x_n) - \bar{e} x_n] dt = k \int_0^T [F_{r_k}(t, x_n) - \bar{e} x_n] dt. \tag{2.88}$$

From (2.75) $|x_n| = |u_n(t + jT)| \rightarrow +\infty$ when $n \rightarrow +\infty$ uniformly in $t \in [0, kT]$. By (L4) we have

$$\lim_{n \rightarrow +\infty} \int_0^{kT} [F_{r_k}(t, u_n(t)) - \bar{e} u_n(t)] dt = k \lim_{|x| \rightarrow +\infty} \int_0^T [F_{r_k}(t, x) - \bar{e} x] dt = +\infty. \tag{2.89}$$

This is a clear contradiction to (2.86). Therefore, $(u_n)_{n \in \mathbb{N}}$ is bounded in H^1_{kT} , and so it has a convergent subsequence. This shows that J_k satisfies the Palais-Smale condition. Next, we show that J_k has a geometry of a Saddle. For, let $u \in H^-$ then we have $u = \widetilde{u}_n$ and $\bar{u} = 0$, so that

$$J_k(\widetilde{u}_n) = \frac{1}{2} \|\widetilde{u}_n\|_{L^2}^2 - \int_0^{kT} [F_{r_k}(t, \widetilde{u}_n(t)) - e(t) \widetilde{u}_n(t)] dt. \tag{2.90}$$

Proceeding as before, we get an inequality similar to (2.72) by replacing u_n by \widetilde{u}_n and \bar{u}_n by 0,

$$J_k(\widetilde{u}_n) \geq \|\widetilde{u}_n\|_{H_{kT}^1} \left(\left[\frac{2\sigma^2}{4\sigma^2 + (kT)^2} - \varepsilon \right] \|\widetilde{u}_n\|_{H_{kT}^1} - \|e\|_{L^2} \right) - C_\varepsilon kT. \quad (2.91)$$

Since ε is chosen arbitrary small, we obtain

$$\lim_{\|\widetilde{u}_n\|_{H_{kT}^1} \rightarrow +\infty} J_k(\widetilde{u}_n) = +\infty, \quad (2.92)$$

which shows that J_k is coercive. Hence, J_k admits a bounded minimizing sequence. Furthermore, J_k is weakly lower semicontinuous on H_{kT}^1 , then

$$\inf_{H^-} J_k > -\infty. \quad (2.93)$$

For s , a constant function, we have $\|s\|_{H_{kT}^1} = |s|\sqrt{kT} \rightarrow +\infty$ if and only if $|s| \rightarrow +\infty$. Then

$$J_k(s) = - \int_0^{kT} [F_{r_k}(t, s) - e(t)s] dt. \quad (2.94)$$

Extending F_{r_k} by T -periodicity we obtain

$$\begin{aligned} J_k(s) &= - \int_0^{kT} [F_{r_k}(t, s) - \bar{e}s] dt = - \sum_{i=0}^{k-1} \int_0^T [F_{r_k}(t + iT, s) - \bar{e}s] dt \\ &= -k \int_0^T [F_{r_k}(t, s) - \bar{e}s] dt. \end{aligned} \quad (2.95)$$

Condition (L4) implies that

$$\lim_{|s| \rightarrow +\infty} [-J_k(s)] = k \lim_{|s| \rightarrow +\infty} \int_0^T [F_{r_k}(t, s) - \bar{e}s] dt = +\infty. \quad (2.96)$$

Hence, for each $k \geq 1$, $(-J_k)$ is coercive on the space of constant functions. Then for each $k \geq 1$, there exists $\eta_k > 0$, large enough, such that

$$J_k(\eta_k) \rightarrow -\infty, \quad \text{and } J_k(-\eta_k) \rightarrow -\infty. \quad (2.97)$$

Thus,

$$\max(J_k(-\eta_k), J_k(\eta_k)) \rightarrow -\infty \quad \text{when } \eta_k \rightarrow +\infty. \quad (2.98)$$

Therefore,

$$\max(J_k(-\eta_k), J_k(\eta_k)) < \inf_{H^-} J_k. \tag{2.99}$$

Let I_{η_k} be the open interval of \mathbb{R} centered at 0 and with radius η_k . Since $H^+ = \mathbb{R}$ it is clear that

$$\partial I_{\eta_k} \cap H^+ = \{-\eta_k, \eta_k\}. \tag{2.100}$$

Therefore, we have

$$\max_{\partial I_{\eta_k} \cap H^+} J_k < \inf_{H^-} J_k, \tag{2.101}$$

with $H^+ \oplus H^- = H_{kT}^1$. Thus J_k has a geometry of Saddle.

Finally, all conditions of the Saddle point theorem are satisfied. Then for each $k \geq 1$, J_k admits a critical point β_k , which is characterized by

$$J_k(\beta_k) = \inf_{\psi \in \Gamma_k} \max_{s \in [-\eta_k, \eta_k]} J_k(\psi(s)), \tag{2.102}$$

where

$$\Gamma_k = \left\{ \psi \in C\left([-\eta_k, \eta_k], H_{kT}^1\right); \psi(-\eta_k) = -\eta_k, \psi(\eta_k) = \eta_k \right\}. \tag{2.103}$$

Thus for each $k \geq 1$, β_k is a weak kT -periodic solution of (2.6) with $\mu = r_k$. If, furthermore, e is assumed continuous, then β_k is a classical solution of (2.6) with $\mu = r_k$.

This completes the proof of Lemma 2.5. □

Remark 2.6. As a consequence of Lemmas 2.2, 2.5 and the Remark 2.3, we conclude that if (H1) and (H2) are satisfied, then (1.1) admits a sequence $(u_k)_{k \geq 1}$ of kT -periodic solutions.

2.3. Existence of Distinct Subharmonic Solutions for (2.6) with $\mu = r_k$

Note that $J_k = J_m$ on $H_{kT}^1 \cap H_{mT}^1$ for each k and m . This justifies the following definition (see [10, Definition 2.1 page 653]).

Definition 2.7. The level of $u \in \cup_k H_{kT}^1$ is defined by $J_m(u)$ when $u \in H_{mT}^1$.

Every functional J_k admits at least a critical level which is given by $\beta_k = \min_{H_{kT}^1} J_k$.

Note that nondistinct subharmonic solutions have the same level. Then we deduce that in order to find the multiplicity of distinct subharmonic solutions, we have to search the multiple critical levels. The sequence $(\beta_k)_{k \geq 1}$ is not always increasing, which means that there exists $m \in \mathbb{N}$ such that if $k \in m\mathbb{N}$ then $\beta_k \leq \beta_m$.

If $\beta_k < \beta_1$ for an integer k , then β_k is not a level for the T -periodic functions and every global minimum of J_k is in fact a subharmonic solution of (2.6).

In our case $J_k(u_k)$ is not necessary a global minimum for J_k and then even if the condition above is verified, it is still insufficient to deduce the existence of true subharmonic solutions. This is why we prove also that the amplitudes and the minimal periods tend to infinity.

Lemma 2.8. *The minimal periods of the solutions u_k of (2.6), with $\mu = r_k$ tend to infinity.*

Proof. Let u_k be a weak solution of (2.6) with $\mu = r_k$. Then u_k is a critical point of J_k . We show that

$$\lim_{k \rightarrow +\infty} \frac{1}{k} J_k(u_k) = -\infty. \quad (2.104)$$

Let $\eta_k \geq k$ and let $\varphi_k \in \Gamma_k$ be defined for all $s \in [-\eta_k, \eta_k]$ and for all $t \in [0, kT]$ by

$$(\varphi_k(s))(t) = s + (\eta_k - |s|). \quad (2.105)$$

We have $(\varphi_k(\eta_k))(t) = \eta_k$ and $(\varphi_k(-\eta_k))(t) = -\eta_k$ for all $t \in [0, kT]$. $(\varphi_k(s))(\cdot)$ is constant with respect to t for all $s \in [-\eta_k, \eta_k]$ and $\varphi_k(s) \in H_{kT}^1$. Let $s_k \in [-\eta_k, \eta_k]$ be such that

$$J_k(\varphi_k(s_k)) = \max_{s \in [-\eta_k, \eta_k]} J_k(\varphi_k(s)). \quad (2.106)$$

We have

$$J_k(u_k) \leq J_k(\varphi_k(s_k)). \quad (2.107)$$

Since $(\varphi_k(s_k))'(t) = 0$ for all $t \in [0, kT]$, (2.54) implies that

$$J_k(\varphi_k(s_k)) = - \int_0^{kT} [F_{r_k}(t, (\varphi_k(s_k))(t)) - e(t)(\varphi_k(s_k))(t)] dt. \quad (2.108)$$

Extending F_{r_k} by T -periodicity, we obtain for $k \geq 2$

$$\begin{aligned} \frac{1}{k} J_k(u_k) &\leq \frac{1}{k} J_k(\varphi_k(s_k)) = -\frac{1}{k} \sum_{j=0}^{k-1} \int_0^{kT} [F_{r_k}(t + jT, (\varphi_k(s_k))(t + jT)) - \bar{e}(\varphi_k(s_k))(t + jT)] dt \\ &= - \int_0^T [F_{r_k}(t, (\varphi_k(s_k))(t)) - \bar{e}(\varphi_k(s_k))(t)] dt. \end{aligned} \quad (2.109)$$

We have $\lim_{k \rightarrow +\infty} |(\varphi_k(s_k))(t)| = +\infty$ for all $t \in [0, kT]$. Apply (L4) with $u = (\varphi_k(s_k))(t)$ to obtain

$$\lim_{k \rightarrow +\infty} \frac{1}{k} J_k(u_k) \leq - \lim_{|u| \rightarrow +\infty} \int_0^T [F_{r_k}(t, u) - \bar{e}u] dt = -\infty. \quad (2.110)$$

Hence (2.104) holds.

Now, assume by contradiction that we can extract from the sequence $(u_k)_{k \geq 1}$, of solutions of (2.6) with $\mu = r_k$ a subsequence whose minimal periods are bounded. Then for this subsequence we can find a common period $k_0 T$. The sequence $(u_n)_{n \geq 1}$ of the critical points of J_{k_0} satisfies

$$J_{k_0}(u_n) = \frac{1}{n} J_n(u_n). \tag{2.111}$$

Assuming $\lim_{n \rightarrow +\infty} \|u_n\|_\infty = +\infty$ and proceeding as before we arrive at the conclusion $(J_{k_0}(u_n))_n$ is bounded and this contradicts (2.104). This completes the proof of Lemma 2.8. \square

Lemma 2.9. *The amplitudes $A_k := (\max_{[0, kT]} u_k - \min_{[0, kT]} u_k)$ of the solutions u_k of equation (2.6) with $\mu = r_k$ tend to infinity.*

Proof. We have to show that $\lim_{k \rightarrow +\infty} A_k = +\infty$. First, we have

$$\lim_{k \rightarrow +\infty} \|u_k\|_\infty = +\infty. \tag{2.112}$$

Otherwise, we can extract from $(u_k)_{k \geq 1}$ a subsequence converging to some u^* with period $n_0 T$, for some $n_0 > 0$. But, this would contradict Lemma 2.8. Next, we must prove that $\lim_{k \rightarrow +\infty} \|\tilde{u}_k\|_\infty = +\infty$. Assume, on the contrary, that $\|\tilde{u}_k\|_\infty$ is bounded. If we suppose further that $(|\tilde{u}_k|)_{k \geq 1}$ is bounded, then $\|u_k\|_\infty$ would be bounded and this contradicts (2.112). Hence there exists a subsequence of $(u_k)_{k \geq 1}$, which we label the same, such that $\lim_{k \rightarrow +\infty} |\bar{u}_k| = +\infty$. This implies $\lim_{k \rightarrow +\infty} \min_{[0, kT]} |u_k(t)| = +\infty$. Then, for k sufficiently large and each $j = 0, 1, \dots, k - 1$, $u_k(t + jT) > d$, uniformly in $t \in (0, T)$. It follows from (L3) that for k large enough $f_{r_k}(t, u_k(t + jT)) - \bar{e} > 0$, uniformly in $t \in (0, T)$. Equation (2.6) with $\mu = r_k$ and the T -periodicity of f_{r_k} with respect to t give

$$\int_0^T \left[\frac{1}{k} \sum_{j=0}^{k-1} f_{r_k}(t, u_k(t + jT)) - \bar{e} \right] dt = 0. \tag{2.113}$$

Hence we can use Fatou's Lemma to obtain

$$0 = \int_0^T \liminf_{k \rightarrow +\infty} \left[\frac{1}{k} \sum_{i=0}^{k-1} f_{r_k}(t, u_k(t + iT)) - \bar{e} \right] dt = \int_0^T \liminf_{x \rightarrow +\infty} [f_{r_k}(t, x) - \bar{e}] dt > 0. \tag{2.114}$$

This is a contradiction. Hence $\lim_{k \rightarrow +\infty} \|\tilde{u}_k\|_\infty = +\infty$, and the proof of Lemma 2.9 is complete. \square





From the above auxiliary results we deduce that equation (2.6) with $\mu = r_k$, and consequently (1.1), admits a sequence $(u_k)_{k \geq 1}$ of distinct kT -periodic solutions whose amplitudes and minimal periods tend to infinity. Thus, the proof of Theorem 1.1 is complete.

Acknowledgments

The authors are grateful to an anonymous referee for suggestions and comments that led to the improvement of the paper. A. Boucherif is grateful to King Fahd University of Petroleum and Minerals for its constant support.

References

- [1] J. Chu and J. J. Nieto, "Recent existence results for second-order singular periodic differential equations," *Boundary Value Problems*, vol. 2009, Article ID 540863, 20 pages, 2009.
- [2] A. Boucherif and N. Daoudi-Merzagui, "Periodic solutions of singular nonautonomous second order differential equations," *Nonlinear Differential Equations and Applications*, vol. 15, no. 1-2, pp. 147–158, 2008.
- [3] N. Daoudi-Merzagui, "Periodic solutions of nonautonomous second order differential equations with a singularity," *Applicable Analysis*, vol. 73, no. 3-4, pp. 449–462, 1999.
- [4] X. Li and Z. Zhang, "Periodic solutions for second-order differential equations with a singular nonlinearity," *Nonlinear Analysis*, vol. 69, no. 11, pp. 3866–3876, 2008.
- [5] J. Mawhin and M. Willem, *Critical Point Theory and Hamiltonian Systems*, vol. 74 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1989.
- [6] P. H. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, vol. 65 of *CBMS Regional Conference Series in Mathematics*, American Mathematical Society, Providence, RI, USA, 1986.
- [7] A. Fonda, *Periodic Solutions of Scalar Second Order Differential Equations with a Singularity*, Académie Royale de Belgique, Brussels, Belgium, 1993.
- [8] A. Fonda, R. Manásevich, and F. Zanolin, "Subharmonic solutions for some second-order differential equations with singularities," *SIAM Journal on Mathematical Analysis*, vol. 24, no. 5, pp. 1294–1311, 1993.
- [9] A. Fonda and M. Ramos, "Large-amplitude subharmonic oscillations for scalar second-order differential equations with asymmetric nonlinearities," *Journal of Differential Equations*, vol. 109, no. 2, pp. 354–372, 1994.
- [10] E. Serra, M. Tarallo, and S. Terracini, "Subharmonic solutions to second-order differential equations with periodic nonlinearities," *Nonlinear Analysis*, vol. 41, pp. 649–667, 2000.
- [11] C.-L. Tang, "Periodic solutions for nonautonomous second order systems with sublinear nonlinearity," *Proceedings of the American Mathematical Society*, vol. 126, no. 11, pp. 3263–3270, 1998.
- [12] J. Yu, "Subharmonic solutions with prescribed minimal period of a class of nonautonomous Hamiltonian systems," *Journal of Dynamics and Differential Equations*, vol. 20, no. 4, pp. 787–796, 2008.
- [13] X. Zhang and X. Tang, "Subharmonic solutions for a class of non-quadratic second order Hamiltonian systems," *Nonlinear Analysis: Real World Applications*, vol. 13, no. 1, pp. 113–130, 2012.
- [14] L. D. Humphreys, P. J. McKenna, and K. M. O'Neill, "High frequency shaking induced by low frequency forcing: periodic oscillations in a spring-cable system," *Nonlinear Analysis. Real World Applications*, vol. 11, no. 5, pp. 4312–4325, 2010.
- [15] P. Omari, G. Villari, and F. Zanolin, "Periodic solutions of the Liénard equation with one-sided growth restrictions," *Journal of Differential Equations*, vol. 67, no. 2, pp. 278–293, 1987.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

