

BOUNDARY REGULARITY OF WEAK SOLUTIONS TO NONLINEAR ELLIPTIC OBSTACLE PROBLEMS

MENG JUNXIA AND CHU YUMING

Received 25 April 2005; Revised 10 September 2005; Accepted 14 September 2005

We study the boundary regularity of weak solutions to nonlinear obstacle problem with $C^{1,\beta}$ -obstacle function, and obtain the $C_{\text{loc}}^{1,\alpha}$ boundary regularity.

Copyright © 2006 M. Junxia and C. Yuming. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

We consider the following variational inequality:

$$\begin{aligned} u \in \mathfrak{R} : \int_{\Omega} A(x, \nabla u) \cdot (\nabla v - \nabla u) dx \\ \geq \int_{\Omega} H(x, u, \nabla u)(v - u) dx + \int_{\Omega} F(x, u) \cdot (\nabla v - \nabla u) dx \end{aligned} \quad (1.1)$$

for all $v \in \mathfrak{R} = \{v \in W_0^{1,p}(\Omega), v \geq \psi \text{ a.e. in } \Omega\}$. Here Ω is a bounded domain in $R^N (N \geq 2)$ with Lipschitz boundary, $2 \leq p \leq N$.

$A(x, \xi) : \Omega \times R^N \rightarrow R^N$ satisfies the following conditions:

- (i) A is a vector valued function, the mapping $x \mapsto A(x, \xi)$ is measurable for all $\xi \in R^N$, $\xi \mapsto A(x, \xi)$ is continuous for a.e. $x \in \Omega$;
- (ii) the homogeneity condition: $A(x, t\xi) = t|t|^{p-2}A(x, \xi)$, $t \in R$, $t \neq 0$;
- (iii) the monotone inequality: $(A(x, \xi) - A(x, \zeta))(\xi - \zeta) \geq a|\xi - \zeta|^p$;
- (iv) $|h||a^{ij}| + |\partial A^i(x, h)/\partial x_j| \leq \tau_1|h|^{p-1}$;
- (v) $\sum_{i,j=1}^N a^{ij}\xi_i\xi_j \geq \tau_2|h|^{p-2}|\xi|^2$;
- (vi) $|A(x, \xi) - A(y, \xi)| \leq b_1(1 + |\xi|^{p-1})|x - y|^{\alpha_0}$;
- (vii) $|A(x, \xi) - A(x, \eta)| \leq b_2||\xi|^{p-2}\xi - |\eta|^{p-2}\eta|$;

where $a^{ij} = \partial A^i / \partial h_j$, a , b_1 , b_2 , τ_1 , τ_2 are positive constants.

2 Boundary regularity

We assume that $H(x, u, \lambda)$, $F(x, u) = \{F_i(x, u)\}_{1 \leq i \leq N}$ in (1.1) are of the form:

$$|H(x, u, \nabla u)| \leq c(|\nabla u|^{p/r'} + |u|^{r-1} + g(x)), \quad (1.2)$$

$$|F(x, u)| \leq c(|u|^{q/p'} + h(x)), \quad (1.3)$$

where $p < q < r$, $r' = r/(r-1)$, $p' = p/(p-1)$, and if $2 \leq p < N$, $r = Np/(N-p)$, while if $p = N$, then r can be some sufficiently large positive number.

Higher regularity of the weak solution to the p -Laplacian obstacle problem

$$I(u) = \inf \left\{ \int_{\Omega} |\nabla u|^p dx : u \in K(\psi) \right\}, \quad (1.4)$$

where

$$K(\psi) = \{v \in W^{1,p}(\Omega) : v \geq \psi \text{ a.e.}\}, \quad (1.5)$$

has been studied by various authors. In the case when ψ is assumed to have only minimal regularity properties, it was shown by [8, 11] that the solution of (1.1) is continuous. In particular, if $\psi \in C^{0,\alpha}(\Omega)$, then the solution u is also an element of $C^{0,\alpha'}(\Omega)$. In the case when $\psi \in C^2(\Omega)$, papers [4, 6, 10, 12] employed different techniques to prove interior $C^{1,\alpha}(\Omega)$ regularity for the solution u to (1.4). Reference [1] gave an interesting result: the condition for $\mathfrak{R}$ to be nonempty is just that ψ should have finite capacity. This implies, among other things, that $\psi^+ = \max(\psi, 0)$ must vanish on $\partial\Omega$, C —almost everywhere. This condition is important for the existence of weak solutions to obstacle problem.

When ψ is smooth (say $C^{1,\alpha}(\Omega)$), the interior regularity of weak solutions to problem (1.1) has been studied extensively by many authors ([3, 13, 14]).

In view of De Giorgi class, paper [2] obtained $C^{0,\alpha}$ interior regularity for solutions of nonlinear elliptic obstacle problem with natural growth in the gradient by taking appropriate test function.

The main concern of these papers is the question of the regularity of the solution u in terms of the given regularity properties of the obstacle ψ and relevant data. This is especially interesting in view of the fact that there is a limit to the amount of regularity that u can inherit from ψ : it is possible for ψ to be real analytic, but u will be at best $C^{1,1}$, that is, have bounded second derivatives.

This paper obtains $C_{\text{loc}}^{1,\alpha}$ boundary regularity of weak solutions to the obstacle problem with $C^{1,\beta}$ -obstacle function under controllable growth condition (1.2). We present a new proof to a useful comparison principle.

2. Notations and preliminaries

Ω is an open bounded subset of R^N , $N \geq 2$; $\partial\Omega$ is the boundary of Ω . If $z \in R^N$, we put

$$\begin{aligned} B_R(z) &= \{x \in R^N : |x - z| < R\}, & \Gamma_R(z) &= \{x \in B_R(z) : x_n = 0\}, \\ B_R^+(z) &= \{x \in B_R(z) : x_n > 0\}, & B_R^-(z) &= \{x \in B_R(z) : x_n < 0\}. \end{aligned} \quad (2.1)$$

We denote by B , B^+ , B^- , Γ , respectively, $B_1(0)$, $B_1^+(0)$, $B_1^-(0)$, $\Gamma_1(0)$. For every set E we denote by $\bar{E}$ its closure, and by $|E|$ its Lebesgue measure. $(f)_R = (1/|B_R|) \int_{B_R} f(x) dx$. The

letter c is used throughout to denote a positive constant, not necessarily the same at each occurrence.

Since $\bar{\Omega}$ is compact, $\partial\Omega$ can be covered by a finite number of neighbourhoods V of its points. It is enough to prove the better regularity of u holds true in $V \cap \Omega$. Since $\partial\Omega$ is a Lipschitz boundary, one can find T which is an invertible Lipschitz mapping such that

$$T(V) = B, \quad T(V \cap \Omega) = B^+, \quad T(V \setminus \bar{\Omega}) = B^-, \quad T(V \cap \partial\Omega) = \Gamma. \quad (2.2)$$

Under the mapping T the variational inequality in Ω is transformed to a variational inequality of the same form in B^+ , for $\bar{u} = u \circ T^{-1}$ which satisfies

$$\begin{aligned} & \int_{B^+} \bar{A}(x, \nabla \bar{u}) \cdot (\nabla v - \nabla \bar{u}) dx \\ & \geq \int_{B^+} \bar{H}(x, \bar{u}, \nabla \bar{u})(v - \bar{u}) dx + \int_{B^+} \bar{F}(x, \bar{u}) \cdot (\nabla v - \nabla \bar{u}) dx, \quad \forall v \in \bar{\mathfrak{R}}, \end{aligned} \quad (2.3)$$

where $\bar{\mathfrak{R}} = \{v \in W_0^{1,p}(B^+), v \geq \psi, \text{ a.e. in } B^+\}$, $\bar{A}$, $\bar{H}$, $\bar{F}$ satisfy assumptions of type (i)–(vii), (1.2), (1.3) with different constants.

In order to simplify the notations, we still denote $\bar{u}$, $\bar{\mathfrak{R}}\bar{A}$, $\bar{H}$, $\bar{F}$ by u , $\mathfrak{R}$, A , H , F , respectively.

Since the original $u \in W_0^{1,p}(\Omega)$, we define then

$$u(x) := \begin{cases} u(x), & \text{if } x \in B^+, \\ -3u(x_1, \dots, x_{n-1}, -x_n) + 4u\left(x_1, \dots, x_{n-1}, -\frac{x_n}{2}\right), & \text{if } x \in B^-. \end{cases} \quad (2.4)$$

In light of Extension theorem [5, page 254], we only need to prove a better regularity of u in B^+ .

Definition 2.1. The function $u \in \mathfrak{R}$ that satisfies (2.3) for all $v \in \mathfrak{R}$ is called a weak solution to the obstacle problem with obstacle ψ .

Definition 2.2. Call $f \in C^{0,\alpha}(\Gamma)$, if for all $x \in \Gamma$, there exists $B_r(x)$ (a ball centered at x of radius r), $r > 0$, such that $f \in C^{0,\alpha}(B_r(x))$.

In the sequel, we will abbreviate $B^+ \cap B_R(y_0) = B_R^+$, $B^+ \cap B_\rho(y_0) = B_\rho^+$, for $0 < \rho < R \leq 1$, the point $y_0 \in \Gamma$ to be understood.

In the following, we will use some lemmas which we state below.

LEMMA 2.3. *Let $w \in W^{1,p}(B_R^+)$ be a solution of the Dirichlet problem*

$$\begin{aligned} & \int_{B_R^+} A(x, \nabla w) \nabla \phi \, dx = 0 \quad \forall \phi \in W_0^{1,p}(B_R^+), \\ & w - u \in W_0^{1,p}(B_R^+). \end{aligned} \quad (2.5)$$

4 Boundary regularity

For $0 < \rho < R/2$, $\sigma \in (0, 1)$,

$$\int_{B_R^+} |\nabla w|^p dx \leq c \int_{B_R^+} |\nabla u|^p dx, \quad (2.6)$$

$$\int_{B_\rho^+} |\nabla w|^p dx \leq c \left(\frac{\rho}{R} \right)^N \int_{B_R^+} |\nabla w|^p dx, \quad (2.7)$$

$$\int_{B_\rho^+} |\nabla w - (\nabla w)_\rho|^p dx \leq c \left(\frac{\rho}{R} \right)^{N+\sigma} \int_{B_R^+} |\nabla w - (\nabla w)_R|^p dx. \quad (2.8)$$

Proof. We can easily get (2.6) by inserting $\phi = w - u$ in (2.5).

An argument similar to the one in [15, Lemma 2.2] shows that (2.7) hold.

The proof of (2.8) is similar to that of [9, Theorem 1.7]. $\square$

LEMMA 2.4. *Let $v \in W^{1,p}(B_R^+)$ be a solution of the Dirichlet problem*

$$\begin{aligned} \int_{B_R^+} A(x, \nabla v) \nabla \phi dx &= \int_{B_R^+} A(x, \nabla \psi) \nabla \phi dx, \\ w - v &\in W_0^{1,p}(B_R^+), \quad \forall \phi \in W_0^{1,p}(B_R^+), \end{aligned} \quad (2.9)$$

then

$$\int_{B_R^+} |\nabla w|^p dx \leq c \int_{B_R^+} |\nabla v|^p dx, \quad (2.10)$$

$$\int_{B_R^+} |\nabla v - \nabla w|^p dx \leq c \int_{B_R^+} |\nabla \psi|^p dx. \quad (2.11)$$

Proof. Formula (2.10) follows immediately from taking $\phi = w - v$ in (2.5).

Inserting $\phi = v - w$ in (2.5) and (2.9), by monotone inequality (iii) and Hölder's inequality, we have

$$\begin{aligned} \int_{B_R^+} |\nabla v - \nabla w|^p dx &\leq c \int_{B_R^+} (A(x, \nabla v) - A(x, \nabla w)) \cdot (\nabla v - \nabla w) dx \\ &= c \int_{B_R^+} A(x, \nabla \psi) \cdot (\nabla v - \nabla w) dx \\ &\leq c \int_{B_R^+} |\nabla \psi|^{p-1} |\nabla v - \nabla w| dx \\ &\leq c \left(\int_{B_R^+} |\nabla \psi|^p dx \right)^{(p-1)/p} \left(\int_{B_R^+} |\nabla v - \nabla w|^p dx \right)^{1/p} \end{aligned} \quad (2.12)$$

from which we get (2.11). $\square$

LEMMA 2.5. *If $v \in W^{1,p}(B_R^+)$ is a solution of the Dirichlet problem (2.9), then $v \geq \psi$ in B_R^+ .*

Proof. It follows from $v = u$ on ∂B_R^+ , $u \in \mathfrak{R}$, that $v \geq \psi$ on ∂B_R^+ . Let $\xi = \min(v, \psi)$, $\xi = \psi$ on ∂B_R^+ , $\xi - \psi \in W_0^{1,p}(B_R^+)$. As test functions in (2.9) we take $\phi = \xi - \psi$, from (2.9) and

monotony inequality (iii), we have

$$\begin{aligned}
0 &= \int_{B_R^+} A(x, \nabla v) - A(x, \nabla \psi) \cdot \nabla (\xi - \psi) dx \\
&= \int_{B_R^+ \cap \{x, v(x) \leq \psi(x)\}} A(x, \nabla v) - A(x, \nabla \psi) \cdot \nabla (v - \psi) dx \\
&\geq a \int_{B_R^+ \cap \{x, v(x) \leq \psi(x)\}} |\nabla v - \nabla \psi|^p dx \\
&= a \int_{B_R^+} |\nabla \xi - \nabla \psi|^p dx
\end{aligned} \tag{2.13}$$

therefore $\xi = \psi$ a.e. in B_R^+ , that is, $v \geq \psi$ a.e. in B_R^+ . $\square$

This lemma is a useful comparison principle, it can be used to obtain the existence or regularity of solutions to elliptic equation or variational inequality.

We extend v to B^+ by setting $v = u$ on $B^+ \setminus B_R^+$, and hence $v \in \mathfrak{R}$. We have the following corollary.

COROLLARY 2.6. *Suppose u is a weak solution to the obstacle problem (2.3), $v \in W^{1,p}(B_R^+)$ is a solution of the Dirichlet problem (2.9), then $v \in \mathfrak{R}$ satisfies the variational inequality*

$$\int_{B^+} A(x, \nabla u) \cdot (\nabla v - \nabla u) dx \geq \int_{B^+} H(x, u, \nabla u)(v - u) dx + \int_{B^+} F(x, u) \cdot (\nabla v - \nabla u) dx. \tag{2.14}$$

LEMMA 2.7. *Assume u is a weak solution to the obstacle problem (2.3), where H, F verify (1.2), (1.3), respectively, $g \in L^t(B^+)$ with $t > N/p$, $h \in L^s(B^+)$ with $s > p'$, v satisfies (2.9), then*

$$\int_{B_R^+} |\nabla v|^p dx \leq c \left[\int_{B_R^+} |\nabla u|^p dx + \int_{B_R^+} |\nabla \psi|^p dx \right], \tag{2.15}$$

$$\begin{aligned}
\int_{B_R^+} |\nabla u - \nabla v|^p dx &\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} \right. \\
&\quad + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} + \left(\int_{B_R^+} |u|^r dx \right)^{q/p} \\
&\quad \left. + R^{Np(1-1/r-1/t)/(p-1)} + R^{N(1-p'/s)} + R^{N(1-p/m)} \right\},
\end{aligned} \tag{2.16}$$

where $\delta = (r - p)/r(p - 1) > 0$.

Proof. By inserting $\phi = v - u$ in (2.9), an application of Hölder's inequality and Young's inequality yields

$$\begin{aligned}
\int_{B_R^+} |\nabla v|^p dx &\leq c \int_{B_R^+} A(x, \nabla v) \cdot \nabla v dx \\
&= c \left[\int_{B_R^+} A(x, \nabla v) \cdot \nabla u dx + \int_{B_R^+} A(x, \nabla \psi) \cdot \nabla (v - u) dx \right]
\end{aligned}$$

6 Boundary regularity

$$\begin{aligned}
& \leq c \left[\int_{B_R^+} |\nabla v|^{p-1} |\nabla u| dx + \int_{B_R^+} |\nabla \psi|^{p-1} |\nabla u - \nabla v| dx \right] \\
& \leq c \left\{ \left(\int_{B_R^+} |\nabla v|^p dx \right)^{(p-1)/p} \left(\int_{B_R^+} |\nabla u|^p dx \right)^{1/p} \right. \\
& \quad \left. + \left(\int_{B_R^+} |\nabla \psi|^p dx \right)^{(p-1)/p} \left(\int_{B_R^+} |\nabla u - \nabla v|^p dx \right)^{1/p} \right\} \\
& \leq c \epsilon_1 \int_{B_R^+} |\nabla v|^p dx + c(\epsilon_1, p) \int_{B_R^+} |\nabla u|^p dx + c \epsilon_2 \int_{B_R^+} |\nabla u - \nabla v|^p dx \\
& \quad + c(\epsilon_2, p) \int_{B_R^+} |\nabla \psi|^p dx \\
& \leq (c \epsilon_1 + c \epsilon_2) \int_{B_R^+} |\nabla v|^p dx + (c(\epsilon_1, p) + c \epsilon_2) \int_{B_R^+} |\nabla u|^p dx \\
& \quad + c(\epsilon_2, p) \int_{B_R^+} |\nabla \psi|^p dx
\end{aligned} \tag{2.17}$$

for $(c \epsilon_1 + c \epsilon_2)$ sufficiently small ($c \epsilon_1 + c \epsilon_2 < 1$), we can get (2.15). $\square$

By $\psi \in W^{1,m}(\Omega)$, $m > N$, we have

$$\int_{B_R} |\nabla \psi|^p dx \leq c \|\nabla \psi\|_{mR}^p R^{N(1-p/m)}. \tag{2.18}$$

Combining monotone inequality (iii), (2.9), (2.14), and (1.2) and using Poincaré's inequality, Hölder's inequality, we have

$$\begin{aligned}
& \int_{B_R^+} |\nabla u - \nabla v|^p dx \\
& \leq c \int_{B_R^+} [A(x, \nabla u) - A(x, \nabla v)] \cdot (\nabla u - \nabla v) dx \\
& \leq c \int_{B_R^+} [H(x, u, \nabla u)(u - v) + (F(x, u) - A(x, \nabla \psi)) \cdot (\nabla u - \nabla v)] dx \\
& \leq c \int_{B_R^+} (|\nabla u|^{p(1-1/r)} + |u|^{r-1} + |g|) |u - v| dx \\
& \quad + c \int_{B_R^+} (|u|^{q/p'} + h(x)) |\nabla u - \nabla v| dx + c \int_{B_R^+} |\nabla \psi|^{p-1} \cdot |\nabla u - \nabla v| dx \\
& \leq c \left[\int_{B_R^+} (|\nabla u|^p + |u|^r + |g|^{r/(r-1)}) dx \right]^{1-1/r} \left(\int_{B_R^+} |u - v|^r dx \right)^{1/r} \\
& \quad + \left(\int_{B_R^+} (|u|^q + |h|^{p'}) dx \right)^{1/p'} \left(\int_{B_R^+} |\nabla u - \nabla v|^p dx \right)^{1/p} \\
& \quad + \int_{B_R^+} |\nabla \psi|^{p-1} \cdot |\nabla u - \nabla v| dx
\end{aligned}$$

$$\begin{aligned}
&\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1-1/r} + \|g\|_t R^{N(1-1/r-1/t)} \right\} \\
&\quad \times R^{1-N(1/p-1/r)} \left(\int_{B_R^+} |\nabla u - \nabla v|^p dx \right)^{1/p} \\
&\quad + c \left\{ \left(\int_{B_R^+} (|u|^p + |\nabla u|^p) dx \right)^{q/pp'} + \|h\|_s R^{N(1/p'-1/s)} + \|\nabla \psi\|_{p/(p-1)}^{p-1} \right\} \\
&\quad \times \left(\int_{B_R^+} |\nabla u - \nabla v|^p dx \right)^{1/p}
\end{aligned} \tag{2.19}$$

since $0 < R \leq 1$, by (2.18), Hölder inequality, Young inequality, we have

$$\begin{aligned}
&\int_{B_R^+} |\nabla u - \nabla v|^p dx \\
&\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} \right. \\
&\quad + \left(\int_{B_R^+} |u|^p dx \right)^{q/p} + \|g\|_t^{p/(p-1)} R^{Np(1-1/r-1/t)/(p-1)} \\
&\quad \left. + \|h\|_s^{p/(p-1)} R^{N(1-p'/s)} + \int_{B_R^+} |\nabla \psi|^p dx \right\} \\
&\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} \right. \\
&\quad + \left(\int_{B_R^+} |u|^p dx \right)^{q/p} + R^{Np(1-1/r-1/t)/(p-1)} + R^{N(1-p'/s)} + R^{N(1-p/m)} \left. \right\} \\
&\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} \right. \\
&\quad + \left(\int_{B_R^+} |u|^r dx \right)^{q/r} |B_R|^{(q/p)(1-p/r)} + R^{Np(1-1/r-1/t)/(p-1)} + R^{N(1-p'/s)} + R^{N(1-p/m)} \left. \right\} \\
&\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} \right. \\
&\quad + \left(\int_{B_R^+} |u|^r dx \right)^{q/p} + R^{N(q/p)} + R^{Np(1-1/r-1/t)/(p-1)} + R^{N(1-p'/s)} + R^{N(1-p/m)} \left. \right\} \\
&\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} \right. \\
&\quad + \left(\int_{B_R^+} |u|^r dx \right)^{q/p} + R^{Np(1-1/r-1/t)/(p-1)} + R^{N(1-p'/s)} + R^{N(1-p/m)} \left. \right\}
\end{aligned} \tag{2.20}$$

which implies (2.16).

8 Boundary regularity

3. $C^{0,\lambda}$ regularity

THEOREM 3.1. *Assume that $H(x, u, \nabla u)$ satisfies (1.2), $g \in L^t(B^+)$ with $t > N/p$, $F(x, u)$ satisfies (1.3), $h \in L^s(B^+)$ with $s > N/(p-1)$, and $\psi \in W^{1,m}(B^+)$ with $m > N$. If $u \in \mathfrak{R}$ makes (2.3) hold, then $u \in C^{0,\lambda}(\Gamma)$ with $\lambda = \min\{1 - N(1/t + 1/r - 1/p)/(p-1), 1 - N/(s(p-1), 1 - N/m)\}$.*

Before proceeding with the formal proof, we make an important observation. It is a well-known result.

PROPOSITION 3.2. *If $f \in W^{1,p}(\Omega)$, then for all constants $k \in \mathbb{R}^N$,*

$$\int_{B_\rho} |\nabla f - (\nabla f)_{\rho, x_0}|^p dx \leq C(p) \int_{B_\rho} |\nabla f - k|^p dx \quad (3.1)$$

for every ρ for which $B_\rho(x_0) \subset \Omega$.

Proof. By elementary inequality, we have

$$\int_{B_\rho} |\nabla f - (\nabla f)_{\rho, x_0}|^p dx \leq C(p) \left[\int_{B_\rho} |\nabla f - k|^p dx + \int_{B_\rho} |k - (\nabla f)_{\rho, x_0}|^p dx \right]. \quad (3.2)$$

Moreover

$$\begin{aligned} \int_{B_\rho} |k - (\nabla f)_{\rho, x_0}|^p dx &= |B_\rho| |k - (\nabla f)_{\rho, x_0}|^p = |B_\rho| \left| k - \frac{1}{|B_\rho|} \int_{B_\rho} \nabla f dx \right|^p \\ &= |B_\rho| \left| \frac{1}{|B_\rho|} \int_{B_\rho} (k - \nabla f) dx \right|^p = |B_\rho|^{1-p} \left| \int_{B_\rho} (k - \nabla f) dx \right|^p \\ &\leq |B_\rho|^{1-p} \int_{B_\rho} |\nabla f - k|^p dx |B_\rho|^{p(1-1/p)} = \int_{B_\rho} |\nabla f - k|^p dx. \end{aligned} \quad (3.3)$$

Therefore (3.1) holds for any $k \in \mathbb{R}^N$. □

Proof of Theorem 3.1. To get the regularity, we need to prove the following inequality:

$$\int_{B_\rho^+} |\nabla u|^p dx \leq c \rho^{N-p+\rho\lambda}. \quad (3.4)$$

Let us consider three different situations.

- (1) If $B_{2R}(y_0) \subset B^+$, inequality (3.4)–(4.1) has been proved in [13], since it is related to interior regularity.
- (2) If $B_R(y_0) \subset B^-$, by Extension theorem [5, page 254], if we can get $C^{1,\alpha}$ regularity of u in B^+ , we can deduce the same result for u in B^- , so we need not care about this situation.

(3) If $B_R(y_0) \cap B^+ \neq \emptyset$, we also give three different situations as follows:

- (a) $y_0 \in \Gamma$,
- (b) $y_0 \in B^-$,
- (c) $y_0 \in B^+$.

We only prove the situation (a), since the others can be transformed into the situation (a) or the interior regularity situation by applying the finitely covered theorem, see [13]. Assume $h \in L^s(B^+, R^N)$ with $s > N/(p-1)$, $\psi \in W^{1,m}(B^+)$ with $m > N$, we see that

$$\begin{aligned} \int_{B_R^+} |h|^{p/(p-1)} &\leq \|h\|_s^{p/(p-1)} R^{N[1-p/s(p-1)]}, \\ \int_{B_R^+} \left| |\nabla \psi|^{p-2} \nabla \psi - (|\nabla \psi|^{p-2} \nabla \psi)_R \right|^{p/(p-1)} dx &\leq c \int_{B_R^+} |\nabla \psi|^p dx \leq c \|\nabla \psi\|_m^p R^{N(1-p/m)}. \end{aligned} \quad (3.5)$$

Combining (2.7), (2.10), (2.11), (2.16), and (3.5), we have

$$\begin{aligned} &\int_{B_\rho^+} |\nabla u|^p dx \\ &\leq c \int_{B_\rho^+} |\nabla w|^p dx + c \int_{B_\rho^+} |\nabla u - \nabla w|^p dx \\ &\leq c \left(\frac{\rho}{R} \right)^N \int_{B_R^+} |\nabla u|^p dx + c \int_{B_R^+} |\nabla u - \nabla v|^p dx + c \int_{B_R^+} |\nabla v - \nabla w|^p dx \\ &\leq c \left(\frac{\rho}{R} \right)^N \int_{B_R^+} |\nabla u|^p dx \\ &\quad + c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} + \left(\int_{B_R^+} |u|^r dx \right)^{q/p} \right. \\ &\quad \left. + R^{Np(1-1/r-1/t)/(p-1)} + R^{N[1-p/s(p-1)]} + R^{N(1-p/m)} \right\} \\ &\leq c \left(\frac{\rho}{R} \right)^N \int_{B_R^+} |\nabla u|^p dx + c \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} \\ &\quad + c \left(\int_{B_R^+} |\nabla u|^p dx \right)^{q/p} + c \left(\int_{B_R^+} |u|^r dx \right)^{q/p} + c R^{N-p+\lambda}, \end{aligned} \quad (3.6)$$

where $\lambda = \min\{1 - N(1/t + 1/r - 1/p)/(p-1), 1 - N/s(p-1), 1 - N/m\}$.

By $t > N/p$, we have the following.

- (i) If $2 \leq p < N$, then $1/t < p/N$, $1/t + 1/r - 1/p < p/N + (N-p)/N - 1/p = (p-1)/N$, $N(1/t + 1/r - 1/p)/(p-1) < 1$.
- (ii) If $p = N$, by $t > 1$, we can assume that r is a positive number sufficiently large, such that: $1/t + 1/r < 1$, so $N(1/t + 1/r - 1/N)/(N-1) < (N/(N-1))(1 - 1/N) = 1$.

Hence, if $2 \leq p \leq N$, we always have $N(1/t + 1/r - 1/p)/(p-1) < 1$.

Using $s > N/(p-1)$, $m > N$, by the definition of λ , we see that: $0 < \lambda < 1$.

10 Boundary regularity

In the meantime, by Poincaré's inequality and Hölder's inequality, we also have

$$\begin{aligned} \int_{B_p^+} |u|^r dx &\leq c \int_{B_p^+} |u_R|^r dx + c \int_{B_R^+} |u - u_R|^r dx \\ &\leq c \left(\frac{\rho}{R} \right)^N \int_{B_R^+} |u|^r dx + c R^{r[1-N(1/p-1/r)]} \left(\int_{B_R^+} |\nabla u|^p dx \right)^{r/p}, \end{aligned} \quad (3.7)$$

where

$$r \left[1 - N \left(\frac{1}{p} - \frac{1}{r} \right) \right] = \begin{cases} \frac{Np}{N-p} \left[1 - N \left(\frac{1}{p} - \frac{N-p}{Np} \right) \right] = 0, & \text{if } 2 \leq p < N; \\ r \left[1 - N \left(\frac{1}{N} - \frac{1}{r} \right) \right] = N, & \text{if } p = N. \end{cases} \quad (3.8)$$

Adding (3.7) to (3.6) and setting

$$\phi(R) = \int_{B_R^+} (|\nabla u|^p + |u|^r) dx, \quad (3.9)$$

we obtain

$$\phi(\rho) \leq c \left[\left(\frac{\rho}{R} \right)^N + \chi(R) \right] \phi(R) + c R^{N-p+p\lambda}, \quad (3.10)$$

where

$$\begin{aligned} \chi(R) &= \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^\delta \\ &\quad + \left(\int_{B_R^+} |\nabla u|^p dx \right)^{(q-p)/p} + \left(\int_{B_R^+} |u|^r dx \right)^{(q-p)/p} \\ &\quad + \begin{cases} \left(\int_{B_R^+} |\nabla u|^p dx \right)^{(r-p)/p}, & \text{if } 2 \leq p < N; \\ R^N \left(\int_{B_R^+} |\nabla u|^p dx \right)^{(r-p)/p}, & \text{if } p = N. \end{cases} \end{aligned} \quad (3.11)$$

We can always get $\chi(R) \rightarrow 0$ as $R \rightarrow 0^+$. Applying [7, page 86, Lemma 2.1], we deduce that for ρ sufficiently small,

$$\int_{B_p^+} |\nabla u|^p dx \leq \phi(\rho) \leq c \rho^{N-p+p\lambda}. \quad (3.12)$$

□

By Dirichlet growth theorem (see [7, page 64, Theorem 1.1]), $u \in C_{\text{loc}}^{0,\lambda}(\Gamma)$.

4. C^{1,α_1} regularity

THEOREM 4.1. *Assume that $H(x, u, \nabla u)$ satisfies (1.2), $g \in L^t(B^+)$ with $t > N$, $F \in C^{0,\beta}(B^+, R)$ with $\beta > 0$, $\psi \in C^{1,\gamma}(B^+)$ for some $\gamma > 0$. If $u \in \mathfrak{X}$ makes (2.3) hold, then $u \in C^{1,\alpha_1}(\Gamma)$ for some $0 < \alpha_1 < \sigma/p$.*

To get our result, we need to prove the following inequality:

$$\int_{B_\rho^+} |\nabla u - (\nabla u)_\rho|^p dx \leq c\rho^{N+p\alpha_1}. \quad (4.1)$$

It is easy to see that $|\nabla \psi|^{p-2} \nabla \psi \in C^{0,\gamma}(B^+)$ if $\psi \in C^{1,\gamma}(B^+)$ and $2 \leq p \leq N$.

Utilizing the conditions of Theorem 4.1, we see that:

$$\int_{B_R^+} |F - F_R|^{p/(p-1)} dx \leq \|F\|_{C^{0,\beta}(B^+)}^{p/(p-1)} R^{N+\beta p/(p-1)}, \quad (4.2)$$

$$\begin{aligned} & |\nabla \psi(x)|^{p-2} \nabla \psi(x) - |\nabla \psi(y)|^{p-2} \nabla \psi(y) \\ & \leq \| |\nabla \psi|^{p-2} \nabla \psi \|_{C^{0,\gamma}(B^+)} |x - y|^\gamma, \quad \forall x, y \in B^+. \end{aligned} \quad (4.3)$$

By $\psi \in C^{1,\gamma}(B^+)$, we can get $|\nabla \psi|^{p-1} \leq c$, so combining condition (vi) we have

$$|A(x, \nabla \psi) - A(y, \nabla \psi)| \leq c|x - y|^{\alpha_0}. \quad (4.4)$$

By condition (vii), (4.3), and (4.4), we see that for all $\phi \in W_0^{1,p}(B_R^+)$, there holds

$$\begin{aligned} & \int_{B_R^+} A(x, \nabla \psi) \cdot \nabla \phi dx \\ &= \int_{B_R^+} (A(x, \nabla \psi) - (A(x, \nabla \psi))_R) \cdot \nabla \phi dx \\ &= \int_{B_R^+} \int_{B_R} [A(x, \nabla \psi(x)) - A(y, \nabla \psi(y))] \cdot \nabla \phi(x) dy dx \\ &= \int_{B_R^+} \int_{B_R} [A(x, \nabla \psi(x)) - A(x, \nabla \psi(y)) \\ & \quad + A(x, \nabla \psi(y)) - A(y, \nabla \psi(y))] \cdot \nabla \phi(x) dy dx \\ &\leq c \int_{B_R^+} \int_{B_R} (|x - y|^{\alpha_0} + ||\nabla \psi(x)|^{p-2} \nabla \psi(x) - |\nabla \psi(y)|^{p-2} \nabla \psi(y)|) |\nabla \phi(x)| dy dx \\ &\leq c \int_{B_R^+} (R^{\alpha_0} + R^\gamma) |\nabla \phi(x)| dx. \end{aligned} \quad (4.5)$$

From last formula, we see that

$$\int_{B_R^+} A(x, \nabla \psi) \cdot \nabla \phi dx \leq c \int_{B_R^+} (R^{\alpha_0} + R^\gamma) |\nabla \phi(x)| dx. \quad (4.6)$$

In the following, we give two lemmas which will be used in the proof of Theorem 4.1.

12 Boundary regularity

LEMMA 4.2. Assume that $\psi \in C^{1,\gamma}(B^+)$, $w, v \in W^{1,p}(B_R^+)$ solve the Dirichlet problem (2.5), (2.9), respectively, then there holds

$$\int_{B_R^+} |\nabla v - \nabla w|^p dx \leq c \left(R^{N+\gamma p/(p-1)} + R^{N+\alpha_0 p/(p-1)} \right). \quad (4.7)$$

Proof. Inserting $\phi = v - w$ in (2.5) and (2.9), by (4.6), monotone inequality (iii), and Hölder inequality, we have

$$\begin{aligned} \int_{B_R^+} |\nabla v - \nabla w|^p dx &\leq c \int_{B_R^+} (A(x, \nabla v) - A(x, \nabla w)) \cdot (\nabla v - \nabla w) dx \\ &= c \int_{B_R^+} A(x, \nabla \psi) \cdot (\nabla v - \nabla w) dx \\ &\leq c \int_{B_R^+} (R^{\alpha_0} + R^\gamma) |\nabla v - \nabla w| dx \\ &\leq c (R^{\gamma+N(p-1)/p} + R^{\alpha_0+N(p-1)/p}) \left(\int_{B_R^+} |\nabla v - \nabla w|^p dx \right)^{1/p}. \end{aligned} \quad (4.8)$$

From last formula, we get (4.7). $\square$

LEMMA 4.3. Assume that $A(x, \xi)$ satisfies condition (i)–(vii), u is a weak solution to obstacle problem (1.1), where H verifies (1.2), $g \in L^t(B^+)$, $t > N$; $F \in C^{0,\beta}(B^+)$, $\beta > 0$; $\psi \in C^{1,\gamma}(B^+)$, $\gamma > 0$. $v \in W^{1,p}(B_R^+)$ solves the Dirichlet problem (2.9), then there holds

$$\begin{aligned} \int_{B_R^+} |\nabla u - \nabla v|^p dx &\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + R^{Np(1-1/r-1/t)/(p-1)} \right. \\ &\quad \left. + R^{N+\beta p/(p-1)} + R^{N+\gamma p/(p-1)} + R^{N+\alpha_0 p/(p-1)} \right\}, \end{aligned} \quad (4.9)$$

where $\delta = (r - p)/r(p - 1) > 0$.

Proof. Combining monotone inequality (iii), and (1.2), (2.14), (2.18), (4.6), Hölder inequality, we have

$$\begin{aligned} \int_{B_R^+} |\nabla u - \nabla v|^p dx &\leq c \int_{B_R^+} [A(x, \nabla u) - A(x, \nabla v)] \cdot (\nabla u - \nabla v) dx \\ &\leq c \int_{B_R^+} [H(x, u, \nabla u)(u - v) + (F(x, u) - A(x, \nabla \psi)) \cdot (\nabla u - \nabla v)] dx \\ &\leq c \int_{B_R^+} (|\nabla u|^{p(1-1/r)} + |u|^{r-1} + |g|) |u - v| dx \\ &\quad + c \int_{B_R^+} |F - F_R| |\nabla u - \nabla v| dx + c \int_{B_R^+} (R^{\alpha_0} + R^\gamma) |\nabla u - \nabla v| dx \end{aligned}$$

$$\begin{aligned}
&\leq c \left\{ \left(\int_{B_R^+} (|\nabla u|^p + |u|^r + |g|^{r/(r-1)}) dx \right)^{1-1/r} \left(\int_{B_R^+} |u-v|^r dx \right)^{1/r} \right. \\
&\quad + \left(\int_{B_R^+} |F-F_R|^{p'} dx \right)^{1/p'} \left(\int_{B_R^+} |\nabla u - \nabla v|^p dx \right)^{1/p} \\
&\quad \left. + \int_{B_R^+} (R^{\alpha_0} + R^\gamma) |\nabla u - \nabla v| dx \right\} \\
&\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1-1/r} + \|g\|_t R^{N(1-1/r-1/t)} \right\} \\
&\quad \times R^{1-N(1/p-1/r)} \left(\int_{B_R^+} |\nabla u - \nabla v|^p dx \right)^{1/p} \\
&\quad + c \left\{ \left(\int_{B_R^+} |F-F_R|^{p'} dx \right)^{1/p'} + R^{\gamma+N(p-1)/p} + R^{\alpha_0+N(p-1)/p} \right\} \\
&\quad \times \left(\int_{B_R^+} |\nabla u - \nabla v|^p dx \right)^{1/p}.
\end{aligned} \tag{4.10}$$

Since $0 < R \leq 1$, by (4.2), and last formula, we have

$$\begin{aligned}
\int_{B_R^+} |\nabla u - \nabla v|^p dx &\leq c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} + R^{Np(1-1/r-1/t)/(p-1)} \right. \\
&\quad \left. + R^{N+\beta p/(p-1)} + R^{N+\gamma p/(p-1)} + R^{N+\alpha_0 p/(p-1)} \right\}.
\end{aligned} \tag{4.11}$$

Hence (4.9) holds. □

Similar to the proof of (3.12), we get for any $\mu \in (0, 1)$

$$\int_{B_\rho^+} |\nabla u|^p dx, \quad \int_{B_\rho^+} |\nabla v|^p dx, \quad \int_{B_\rho^+} |u|^r dx \leq c \rho^{N-p+p\mu}. \tag{4.12}$$

In view of $t > N$, we have the following.

- (i) If $2 \leq p < N$, $1 - 1/r - 1/t > 1 - 1/r - 1/N = (p-1)/p$, from which we get $Np(1 - 1/r - 1/t)/(p-1) > N$.
- (ii) If $p = N$, we can assume that r is a positive number large such that $1/t + 1/r < 1/N$, $Np(1 - 1/r - 1/t)/(p-1) > N^2(1 - 1/N)/(N-1) = N$.

Hence, we always have $Np(1 - 1/r - 1/t)/(p-1) > N$ when $2 \leq p \leq N$.

In the following, we prove Theorem 4.1.

14 Boundary regularity

By (4.2), (4.3), (4.9), Lemmas 2.3, 4.2, and 4.3, we have

$$\begin{aligned}
& \int_{B_\rho^+} |\nabla u - (\nabla u)_\rho|^p dx \\
& \leq c \int_{B_\rho^+} |\nabla u - (\nabla w)_\rho|^p dx \\
& \leq c \int_{B_\rho^+} |\nabla w - (\nabla w)_\rho|^p dx + c \int_{B_\rho^+} |\nabla u - \nabla v|^p dx + c \int_{B_\rho^+} |\nabla v - \nabla w|^p dx \\
& \leq c \int_{B_\rho^+} |\nabla w - (\nabla w)_\rho|^p dx \\
& \quad + c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} \right. \\
& \quad \left. + R^{Np(1-1/r-1/t)/(p-1)} + R^{N+\beta p/(p-1)} + R^{N+\gamma p/(p-1)} + R^{N+\alpha_0 p/(p-1)} \right\} \\
& \quad + c \left(R^{N+\gamma p/(p-1)} + R^{N+\alpha_0 p/(p-1)} \right) \\
& \leq c \left(\frac{\rho}{R} \right)^{N+\sigma} \int_{B_R^+} |\nabla u - (\nabla u)_R|^p dx \\
& \quad + c \left\{ \left[\int_{B_R^+} (|\nabla u|^p + |u|^r) dx \right]^{1+\delta} \right. \\
& \quad \left. + R^{Np(1-1/r-1/t)/(p-1)} + R^{N+\beta p/(p-1)} + R^{N+\gamma p/(p-1)} + R^{N+\alpha_0 p/(p-1)} \right\} \\
& \leq c \left(\frac{\rho}{R} \right)^{N+\sigma} \int_{B_R^+} |\nabla u - (\nabla u)_R|^p dx \\
& \quad + c \left(R^{(1+\delta)(N-p+p\mu)} + R^{Np(1-1/r-1/t)/(p-1)} + R^{N+\beta p/(p-1)} \right. \\
& \quad \left. + R^{N+\gamma p/(p-1)} + R^{N+\alpha_0 p/(p-1)} \right).
\end{aligned} \tag{4.13}$$

We can select μ sufficiently close to 1, such that $(1+\delta)(N-p+p\mu) > N$. Hence we get

$$\int_{B_\rho^+} |\nabla u - (\nabla u)_\rho|^p dx \leq c \left(\frac{\rho}{R} \right)^{N+\sigma} \int_{B_R^+} |\nabla u - (\nabla u)_R|^p dx + c R^{N+p\alpha_1} \tag{4.14}$$

for some $0 < \alpha_1 < \sigma/p$.

Applying [7, page 86, Lemma 2.1], we get

$$\int_{B_\rho^+} |\nabla u - (\nabla u)_\rho|^p dx \leq c \rho^{N+p\alpha_1} \tag{4.15}$$

for ρ sufficiently small. By [7, page 72, Theorem 1.3], we obtain that $u \in C_{\text{loc}}^{1,\alpha_1}(\Gamma)$.

Acknowledgments

We would like to thank professor Fang AiNong for many useful comments on the subject, the project supported by the Natural Science Foundation of Hebei University, the National Natural Science Foundation of China under Grant 10471039, and the Zhejiang Province Natural Science Foundation of China under Grant M103087.

References

- [1] D. R. Adams, *Capacity and the obstacle problem*, Applied Mathematics and Optimization. An International Journal **8** (1982), no. 1, 39–57.
- [2] F. Chiaccchio, *Regularity for solutions of nonlinear elliptic equations with natural growth in the gradient*, Bulletin des Sciences Mathématiques **124** (2000), no. 1, 57–74.
- [3] H. J. Choe, *A regularity theory for a general class of quasilinear elliptic partial differential equations and obstacle problems*, Archive for Rational Mechanics and Analysis **114** (1991), no. 4, 383–394.
- [4] H. J. Choe and J. L. Lewis, *On the obstacle problem for quasilinear elliptic equations of p Laplacian type*, SIAM Journal on Mathematical Analysis **22** (1991), no. 3, 623–638.
- [5] L. C. Evans, *Partial Differential Equations*, Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Rhode Island, 1998.
- [6] M. Fuchs, *Hölder continuity of the gradient for degenerate variational inequalities*, Nonlinear Analysis. Theory, Methods & Applications. An International Multidisciplinary Journal. Series A: Theory and Methods **15** (1990), no. 1, 85–100.
- [7] M. Giaquinta, *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Annals of Mathematics Studies, vol. 105, Princeton University Press, New Jersey, 1983.
- [8] J. Heinonen and T. Kilpeläinen, *On the Wiener criterion and quasilinear obstacle problems*, Transactions of the American Mathematical Society **310** (1988), no. 1, 239–255.
- [9] G. M. Lieberman, *The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations*, Communications in Partial Differential Equations **16** (1991), no. 2-3, 311–361.
- [10] P. Lindqvist, *Regularity for the gradient of the solution to a nonlinear obstacle problem with degenerate ellipticity*, Nonlinear Analysis. Theory, Methods & Applications. An International Multidisciplinary Journal. Series A: Theory and Methods **12** (1988), no. 11, 1245–1255.
- [11] J. H. Michael and W. P. Ziemer, *Interior regularity for solutions to obstacle problems*, Nonlinear Analysis. Theory, Methods & Applications. An International Multidisciplinary Journal. Series A: Theory and Methods **10** (1986), no. 12, 1427–1448.
- [12] J. Mu, *Higher regularity of the solution to the p -Laplacian obstacle problem*, Journal of Differential Equations **95** (1992), no. 2, 370–384.
- [13] Z. Tan and Z. Q. Yan, *Regularity of weak solutions to some degenerate elliptic equations and obstacle problems*, Northeastern Mathematical Journal. Dongbei Shuxue **9** (1993), no. 2, 143–156.
- [14] J. Yang, *Regularity of weak solutions to quasilinear elliptic obstacle problems*, Acta Mathematica Scientia. Series B. English Edition **17** (1997), no. 2, 159–166.
- [15] S. Z. Zheng, *Partial regularity of A -harmonic systems of equations and quasiregular mappings*, Chinese Annals of Mathematics. Series A. Shuxue Niankan. A Ji **19** (1998), no. 1, 63–72 (Chinese), translation in Chinese Journal of Contemporary Mathematics **19** (1998), no. 1, 19–30.

Meng Junxia: College of Mathematics and Computer Science, Hebei University, Hebei 071002, China
E-mail address: junxia@hbu.cn

Chu Yuming: College of Science, Huzhou University, Zhejiang 313000, China
E-mail address: chuyuming@hutc.zj.cn

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskkklai@cityu.edu.hk