

## Research Article

# Multiple Nodal Solutions for Some Fourth-Order Boundary Value Problems via Admissible Invariant Sets

Yang Yang,<sup>1,2</sup> Jihui Zhang,<sup>1</sup> and Zhitao Zhang<sup>3</sup>

<sup>1</sup> Institute of Mathematics, School of Mathematics and Computer Sciences, Nanjing Normal University, Nanjing 210097, China

<sup>2</sup> School of Science, Jiangnan University, Wuxi 214122, China

<sup>3</sup> Academy of Mathematics and System Sciences, Institute of Mathematics, the Chinese Academy of Sciences, Beijing 100080, China

Correspondence should be addressed to Zhitao Zhang, zzt@math03.math.ac.cn

Received 6 May 2008; Accepted 16 September 2008

Recommended by Donal O'Regan

Existence and multiplicity results for nodal solutions are obtained for the fourth-order boundary value problem (BVP)  $u^{(4)}(t) = f(t, u(t))$ ,  $0 < t < 1$ ,  $u(0) = u(1) = u''(0) = u''(1) = 0$ , where  $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous. The critical point theory and admissible invariant sets are employed to discuss this problem.

Copyright © 2008 Yang Yang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

In this paper, we consider the existence of nodal solutions to the semilinear fourth-order equation:

$$u^{(4)}(t) = f(t, u(t)), \quad 0 < t < 1, \quad (1.1)$$

$$u(0) = u(1) = u''(0) = u''(1) = 0, \quad (1.2)$$

where  $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous.

Owing to the importance of *higher-order* differential equations in physics, the existence and multiplicity of the solutions to such problems have been studied by many authors. They obtained the existence of solutions by the cone expansion or compression fixed point theorem [1–6]; sub-sup solution method [7–9]; critical point theory [10–13]; Morse theory [14, 15]; and eta [16, 17]. There are also papers which study nodal solutions for elliptic

equations [18, 19]. In particular, in [20], Han and Li obtained multiple positive, negative, and sign-changing solutions by combining the critical point theory and the method of sub-sup solutions for the (BVP) (1.2). The main result is as follows:

(H<sub>1</sub>) there exist a strict subsolution  $\alpha$  and a strict supersolution  $\beta$  of (BVP) (1.2) with  $\alpha < \beta$ ,  $\alpha(0) = \alpha(1) = \alpha''(0) = \alpha''(1) = 0$ , and  $\beta(0) = \beta(1) = \beta''(0) = \beta''(1) = 0$ ;

(H<sub>2</sub>)  $f(t, u)$  is strictly increasing in  $u$ ;

(H<sub>3</sub>)  $f(t, u)$  is locally Lipschitz continuous in  $u$ ;

(H<sub>4</sub>) there exist  $\mu \in (0, 1/2)$  and  $\Lambda > 0$  such that  $0 < F(t, u) = \int_0^1 f(t, v) dv \leq \mu u f(t, u)$  for all  $|u| \geq \Lambda$  and  $t \in [0, 1]$ .

**Theorem 1.1** (see [20]). *Assume that (H<sub>1</sub>)–(H<sub>4</sub>) hold. Then, (BVP) (1.2) has at least four solutions.*

Motivated by their ideas, we cannot help wondering if there are no strict subsolution and supersolution of (BVP) (1.2), can we still get the nodal solutions just by critical point theory? In this paper, we will use the admissible invariant sets and critical point theory to settle this problem. But we should point out that in all theorems of our paper, the nonlinearity  $f(t, u)$  is assumed to be odd in  $u$ , while no such symmetry is required in [20].

The paper is organized as follows: in Section 2, we give some preliminaries, including the critical point theorems which will be used in our main results and some concepts concerning the partially ordered Banach space. The main results and proofs are established in Section 3.

## 2. Preliminaries

Let  $E$  be a Hilbert space and  $X \subset E$  a Banach space densely embedded in  $E$ . Assume that  $E$  has a closed convex cone  $P_E$  and that  $P := P_E \cap X$  has interior points in  $X$ , that is,  $P = \overset{\circ}{P} \cup \partial P$  with  $\overset{\circ}{P}$  the interior and  $\partial P$  the boundary of  $P$  in  $X$ .

Let  $J \in C^1(E, \mathbb{R})$  and  $J'(u) = u - A(u)$  for  $u \in E$ . We use the following notation:  $K = K(J) = \{u \in E : J'(u) = 0\}$ ,  $J^b = \{u \in E : J(u) \leq b\}$ ,  $K_c = \{u \in E : J(u) = c, J'(u) = 0\}$ ,  $K([a, b]) = \{u \in E : J(u) \in [a, b], J'(u) = 0\}$  for  $a, b, c \in \mathbb{R}$ . Let  $\|\cdot\|$  and  $\|\cdot\|_X$  denote the norms in  $E$  and  $X$ , respectively.

**Lemma 2.1** (see [21]). *Assume  $E$  is a Hilbert space, and  $M$  is a closed convex set of  $E$ ,  $J'(u) = u - A(u)$ , and  $A(M) \subset M$ . Then, there exists a pseudogradient vector field  $W := -\text{id} + B$  for  $J$ , and  $B(M) \subset M$ . Furthermore, if  $J$  is even,  $M = -M$ , then  $W$  is odd.*

Consider the pseudogradient flow  $\sigma$  on  $E$  associated with the vector field  $W = -\text{id} + B$ ,

$$\begin{aligned} \frac{d}{dt}\sigma(t, u) &= -W(\sigma(t, u)), \quad t \geq 0; \\ \sigma(0, u) &= u. \end{aligned} \tag{2.1}$$

We see that  $\sigma$  is odd in  $u$ , if  $W$  is odd in  $u$ . Since  $u + \lambda(-W(u)) = (1 - \lambda)u + \lambda B(u) \in M$  for  $u \in M \setminus K$  and  $0 \leq \lambda \leq 1$ , the Brezis-Martin theorem [22] implies that  $\sigma(t, M) \subset M$  for  $t \geq 0$ .

**Definition 2.2** (see [21, 23]). With the flow  $\sigma$ , a subset  $M \subset E$  is called an invariant set if  $\sigma(t, M) \subset M$  for  $t \geq 0$ .

Let us assume that

( $\Phi$ )  $K(J) \subset X$ ,  $J'(u) = u - A(u)$  for  $u \in E$ ,  $A : X \rightarrow X$  is continuous.

Under condition ( $\Phi$ ), we have  $\sigma(t, x) \in X$  for  $x \in X$  and  $\sigma$  is continuous in  $(t, x) \in \mathbb{R} \times X$ .

*Definition 2.3* (see [21]). Let  $M \subset X$  be an invariant set under  $\sigma$ .  $M$  is said to be an admissible invariant set for  $J$  if (a)  $M$  is the closure of an open set in  $X$ , that is,  $M = \dot{M} \cup \partial M$ ; (b) if  $u_n = \sigma(t_n, v)$  for some  $v \notin M$  and  $u_n \rightarrow u$  in  $E$  as  $t_n \rightarrow \infty$  for some  $u \in K$ , then  $u_n \rightarrow u$  in  $X$ ; (c) if  $u_n \in K \cap M$  such that  $u_n \rightarrow u$  in  $E$ , then  $u_n \rightarrow u$  in  $X$ ; (d) for any  $u \in \partial M \setminus K$ ,  $\sigma(t, u) \in \dot{M}$  for  $t > 0$ .

**Lemma 2.4** (see [24]). Let  $J \in C^1(E, \mathbb{R})$  and ( $\Phi$ ) hold. Assume  $J$  is even, bounded from below,  $J(0) = 0$  and satisfies (PS) condition. Assume that the positive cone  $P$  is an admissible invariant set for  $J$  and  $K_c \cap \partial P = \emptyset$  for all  $c < 0$ . Suppose there is a linear subspace  $F \subset X$  with  $\dim F = n$ , such that  $\sup_{F \cap \partial B_\rho} J(u) < 0$  for some  $\rho > 0$ , where  $B_\rho = \{u \in X : \|u\|_X \leq \rho\}$ . Then,  $J$  has at least  $n$  pairs of critical points with negative critical values. More precisely,

- (i) if  $\inf_P J \leq \inf_E J$ ,  $J$  has at least one pair of critical points in  $\dot{P} \cup (-\dot{P})$ , and at least  $n - 1$  pairs of critical points in  $X \setminus (P \cup (-P))$ ;
- (ii) if  $\inf_E J < \inf_P J$ ,  $J$  has at least one pair of critical points in  $\dot{P} \cup (-\dot{P})$ , and at least  $n$  pairs of critical points in  $X \setminus (P \cup (-P))$ .

**Lemma 2.5** (see [21]). Let  $J \in C^1(E, \mathbb{R})$  and ( $\Phi$ ) hold. Assume  $J$  is even,  $J(0) = 0$ , and  $J$  satisfies (PS) condition. Assume that the positive cone  $P$  is an admissible invariant set for  $J$  and  $K_c \cap \partial P = \emptyset$  for all  $c < 0$ . Suppose there exist linear subspaces  $F \subset X$  and  $H \subset E$  with  $\dim F = n$ ,  $\text{codim} H = k \geq 1$  ( $k = 0$ , resp.),  $n > k$ , such that for some  $\rho > 0$ ,  $\sup_{F \cap \partial B_\rho(0)} J(u) < 0$  and  $\inf_H J(u) > -\infty$ . Then,  $J$  has at least  $(n - k)$  ( $(n - 1)$ , resp.) pairs of critical points in  $X \setminus (P \cup (-P))$  with negative critical values.

**Lemma 2.6** (see [21]). Let  $J \in C^1(E, \mathbb{R})$  and ( $\Phi$ ) hold. Assume  $J$  is even,  $J(0) = 0$  and  $J$  satisfies (PS) condition. Assume that the positive cone  $P$  is an admissible invariant set for  $J$  and  $K_c \cap \partial P = \emptyset$  for all  $c > 0$ . Suppose there exist linear subspaces  $F \subset X$  and  $H \subset E$  with  $\dim F = n$ ,  $\text{codim} H = k \geq 1$ ,  $n > k + 1$ , such that for some  $\rho > \gamma > 0$ ,  $\sup_{F \cap \partial B_\rho(0)} J(u) \leq 0$  and  $\inf_{H \cap \partial B_\gamma(0)} J(u) > 0$ . Then for  $k \geq 1$  ( $k = 0$ , resp.),  $J$  has at least  $(n - k - 1)$  ( $(n - 1)$ , resp.) pairs of critical points in  $X \setminus (P \cup (-P))$  with positive critical values.

**Lemma 2.7** (see [21, 25]). Assume  $J \in C^1(E, \mathbb{R})$  is even,  $J(0) = 0$ , satisfies ( $\Phi$ ) and  $(PS)_c$  condition for  $c > 0$ . Assume that  $P$  is an admissible invariant set for  $J$ ,  $K_c \cap \partial P = \emptyset$  for all  $c > 0$ .  $E = \overline{\bigoplus_{j=1}^{\infty} E_j}$ , where  $E_j$  are finite-dimensional subspaces of  $X$ , and for each  $k$ , let  $Y_k = \bigoplus_{j=1}^k E_j$  and  $Z_k = \overline{\bigoplus_{j=k}^{\infty} E_j}$ . Assume for each  $k$  there exist  $\rho_k > \gamma_k > 0$  such that  $\lim_{k \rightarrow \infty} a_k < \infty$ , where  $a_k = \max_{Y_k \cap \partial B_{\rho_k}(0)} J(u)$ ,  $b_k = \inf_{Z_k \cap \partial B_{\gamma_k}(0)} J(u) \rightarrow \infty$  as  $k \rightarrow \infty$ . Then,  $J$  has a sequence of critical points  $u_n \in X \setminus (P \cup (-P))$  such that  $J(u_n) \rightarrow \infty$  as  $n \rightarrow \infty$ , provided  $Z_k \cap \partial B_{\gamma_k}(0) \cap P = \emptyset$  for large  $k$ .

Next, we need some basic concepts of ordered Banach spaces.

*Definition 2.8.* An ordered real Banach space is a pair  $(X, P)$ , where  $X$  is a real Banach space and  $P$  a closed convex subset of  $X$  such that  $(-P) \cap P = \{0\}$  and  $\mathbb{R}^+ \cdot P \subset P$ . The partial order

on  $X$  is given by the cone  $P$ . For  $u, v \in X$ , we write

$$\begin{aligned} u \leq v &\iff v - u \in P; \\ u < v &\iff u \leq v, \quad \text{but } u \neq v; \\ u \ll v &\iff v - u \in \dot{P}. \end{aligned} \tag{2.2}$$

If  $P$  has nonempty interior, then it is called a solid cone. If every ordered interval is bounded, then  $P$  is called a normal cone. An operator  $A : D(A) \rightarrow X$  is called order preserving (in the literature sometimes increasing) if

$$u \leq v \implies Au \leq Av; \tag{2.3}$$

strictly order preserving if

$$u < v \implies Au < Av; \tag{2.4}$$

and strongly order preserving if

$$u < v \implies Au \ll Av. \tag{2.5}$$

### 3. Main results

In this section, we will employ the abstract results in Section 2 to establish some existence theorems on sign-changing solutions of (BVP) (1.2). Firstly, we give some lemmas to change (BVP) (1.2) to a variational problem. Let  $C[0, 1]$  be the usual real Banach space with the norm  $\|u\|_C = \max_{t \in [0, 1]} |u(t)|$  for all  $u \in C[0, 1]$ . We can easily verify that

$$C_0[0, 1] = \{u \in C[0, 1] : u(0) = u(1) = 0\} \tag{3.1}$$

is also a Banach space with respect to  $\|\cdot\|_C$ . Let

$$P = \{u \in C_0[0, 1] : u(t) \geq 0 \forall t \in [0, 1]\}, \tag{3.2}$$

then  $P$  is a normal solid cone in  $C_0[0, 1]$  and

$$\dot{P} = \{u \in C_0[0, 1] : u(t) > 0 \forall t \in (0, 1)\}. \tag{3.3}$$

By  $L^2[0, 1]$ , we denote the usual real Hilbert space with the inner product  $(u, v) = \int_0^1 u(t)v(t)dt$  for all  $u, v \in L^2[0, 1]$ .

It is well known that the solution of (BVP) (1.2) in  $C^4[0, 1]$  is equivalent to the solution of the following integral equation in  $C[0, 1]$ :

$$u(t) = \int_0^1 G(t, s) \int_0^1 G(s, \tau) f(\tau, u(\tau)) d\tau ds, \quad t \in [0, 1], \tag{3.4}$$

where  $G(t, s)$  is the Green's function of the linear boundary value problem  $-u''(t) = 0$  for all  $t \in [0, 1]$  subject to  $u(0) = u(1) = 0$ , that is,

$$G(t, s) = \begin{cases} s(1-t), & 0 \leq s \leq t \leq 1; \\ t(1-s), & 0 \leq t \leq s \leq 1. \end{cases} \quad (3.5)$$

Define operators  $T, A_f : C[0, 1] \rightarrow C[0, 1]$  by

$$\begin{aligned} Tu(t) &= \int_0^1 G(t, s)u(s)ds, \\ A_f u(t) &= f(t, u(t)). \end{aligned} \quad (3.6)$$

Since  $T : C[0, 1] \rightarrow C_0[0, 1]$ , (3.4) is equivalent to the following operator equation in  $C_0[0, 1]$ :

$$u = T^2 A_f u. \quad (3.7)$$

*Remark 3.1.* It is easy to see that

- (i)  $G : [0, 1] \times [0, 1] \rightarrow [0, 1]$  is nonnegative continuous;
- (ii)  $\max_{(t,s) \in [0,1] \times [0,1]} G(t, s) = 1/4$ ;
- (iii)  $A_f : C[0, 1] \rightarrow C[0, 1]$  is bounded and continuous.

**Lemma 3.2** (see [20]).  $T : L^2[0, 1] \rightarrow C_0[0, 1]$  is a linear completely continuous operator and also a linear completely continuous operator from  $L^2[0, 1]$  to  $L^2[0, 1]$ . In addition,  $T : C_0[0, 1] \rightarrow C_0[0, 1]$  is strongly order-preserving.

From the definition of  $T$ , we can obtain that  $Tu \neq 0$  for all  $u \in L^2[0, 1]$  with  $u \neq 0$ . Therefore,  $Tu_1 \neq Tu_2$  for all  $u_1, u_2 \in L^2[0, 1]$  with  $u_1 \neq u_2$ . It is well known that all eigenvalues of  $T$  are

$$\{\lambda_k\}_{k \in \mathbb{N}} = \frac{1}{k^2 \pi^2}, \quad (3.8)$$

which have the corresponding orthonormal eigenfunctions

$$\{e_k\}_{k \in \mathbb{N}} = \{\sqrt{2} \sin k\pi t\}_{k \in \mathbb{N}}, \quad (3.9)$$

and  $\lambda_1 > \lambda_2 > \dots > \lambda_k > \dots > 0 \quad \forall k \in \mathbb{N}$ .

**Lemma 3.3** (see [10]). (i) *The operator equation*

$$u = T^2 A_f u \quad (3.10)$$

has a solution in  $C[0,1]$  if and only if the operator equation

$$v = TA_fTv \quad (3.11)$$

has a solution in  $L^2[0,1]$ .

(ii) The uniqueness of the solution for these two above equations is also equivalent.

*Remark 3.4.* From the proof of Lemma 3.3 [10], it is very clear if  $u \in L^2[0,1]$  is a solution for (3.11), then  $Tu \in C_0[0,1]$  is a solution for (3.7). Furthermore, if  $u \in C_0[0,1]$  is a solution for (3.11), then  $Tu \in C_0[0,1]$  is a solution for (3.7) with the same sign, which follows from Lemma 3.2.

**Lemma 3.5** (see [10]). Let  $\Psi(u) = \int_0^1 \int_0^{u(t)} f(t,v)dv dt$ ,  $u \in C[0,1]$ . Then,

- (i)  $\Psi$  is Fréchet differentiable on  $C[0,1]$  and  $(\Psi'(u))(w) = (A_f, w)$  for all  $u, w \in C[0,1]$ ;
- (ii)  $\Psi \circ T$  is Fréchet differentiable on  $L^2[0,1]$  and  $(\Psi \circ T)'(v) = TA_fTv$  for all  $v \in L^2[0,1]$ .

Choose  $E = L^2[0,1]$  and  $X = C_0[0,1]$  to be our Hilbert space and Banach space, respectively. Define a functional  $J : E \rightarrow R$ :

$$J(u) = \frac{1}{2}\|u\|^2 - \Psi(Tu), \quad u \in E. \quad (3.12)$$

Then, according to Lemma 3.5, we have

$$J'(u) = u - TA_fTu \quad \forall u \in E. \quad (3.13)$$

Hence, Lemma 3.3 implies that the operator equation  $u = T^2A_fu$  has a solution in  $X$  if and only if the functional  $J$  has a critical point in  $E$ . Thus, (BVP) (1.2) has been transformed into a variational problem.

We refer the following assumption:

( $f_1$ )  $f : [0,1] \times R \rightarrow R$  is continuous and increasing in  $u$ .

**Lemma 3.6.** Under ( $f_1$ ),  $(\Phi)$  is satisfied, and  $A := TA_fT : C_0[0,1] \rightarrow C_0[0,1]$  is strongly order-preserving.

*Proof.* The proof is similar to [20], and we omit it here. □

**Lemma 3.7.** Under ( $f_1$ ),  $M = P$  is an admissible invariant set for  $J$ .

*Proof.* We know that  $A : C_0[0,1] \rightarrow C_0[0,1]$  is strongly order-preserving, so does  $B$  given in Lemma 2.1. The Brezis-Martin theory implies that  $P$  and  $-P$  are invariant sets under the negative pseudogradient flow of  $J$ . Requirement (a) is satisfied automatically. For (d), we note that for all  $v \in P \setminus \{0\}$ , we have  $B(v) \in \dot{P}$ , similar to the proof in [23],  $\sigma(t, \partial P) \in \dot{P}$ . To prove (b), let  $u_n = \sigma(t_n, v)$  for some  $v \in X \setminus (P \cup (-P))$ , so  $u_n \in X = C_0[0,1]$ , let  $t_n \rightarrow \infty$  be a sequence such that  $u_n \rightarrow u$  in  $E = L^2[0,1]$  for some  $u \in K(J) \subset X = C_0[0,1]$ , then  $u_n \rightarrow u$  in  $X = C_0[0,1]$ . For (c), if  $u_n \in K(J) \cap (P \cup (-P)) \subset X$ , then  $J'(u_n) = 0$ , if  $u_n \rightarrow u$  in  $E = L^2[0,1]$ , for  $J \in C^1(E, R)$ , then  $J'(u) = 0$  and  $u \in K(J) \subset X = C_0[0,1]$ , so  $u_n \rightarrow u$  in  $X = C_0[0,1]$ , and the proof is completed. □

**Lemma 3.8** (see [15]). *Any bounded sequence  $\{u_n\} \subset L^2[0, 1]$  such that  $J'(u) \rightarrow 0$  as  $n \rightarrow \infty$  has a convergent subsequence.*

Next, we make more assumptions:

- (f<sub>2</sub>)  $\lim_{|u| \rightarrow \infty} f(t, u)/u < \pi^4$ , uniformly for  $t \in [0, 1]$ ;
- (f<sub>3</sub>)  $\lim_{|u| \rightarrow 0} f(t, u)/u > k^4 \pi^4$ , uniformly for  $t \in [0, 1]$  and some  $k \geq 2$ ;
- (f<sub>4</sub>)  $f(t, u)$  is odd in  $u$ .

**Theorem 3.9** (sublinear nonlinearity). *Under (f<sub>1</sub>)–(f<sub>4</sub>), (BVP) (1.2) has at least one pair of one-sign solutions  $u_1 > 0$ ,  $-u_1 < 0$ , and at least  $k - 1$  pairs of nodal solutions  $u_i$  for  $i = 2, \dots, k$ .*

*Proof.* It is easy to see that  $J \in C^1(E, R)$  and  $(\Phi)$  holds.  $P$  is an admissible invariant set for  $J$ , and  $K_c(J) \cap \partial P = \emptyset$  for  $c \neq 0$ . Also,  $J$  is even,  $J(0) = 0$ . By (f<sub>2</sub>), there exist  $\delta > 0$ ,  $\Lambda > 0$  such that  $F(t, u) \leq (1/2)(\pi^4 - \delta)u^2 + \Lambda$  for all  $u \in R$ , then

$$\begin{aligned}
 J(u) &= \frac{1}{2} \|u\|^2 - \int_0^1 F(t, Tu(t)) dt \\
 &\geq \frac{1}{2} \|u\|^2 - \frac{1}{2} (\pi^4 - \delta) \int_0^1 (Tu)^2 dt - \Lambda \\
 &\geq \frac{1}{2} \|u\|^2 - \frac{\pi^4 - \delta}{2\pi^4} \|u\|^2 - \Lambda \\
 &= \frac{\delta}{2\pi^4} \|u\|^2 - \Lambda \geq -\Lambda.
 \end{aligned} \tag{3.14}$$

So  $J$  is coercive, bounded from below, and satisfies (PS) condition.

Take  $F = \bigoplus_{i=1}^k \{e_i\}$ ; from (f<sub>3</sub>), there exist  $\eta > 0$ ,  $\delta_1 > 0$  such that  $|s| < \eta$ ,  $F(t, s) \geq (1/2)(k^4 \pi^4 + \delta_1)s^2$ , choose  $\rho = 4\eta$ , then  $|u| \leq \rho \Rightarrow |Tu| = |\int_0^1 G(t, s)u(s)ds| \leq (1/4)|u| \leq \eta$ , and

$$\begin{aligned}
 J(u) &= \frac{1}{2} \|u\|^2 - \int_0^1 F(t, Tu) dt \\
 &\leq \frac{1}{2} \|u\|^2 - \frac{1}{2} (k^4 \pi^4 + \delta_1) \int_0^1 (Tu)^2 dt \\
 &\leq \frac{1}{2} \|u\|^2 - \frac{k^4 \pi^4 + \delta_1}{2k^4 \pi^4} \|u\|^2 = -\frac{\delta_1}{2k^4 \pi^4} \|u\|^2 < 0,
 \end{aligned} \tag{3.15}$$

so  $\sup_{F \cap \partial B_\rho} J(u) < 0$  for  $\rho > 0$  small. Result follows from Lemma 2.4.  $\square$

Next, we consider an asymptotically linear problem:

- (f<sub>5</sub>)  $\lim_{|u| \rightarrow \infty} f(t, u)/u \in (k^4 \pi^4, (k+1)^4 \pi^4)$ , uniformly for  $t \in [0, 1]$ ;
- (f<sub>6</sub>)  $\lim_{|u| \rightarrow 0} f(t, u)/u \in (l^4 \pi^4, (l+1)^4 \pi^4)$ , uniformly for  $t \in [0, 1]$ .

**Theorem 3.10** (asymptotically linear case). *Under (f<sub>1</sub>), (f<sub>4</sub>), (f<sub>5</sub>), and (f<sub>6</sub>), (BVP) (1.2) has at least  $n$  pairs of nodal solutions provided  $k > l + 2$  or  $l > k + 1$ . Here,  $n = k - l - 2$ , if  $k > l + 2$ ; and  $n = l - k - 1$ , if  $l > k + 1$ .*

*Proof.* Take  $k^4\pi^4 < b_1 \leq b_2 < (k+1)^4\pi^4$  and  $\Lambda > 0$  such that for  $|u| \geq \Lambda$ ,  $b_1 \leq f(t, u)/u \leq b_2$ . Now let  $\{u_n\}$  be a (PS) sequence for  $J(u)$ . Writing  $u_n = v_n + w_n$  with  $v_n \in E_k = \bigoplus_{i=1}^k \{e_i\}$ ,  $w_n \in E_k^\perp$ , and taking inner product of  $J'(u_n)$  and  $v_n - w_n$ , we see that

$$\begin{aligned}
o(1) \cdot \|u_n\| &= \langle J'(u_n), v_n - w_n \rangle \\
&= \langle u_n, v_n - w_n \rangle - \int_0^1 f(t, Tu_n)(Tv_n - Tw_n) dt \\
&= \langle v_n + w_n, v_n - w_n \rangle - \int_{|Tu_n| \geq \Lambda} \frac{f(t, Tu_n)}{Tu_n} ((Tv_n)^2 - (Tw_n)^2) dt \\
&\quad - \int_{|Tu_n| < \Lambda} f(t, Tu_n)(Tv_n - Tw_n) dt \\
&\leq \|v_n\|^2 - \|w_n\|^2 - b_1 \int_{|Tu_n| \geq \Lambda} (Tv_n)^2 dt + b_2 \int_{|Tu_n| \geq \Lambda} (Tw_n)^2 dt \\
&\quad - \int_{|Tu_n| < \Lambda} f(t, Tu_n)(Tv_n - Tw_n) dt \\
&= \|v_n\|^2 - \|w_n\|^2 - b_1 \int_0^1 (Tv_n)^2 dt + b_1 \int_{|Tu_n| < \Lambda} (Tv_n)^2 + b_2 \int_0^1 (Tw_n)^2 dt \\
&\quad - b_2 \int_{|Tu_n| < \Lambda} (Tw_n)^2 dt - \int_{|Tu_n| < \Lambda} f(t, Tu_n)(Tv_n - Tw_n) dt \\
&\leq \|v_n\|^2 - \|w_n\|^2 - \frac{b_1}{k^4\pi^4} \|v_n\|^2 + \frac{b_2}{(k+1)^4\pi^4} \|w_n\|^2 + b_1 \int_{|Tu_n| < \Lambda} (Tv_n)^2 dt \\
&\quad - b_2 \int_{|Tu_n| < \Lambda} (Tw_n)^2 dt - \int_{|Tu_n| < \Lambda} f(t, Tu_n)(Tv_n - Tw_n) dt \\
&\leq \left(1 - \frac{b_1}{k^4\pi^4}\right) \|v_n\|^2 + \left(\frac{b_2}{(k+1)^4\pi^4} - 1\right) \|w_n\|^2 + \frac{b_1 b_2}{b_2 - b_1} \int_{|Tu_n| < \Lambda} |Tu_n|^2 dt \\
&\quad + \left(\int_{|Tu_n| \leq \Lambda} |f(t, Tu_n)|^2 dt\right)^{1/2} \left(\int_{|Tu_n| < \Lambda} (Tv_n - Tw_n)^2 dt\right)^{1/2} \\
&\leq \left(1 - \frac{b_1}{k^4\pi^4}\right) \|v_n\|^2 + \left(\frac{b_2}{(k+1)^4\pi^4} - 1\right) \|w_n\|^2 + \frac{b_1 b_2}{b_2 - b_1} \Lambda^2 + C \left(\int_0^1 |Tu_n|^2 dt\right)^{1/2} \\
&\leq \left(1 - \frac{b_1}{k^4\pi^4}\right) \|v_n\|^2 + \left(\frac{b_2}{(k+1)^4\pi^4} - 1\right) \|w_n\|^2 + \frac{b_1 b_2}{b_2 - b_1} \Lambda^2 + C \|u_n\| \\
&\leq -a \|u_n\|^2 + C \|u_n\| + C_1.
\end{aligned} \tag{3.16}$$

So  $\{u_n\}$  is bounded, where  $a = \min\{b_1/k^4\pi^4 - 1, 1 - b_2/(k+1)^4\pi^4\} > 0$ . Then,  $J(u)$  satisfies the (PS) condition.

If  $k > l + 2$ , let  $F = \bigoplus_{i=1}^k \{e_i\}$ , and  $H = \bigoplus_{i=l+2}^\infty \{e_i\}$ , then  $\dim F = k$ , and  $\text{codim} H = l + 1$ .



From  $(f_6)$ , we know that there exist  $\delta > 0$ , and  $\eta > 0$  such that

$$\frac{1}{2}(l^4\pi^4 + \delta)u^2 \leq F(t, u) \leq \frac{1}{2}((l+1)^4\pi^4 - \delta)u^2 \quad \text{for } |u| \leq \eta. \quad (3.17)$$

Then, for  $\|u\| \leq (1/4)\eta$ ,  $|Tu| \leq \eta$ , we can obtain, when  $u \in H$ ,

$$\begin{aligned} J(u) &= \frac{1}{2}\|u\|^2 - \frac{1}{2}\int_0^1 F(t, Tu)dt \\ &\geq \frac{1}{2}\|u\|^2 - \frac{1}{2}((l+1)^4\pi^4 - \delta)\int_0^1 (Tu)^2 dt \\ &\geq \frac{1}{2}\|u\|^2 - \frac{(l+1)^4\pi^4 - \delta}{2(l+2)^4\pi^4}\|u\|^2 \\ &= \left(\frac{1}{2} - \frac{(l+1)^4\pi^4 - \delta}{2(l+2)^4\pi^4}\right)\|u\|^2 > 0. \end{aligned} \quad (3.18)$$

So, choose  $\gamma = (1/4)\eta$ , then  $\inf_{H \cap \partial B_\gamma(0)} J(u) > 0$ .

From  $(f_5)$ , we can get there exist  $\theta > 0$ ,  $\Lambda > 0$  such that

$$\frac{1}{2}(k^4\pi^4 + \theta)u^2 - \Lambda \leq F(t, u) \leq \frac{1}{2}((k+1)^4\pi^4 - \theta)u^2 + \Lambda \quad \forall u \in \mathbb{R}. \quad (3.19)$$

Then, when  $u \in F$ , we have

$$\begin{aligned} J(u) &= \frac{1}{2}\|u\|^2 - \int_0^1 F(t, Tu)dt \\ &\leq \frac{1}{2}\|u\|^2 - \frac{1}{2}(k^4\pi^4 + \theta)\int_0^1 (Tu)^2 dt + \Lambda \\ &\leq \frac{1}{2}\|u\|^2 - \frac{k^4\pi^4 + \theta}{2k^4\pi^4}\|u\|^2 + \Lambda \\ &= -\frac{\theta}{2k^4\pi^4}\|u\|^2 + \Lambda. \end{aligned} \quad (3.20)$$

Choose  $\rho$  large enough such that  $\rho > \gamma > 0$ , and  $\sup_{F \cap \partial B_\rho(0)} J(u) \leq 0$ , result follows from Lemma 2.6.

If  $l > k + 1$ , let  $F = \bigoplus_{i=1}^l \{e_i\}$ ,  $H = \bigoplus_{i=k+2}^\infty \{e_i\}$ , then  $\dim F = l$ ,  $\text{codim} H = k + 1$ . From (3.17), when  $u \in F$ ,

$$\begin{aligned} J(u) &= \frac{1}{2}\|u\|^2 - \int_0^1 F(t, Tu)dt \\ &\leq \frac{1}{2}\|u\|^2 - \frac{1}{2}(l^4\pi^4 + \delta)\int_0^1 (Tu)^2 dt \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{2}\|u\|^2 - \frac{l^4\pi^4 + \delta}{2l^4\pi^4}\|u\|^2 \\
&= -\frac{\delta}{2l^4\pi^4}\|u\|^2 < 0.
\end{aligned} \tag{3.21}$$

When  $u \in H$ , we know from (3.19),

$$\begin{aligned}
J(u) &= \frac{1}{2}\|u\|^2 - \int_0^1 F(t, Tu) dt \\
&\geq \frac{1}{2}\|u\|^2 - \frac{1}{2}((k+1)^4\pi^4 - \theta) \int_0^1 (Tu)^2 dt - \Lambda \\
&\geq \frac{1}{2}\|u\|^2 - \frac{(k+1)^4\pi^4 - \theta}{2(k+2)^4\pi^4}\|u\|^2 - \Lambda \\
&= \left(\frac{1}{2} - \frac{(k+1)^4\pi^4 - \theta}{2(k+2)^4\pi^4}\right)\|u\|^2 - \Lambda \geq -\Lambda,
\end{aligned} \tag{3.22}$$

which means  $\inf_H J(u) > -\infty$ , then result follows from Lemma 2.5.  $\square$

Next, we consider a superlinear problem. Assume that

(f<sub>7</sub>) there is  $\mu > 2$  such that  $0 < \mu F(t, u) \leq f(t, u)u$  for  $|u|$  large;

(f<sub>8</sub>) there are  $p > \mu$ ,  $C > 0$  such that  $F(t, u) \leq C|u|^p$  for  $|u|$  large.

**Theorem 3.11** (superlinear nonlinearity). *Under (f<sub>1</sub>), (f<sub>4</sub>), (f<sub>7</sub>), and (f<sub>8</sub>), (BVP) (1.2) has infinitely many nodal solutions.*

*Proof.* From condition (f<sub>7</sub>) by the standard argument,  $J$  satisfies (PS)<sub>c</sub> condition for every  $c \in \mathbb{R}$ . Let  $Z_k = \bigoplus_{i=k}^{\infty} \{e_i\}$ . From (f<sub>8</sub>), we obtain  $|F(t, u)| \leq C|u|^p + C_1$  for all  $u \in \mathbb{R}$ . Define  $\beta_k = \sup_{u \in Z_k, \|Tu\|=1} |Tu|$ , it is very clear  $\beta_k < \infty$  and  $0 < \beta_{k+1} \leq \beta_k$ , so  $\beta_k \rightarrow \beta \geq 0$  and  $\beta \neq \pm\infty$ . So if  $u \in Z_k$ ,

$$\begin{aligned}
J(u) &= \frac{1}{2}\|u\|^2 - \int_0^1 F(t, Tu) dt \\
&\geq \frac{1}{2}\|u\|^2 - \int_0^1 (C|Tu|^p + C_1) dt \\
&\geq \frac{1}{2}\|u\|^2 - C\beta_k^p \|Tu\|^p - C_1 \\
&\geq \frac{1}{2}\|u\|^2 - C \frac{\beta_k^p}{(k^2\pi^2)^p} \|u\|^p - C_1.
\end{aligned} \tag{3.23}$$

Choosing  $r_k = (4C\beta_k^p)^{1/(2-p)} (k^2\pi^2)^{p/(p-2)}$ , we obtain, if  $u \in Z_k$  and  $\|u\| = r_k$ ,

$$J(u) \geq \frac{(k^2\pi^2)^{2p/(p-2)}}{4(4C\beta_k^p)^{2/(p-2)}} - C_1 \rightarrow \infty, \quad \text{if } k \rightarrow \infty. \tag{3.24}$$

Let  $Y_k = \bigoplus_{i=1}^k \{e_i\}$ . From (f7), after integrating, we obtain the existence of  $C_2 > 0$  such that  $F(t, u) \geq C_2|u|^\mu$  for  $|u| \geq R$ . Hence, we have  $F(t, u) \geq C_2|u|^\mu - C_3$  for  $u \in R$  and  $C_3 > 0$  is constant. Therefore, when  $u \in Y_k$ ,

$$\begin{aligned} J(u) &= \frac{1}{2} \|u\|^2 - \int_0^1 F(t, Tu) dt \\ &\leq \frac{1}{2} \|u\|^2 - \int_0^1 (C_2|Tu|^\mu - C_3) dt \\ &\leq \frac{1}{2} \|u\|^2 - C \|Tu\|^\mu + C_3 \\ &\leq \frac{1}{2} \|u\|^2 - \frac{C}{(k^2\pi^2)^\mu} \|u\|^\mu + C_3. \end{aligned} \tag{3.25}$$

Noting  $\mu > 2$ , choose  $\rho_k > r_k > 0$  large enough, such that  $J(u)|_{u \in Y_k \cap \partial B_{\rho_k}} < 0$ , and

$$\lim_{k \rightarrow \infty} \max_{u \in Y_k \cap \partial B_{\rho_k}} J(u) < \infty. \tag{3.26}$$

Result follows from Lemma 2.7. □

*Remark 3.12.* If there exist no strict subsolution and supersolution required in [20], just only using the functional  $J$  to get the critical point [10, 11], then we just know that (BVP) (1.2) has solutions, even we can know the sign of the critical point of the functional  $J$  because  $Tu$  is not strongly order-preserving in  $L^2[0, 1]$ . In our paper, using admissible invariant sets in  $C_0[0, 1]$ , we can settle the problem.

## Acknowledgments

The authors are grateful to the referees for their useful suggestions which have improved the writing of the paper. Jihui Zhang thanks Z. Zhang and the members of AMSS very much for their hospitality and invitation to visit the Academy of Mathematics and Systems Sciences (AMSS), Academia Sinica, in January 2008. The authors also would like to thank Professor D. Cao, Professor S. Li, Professor Y. Ding, and Professor H. Yin for their help and many valuable discussions. This research was supported by the NNSF of China (Grant no.10871096), Foundation of Major Project of Science and Technology of Chinese Education Ministry, SRFDP of Higher Education, and NSF of Education Committee of Jiangsu Province. Zhitao Zhang was supported by NNSF of China (Grant no.10671195).

## References

- [1] J. M. Davis, P. W. Eloe, and J. Henderson, "Triple positive solutions and dependence on higher order derivatives," *Journal of Mathematical Analysis and Applications*, vol. 237, no. 2, pp. 710–720, 1999.
- [2] J. M. Davis, J. Henderson, and P. J. Y. Wong, "General Lidstone problems: multiplicity and symmetry of solutions," *Journal of Mathematical Analysis and Applications*, vol. 251, no. 2, pp. 527–548, 2000.
- [3] Z. Bai and H. Wang, "On positive solutions of some nonlinear fourth-order beam equations," *Journal of Mathematical Analysis and Applications*, vol. 270, no. 2, pp. 357–368, 2002.

- [4] J. R. Graef, C. Qian, and B. Yang, "Multiple symmetric positive solutions of a class of boundary value problems for higher order ordinary differential equations," *Proceedings of the American Mathematical Society*, vol. 131, no. 2, pp. 577–585, 2003.
- [5] Y. Li, "Positive solutions of fourth-order periodic boundary value problems," *Nonlinear Analysis: Theory, Methods Applications*, vol. 54, no. 6, pp. 1069–1078, 2003.
- [6] Q. Yao, "Positive solutions for eigenvalue problems of fourth-order elastic beam equations," *Applied Mathematics Letters*, vol. 17, no. 2, pp. 237–243, 2004.
- [7] M. Ruyun, Z. Jihui, and F. Shengmao, "The method of lower and upper solutions for fourth-order two-point boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 215, no. 2, pp. 415–422, 1997.
- [8] Z. Bai, "The method of lower and upper solutions for a bending of an elastic beam equation," *Journal of Mathematical Analysis and Applications*, vol. 248, no. 1, pp. 195–202, 2000.
- [9] S. Charkrit and A. Kananthai, "Existence of solutions for some higher order boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 2, pp. 830–850, 2007.
- [10] F. Li, Q. Zhang, and Z. Liang, "Existence and multiplicity of solutions of a kind of fourth-order boundary value problem," *Nonlinear Analysis: Theory, Methods Applications*, vol. 62, no. 5, pp. 803–816, 2005.
- [11] X.-L. Liu and W.-T. Li, "Existence and multiplicity of solutions for fourth-order boundary value problems with parameters," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 1, pp. 362–375, 2007.
- [12] F. Li, Y. Li, and Z. Liang, "Existence of solutions to nonlinear Hammerstein integral equations and applications," *Journal of Mathematical Analysis and Applications*, vol. 323, no. 1, pp. 209–227, 2006.
- [13] F. Li, Y. Li, and Z. Liang, "Existence and multiplicity of solutions to  $2m$ th-order ordinary differential equations," *Journal of Mathematical Analysis and Applications*, vol. 331, no. 2, pp. 958–977, 2007.
- [14] G. Han and Z. Xu, "Multiple solutions of some nonlinear fourth-order beam equations," *Nonlinear Analysis: Theory, Methods Applications*, vol. 68, no. 12, pp. 3646–3656, 2008.
- [15] Y. Yang and J. Zhang, "Existence of solutions for some fourth-order boundary value problems with parameters," *Nonlinear Analysis: Theory, Methods Applications*, vol. 69, no. 4, pp. 1364–1375, 2008.
- [16] R. P. Agarwal and I. Kiguradze, "Two-point boundary value problems for higher-order linear differential equations with strong singularities," *Boundary Value Problems*, vol. 2006, Article ID 83910, 32 pages, 2006.
- [17] K. Perera and Z. Zhang, "Multiple positive solutions of singular  $p$ -Laplacian problems by variational methods," *Boundary Value Problems*, vol. 2005, no. 3, pp. 377–382, 2005.
- [18] D. Cao and E. S. Noussair, "Multiplicity of positive and nodal solutions for nonlinear elliptic problems in  $R_N$ ," *Annales de l'Institut Henri Poincaré. Analyse Non Linéaire*, vol. 13, no. 5, pp. 567–588, 1996.
- [19] D. Cao, E. S. Noussair, and S. Yan, "Solutions with multiple peaks for nonlinear elliptic equations," *Proceedings of the Royal Society of Edinburgh. Section A*, vol. 129, no. 2, pp. 235–264, 1999.
- [20] G. Han and F. Li, "Multiple solutions of some fourth-order boundary value problems," *Nonlinear Analysis: Theory, Methods Applications*, vol. 66, no. 11, pp. 2591–2603, 2007.
- [21] A. Qian and S. Li, "Multiple nodal solutions for elliptic equations," *Nonlinear Analysis: Theory, Methods Applications*, vol. 57, no. 4, pp. 615–632, 2004.
- [22] K. C. Chang, *Infinite-Dimensional Morse Theory and Multiple Solution Problems*, Progress in Nonlinear Differential Equations and Their Applications, Birkhäuser, Boston, Mass, USA, 1993.
- [23] Z. Liu and J. Sun, "Invariant sets of descending flow in critical point theory with applications to nonlinear differential equations," *Journal of Differential Equations*, vol. 172, no. 2, pp. 257–299, 2001.
- [24] S. Li and Z.-Q. Wang, "Ljusternik-Schnirelman theory in partially ordered Hilbert spaces," *Transactions of the American Mathematical Society*, vol. 354, no. 8, pp. 3207–3227, 2002.
- [25] P. H. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, vol. 65 of CBMS Regional Conference Series in Mathematics, American Mathematical Society, Washington, DC, USA, 1986.