PERMANENCE AND GLOBAL STABILITY OF POSITIVE SOLUTIONS OF A NONAUTONOMOUS DISCRETE RATIO-DEPENDENT PREDATOR-PREY MODEL

HAI-FENG HUO AND WAN-TONG LI

Received 28 October 2004

We first give sufficient conditions for the permanence of nonautonomous discrete ratio-dependent predator-prey model. By linearization of the model at positive solutions and construction of Lyapunov function, we also obtain some conditions which ensure that a positive solution of the model is stable and attracts all positive solutions.

1. Introduction

In the theoretical ecology, permanence and global stability of the population model are very important. There are extensive literature related to these topics for differential equation models (see [3, 6, 7, 8, 9, 12] and the references cited therein). Recently, there has been a tendency for some researchers in the field of difference equations to develop some new methods which are analogous to those used in the study of differential equations. (See, e.g., [1, 2, 4, 5, 10, 11] and the references therein.)

In [5], Fan and Wang considered the following discrete periodic ratio-dependent predator-prey model:

$$\begin{aligned} x_1(k+1) &= x_1(k)\exp\left\{a(k)-b(k)x_1(k)-\frac{c(k)x_2(k)}{m(k)x_2(k)+x_1(k)}\right\},\\ x_2(k+1) &= x_2(k)\exp\left\{-d(k)+\frac{f(k)x_1(k)}{m(k)x_2(k)+x_1(k)}\right\}, \end{aligned} \tag{1.1}$$

and establish sufficient conditions for the existence of a positive periodic solution of the periodic system (1.1). In this paper, we will establish sufficient conditions for the permanence of system (1.1) and also obtain some conditions which ensure that a positive solution of the model is stable and attracts all positive solutions.

First, we present two definitions.

Definition 1.1. System (1.1) is defined to be permanent if there are positive constants M and m such that each positive solution $\{x_1(k), x_2(k)\}$ of system (1.1) satisfies

$$m \le \liminf_{k\to\infty} x_i(k) \le \limsup_{k\to\infty} x_i(k) \le M, \quad i = 1,2. \tag{1.2}$$

Copyright © 2005 Hindawi Publishing Corporation
Discrete Dynamics in Nature and Society 2005:2 (2005) 135–144
DOI: 10.1155/DDNS.2005.135

Definition 1.2. System (1.1) is defined to be globally asymptotically stable if a positive solution of system (1.1) is stable and this solution attracts all positive solutions.

Throughout this paper, we will assume that $a(k)$, $b(k)$, $c(k)$, $d(k)$, $m(k)$, and $f(k)$ are bounded nonnegative sequences, and use the following notations: for any bounded sequence $\{u(k)\}$,

$$u^M = \sup_{k\in\mathbb{N}} u(k), \qquad u^L = \inf_{k\in\mathbb{N}} u(k). \tag{1.3}$$

For biological reasons, we only consider solution $\{x_1(k), x_2(k)\}$, with $x_1(0) > 0$, $x_2(0) > 0$.

The organization of this paper is the following. In the next section, we establish the permanence of system (1.1). In Section 3, we obtain the sufficient conditions which ensure that a positive solution of system (1.1) is stable and attracts all positive solutions.

2. Permanence

In this section, we establish a permanence result for system (1.1).

Lemma 2.1. *For every solution $\{x_1(k), x_2(k)\}$ of (1.1),*

$$\limsup_{k\to\infty} x_1(k) \le B_1, \qquad \limsup_{k\to\infty} x_2(k) \le B_2, \tag{2.1}$$

where

$$B_1 = \max\left\{\frac{a^M}{b^L}, \frac{\exp(a^M - 1)}{b^L}\right\}, \qquad B_2 = \left\{\frac{f^M B_1}{m^L d^L}, \frac{f^M B_1}{m^L d^L}\exp\{-d^L + f^M\}\right\}. \tag{2.2}$$

Proof. Clearly, $x_1(k) > 0$ and $x_2(k) > 0$ for $k \ge 0$. We first prove that

$$\limsup_{k\to\infty} x_1(k) \le B_1. \tag{2.3}$$

To prove (2.3), we first assume that there exists an $l_0 \in \mathbb{N}$ such that $x_1(l_0 + 1) \ge x_1(l_0)$. Then,

$$a(l_0) - b(l_0)x_1(l_0) - \frac{c(l_0)x_2(l_0)}{m(l_0)x_2(l_0) + x_1(l_0)} \ge 0. \tag{2.4}$$

Hence,

$$x_1(l_0) \le \frac{a(l_0)}{b(l_0)} \le \frac{a^M}{b^L}. \tag{2.5}$$

It follows that

$$\begin{aligned} x_1(l_0+1) &= x_1(l_0)\exp\left\{a(l_0) - b(l_0)x_1(l_0) - \frac{c(l_0)x_2(l_0)}{m(l_0)x_2(l_0) + x_1(l_0)}\right\} \\ &\le x_1(l_0)\exp\{a(l_0) - b(l_0)x_1(l_0)\} \\ &\le x_1(l_0)\exp\{a^M - b^L x_1(l_0)\} \le \frac{\exp(a^M - 1)}{b^L}, \end{aligned} \tag{2.6}$$

here we used

$$\max_{x\in\mathbb{R}}\{x\exp(a-bx)\}=\frac{\exp(a-1)}{b},\quad \text{for } a,b>0. \tag{2.7}$$

We claim that

$$x_1(k)\le B_1,\quad \text{for } k\ge l_0. \tag{2.8}$$

By way of contradiction, assume that there exists a $p_0>l_0$ such that $x_1(p_0)>B_1$. Then $p_0\ge l_0+2$. Let $\widetilde{p}_0\ge l_0+2$ be the smallest integer such that $x_1(\widetilde{p}_0)>B_1$. Then $x_1(\widetilde{p}_{0-1})<x_1(\widetilde{p}_0)$. The above argument produces that $x_1(\widetilde{p}_0)\le B_1$, a contradiction. This proves the claim. Now, we assume that $x_1(k+1)<x_1(k)$ for all $k\in\mathbb{N}$. In particular, $\lim_{k\to\infty}x_1(k)$ exists, denoted by $\overline{x}_1$. We claim that $\overline{x}_1\le a^M/b^L$. By way of contradiction, assume that $\overline{x}_1>a^M/b^L$. Taking limit in the first equation in system (1.1) gives

$$\lim_{k\to\infty}\left(a(k)-b(k)x_1(k)-\frac{c(k)x_2(k)}{m(k)x_2(k)+x_1(k)}\right)=0, \tag{2.9}$$

which is a contradiction since

$$a(k)-b(k)x_1(k)-\frac{c(k)x_2(k)}{m(k)x_2(k)+x_1(k)}\le a(k)-b(k)x_1(k)\le a^M-b^L\overline{x}_1<0,\quad \text{for } n\in\mathbb{N}. \tag{2.10}$$

This proves the claim. Note that $a^M/b^L\le B_1$. It follows that (2.3) holds.

Next, we prove that

$$\limsup_{k\to\infty}x_2(k)\le B_2. \tag{2.11}$$

At first, we assume that there exists an $n_0\in\mathbb{N}$ such that $x_2(n_0+1)\ge x_2(n_0)$. Then

$$-d(n_0)+\frac{f(n_0)x_1(n_0)}{m(n_0)x_2(n_0)+x_1(n_0)}\ge 0. \tag{2.12}$$

Hence,

$$\begin{aligned}-d(n_0)+\frac{f(n_0)x_1(n_0)}{m(n_0)x_2(n_0)}&\ge 0,\\ x_2(n_0)\le\frac{f(n_0)x_1(n_0)}{m(n_0)d(n_0)}&\le\frac{f^MB_1}{m^Ld^L}.\end{aligned} \tag{2.13}$$

It follows that

$$\begin{aligned}x_2(n_0+1)&=x_2(n_0)\exp\left\{-d(n_0)+\frac{f(n_0)x_1(n_0)}{m(n_0)x_2(n_0)+x_1(n_0)}\right\}\\ &\le\frac{f^MB_1}{m^Ld^L}\exp\{-d^L+f^M\}.\end{aligned} \tag{2.14}$$

We claim that $x_2(k) \le B_2$ for $k \ge n_0$. By way of contradiction, assume that there exists a $q_0 > n_0$ such that $x_2(q_0) > B_2$. Then $q_0 = n_0 + 2$. Let $\widetilde{q}_0 \ge n_0 + 2$ be the smallest integer such that $x_2(\widetilde{q}_0) > B_2$. Then $x_2(\widetilde{q}_{0-1}) < x_2(\widetilde{q}_0)$. The above argument produces that $x_2(\widetilde{q}_0) \le B_2$, a contradiction. This proves the claim. Now, we assume that $x_2(k+1) < x_2(k)$ for all $k \in \mathbb{N}$. In particular, $\lim_{k\to\infty} x_2(k)$ exists, denoted by $\overline{x}_2$. We claim that $\overline{x}_2 \le f^M B_1/m^L d^L$. By way of contradiction, assume that $\overline{x}_2 > f^M B_1/m^L d^L$. Taking limit in the second equation in system (1.1) gives

$$\lim_{k\to\infty}\left(-d(k)+\frac{f(k)x_1(k)}{m(k)x_2(k)+x_1(k)}\right)=0, \tag{2.15}$$

which is a contradiction since

$$-d(k)+\frac{f(k)x_1(k)}{m(k)x_2(k)+x_1(k)} \le -d^L+\frac{f^M B_1}{m^L \overline{x}_2} < 0. \tag{2.16}$$

It follows that (2.11) holds. This completes the proof. □

Lemma 2.2. *Assume that*

$$a^L > \frac{c^M}{m^L}, \qquad f^L > d^M. \tag{2.17}$$

Then

$$\liminf_{k\to\infty} x_1(k) \ge D_1, \qquad \liminf_{k\to\infty} x_2(k) \ge D_2, \tag{2.18}$$

where

$$\begin{aligned} D_1 &= \min\left\{\frac{a^L - c^M/m^L}{b^M}\exp\left\{a^L - b^M B_1 - \frac{c^M}{m^L}\right\}, \frac{a^L - c^M/m^L}{b^M}\right\},\\ D_2 &= \min\left\{\frac{f^L - d^M}{m^M d^M}D_1, \frac{f^L - d^M}{m^M d^M}\exp\left\{-d^M + \frac{f^L D_1}{m^M B_2 + D_1}\right\}\right\}. \end{aligned} \tag{2.19}$$

Proof. We first show that

$$\liminf_{k\to\infty} x_1(k) \ge D_1. \tag{2.20}$$

According to Lemma 2.1, there exists a $k^* \in \mathbb{N}$ such that

$$x_1(k) \le B_1 + \epsilon, \quad x_2(k) \le B_2 + \epsilon, \quad \text{for } k \ge k^*. \tag{2.21}$$

Firstly, we assume that there exists an $l_0 \geq k^*$ such that $x_1(l_0+1) \leq x_1(l_0)$. Note that, for $k \geq l_0$,

$$\begin{aligned} x_1(k+1) &= x_1(k)\exp\left\{a(k) - b(k)x_1(k) - \frac{c(k)x_2(k)}{m(k)x_2(k)+x_1(k)}\right\} \\ &\geq x_1(k)\exp\left\{a(k) - b(k)x_1(k) - \frac{c(k)}{m(k)}\right\} \\ &\geq x_1(k)\exp\left\{a^L - b^M x_1(k) - \frac{c^M}{m^L}\right\}. \end{aligned} \tag{2.22}$$

In particular, with $k = l_0$, we have

$$a^L - b^M x_1(l_0) - \frac{c^M}{m^L} \leq 0, \tag{2.23}$$

which implies that

$$x_1(l_0) \geq \frac{a^L - c^M/m^L}{b^M}. \tag{2.24}$$

Then

$$x_1(l_0+1) \geq \frac{a^L - c^M/m^L}{b^M}\exp\left\{a^L - b^M(B_1+\epsilon) - \frac{c^M}{m^L}\right\}. \tag{2.25}$$

Let

$$x_{1\epsilon} = \frac{a^L - c^M/m^L}{b^M}\exp\left\{a^L - b^M(B_1+\epsilon) - \frac{c^M}{m^L}\right\}. \tag{2.26}$$

We claim that

$$x_1(k) \leq x_{1\epsilon}, \quad \text{for } k \geq l_0. \tag{2.27}$$

By way of contradiction, assume that there exists a $p_0 \geq l_0$ such that $x_1(p_0) < x_{1\epsilon}$. Then $p_0 \geq l_0 + 2$. Let $\widetilde{p}_0 \geq l_0 + 2$ be the smallest integer such that $x_1(\widetilde{p}_0) < x_{1\epsilon}$. Then $x_1(\widetilde{p}_0 - 1) > x_1(\widetilde{p}_0)$. The above argument produces that $x_1(\widetilde{p}_0) \geq x_{1\epsilon}$, a contradiction. This proves the claim. Now, we assume that $x_1(k+1) > x_1(k)$ for all $k \in \mathbb{N}$. In particular, $\lim_{k\to\infty} x_1(k)$ exists, denoted by $\underline{x}_1$. We claim that

$$\underline{x}_1 \geq \frac{(a^L - c^M/m^L)}{b^M}. \tag{2.28}$$

By way of contradiction, assume that

$$\underline{x}_1 < \frac{(a^L - c^M/m^L)}{b^M}. \tag{2.29}$$

Taking limit in the first equation in system (1.1) gives

$$\lim_{k\to\infty}\left(a(k)-b(k)x_1(k)-\frac{c(k)x_2(k)}{m(k)x_2(k)+x_1(k)}\right)=0, \tag{2.30}$$

which is a contradiction since

$$\lim_{k\to\infty}\left(a(k)-b(k)x_1(k)-\frac{c(k)x_2(k)}{m(k)x_2(k)+x_1(k)}\right)\geq a^L-b^M\underline{x}_1-\frac{c^M}{m^L}>0. \tag{2.31}$$

This proves the claim. It follows that (2.20) holds.

Next, we prove that

$$\liminf_{k\to\infty} x_2(k)\geq D_2. \tag{2.32}$$

At first, we assume that there exists an $n_0\in\mathbb{N}$ such that $x_2(n_0+1)\geq x_2(n_0)$. Note that, for $k\geq n_0$,

$$\begin{aligned} x_2(k+1)&=x_2(k)\exp\left\{-d(k)+\frac{f(k)x_1(k)}{m(k)x_2(k)+x_1(k)}\right\}\\ &\geq x_2(k)\exp\left\{-d(k)+\frac{f(k)D_1}{m(k)x_2(k)+D_1}\right\}. \end{aligned} \tag{2.33}$$

In particular, with $k=n_0$, we get

$$-d(n_0)+\frac{f(n_0)D_1}{m(n_0)x_2(n_0)+D_1}\leq 0, \tag{2.34}$$

which implies that

$$x_2(n_0)\geq\frac{f^L-d^M}{m^Md^M}D_1. \tag{2.35}$$

Then

$$x_2(n_0+1)\geq\frac{f^L-d^M}{m^Md^M}D_1\exp\left\{-d^M+\frac{f^LD_1}{m^M(B_2+\epsilon)+D_1}\right\}. \tag{2.36}$$

Let

$$x_{2\epsilon}=\frac{f^L-d^M}{m^Md^M}D_1\exp\left\{-d^M+\frac{f^LD_1}{m^M(B_2+\epsilon)+D_1}\right\}. \tag{2.37}$$

We claim that $x_2(k)\geq x_{2\epsilon}$ for $k\geq n_0$. By way of contradiction, assume that there exists a $q_0\geq n_0$ such that $x_2(q_0)<x_{2\epsilon}$. Then $q_0\geq n_0+2$. Let $\widetilde{q}_0\geq n_0+2$ be the smallest integer such that $x_2(\widetilde{q}_0)<x_{2\epsilon}$. Then $x_2(\widetilde{q}_0-1)>x_2(\widetilde{q}_0)$. The above argument produces that $x_2(\widetilde{q}_0)\geq x_{2\epsilon}$, a contradiction. This proves the claim. Now, we assume that $x_2(k+1)<x_2(k)$ for all $k\in\mathbb{N}$. In particular, $\lim_{k\to\infty}x_2(k)$ exists, denoted by $\underline{x}_2$. We claim that

$$\underline{x}_2\geq\frac{f^L-d^M}{m^Md^M}D_1. \tag{2.38}$$

By way of contradiction, assume that

$$\underline{x}_2 < \frac{f^L - d^M}{m^M d^M} D_1. \tag{2.39}$$

Taking limit in the second equation in system (1.1) gives

$$\lim_{k\to\infty} \left(-d(k) + \frac{f(k)x_1(k)}{m(k)x_2(k) + x_1(k)} \right) = 0, \tag{2.40}$$

which is a contradiction since

$$-d(k) + \frac{f(k)x_1(k)}{m(k)x_2(k) + x_1(k)} \geq -d^M + \frac{f^L D_1}{m^M \underline{x}_2 + D_1} > 0. \tag{2.41}$$

It follows that (2.32) holds. This completes the proof. □

Now, by Lemmas 2.1 and 2.2, we can easily obtain the following result.

Theorem 2.3. *Assume that*

$$a^L > \frac{c^M}{m^L}, \qquad f^L > d^M. \tag{2.42}$$

Then system (1.1) is permanent.

3. Global stability

In this section, we derive sufficient conditions which guarantee that the positive solution of (1.1) is globally stable. Our strategy in the proof of the global stability of the positive solution of (1.1) is to construct suitable Lyapunov functions

Theorem 3.1. *In addition to the assumptions made in Theorem 2.3, assume further that*

(i) *there exist positive constant ν and positive constants n_i, $i = 1, 2$, such that*

$$\min\left\{ n_1 b(k) - n_1 \frac{c(k)}{4m(k)D_1} - n_2 \frac{f(k)}{4D_1}, n_2 \frac{f(k)D_1}{\left(m(k)B_2 + B_1\right)^2} - n_1 \frac{c(k)}{4m(k)D_2} \right\} > \nu, \tag{3.1}$$

for all large k, where D_i and B_i are given in Lemmas 2.1 and 2.2,

(ii) *$b(k)B_1 \leq 1$ and $f(k) \leq 4$ for all large k, where B_1 is given in Lemma 2.1.*

Then system (1.1) is globally asymptotically stable, that is, a positive solution of (1.1) is stable and attracts all positive solutions.

Proof. Let $\{x_1^*(k), x_2^*(k)\}$ be a positive solution of (1.1). We prove below that it is uniformly asymptotically stable. To this end, we introduce the change of variables

$$u_1(k) = x_1(k) - x_1^*(k), \qquad u_2(k) = x_2(k) - x_2^*(k). \tag{3.2}$$

System (1.1) is then transformed into

$$\begin{aligned}
u_1(k+1) &= x_1(k)\exp\left\{a(k)-b(k)x_1(k)-\frac{c(k)x_1(k)x_2(k)}{m(k)x_2(k)+x_1(k)}\right\}\\
&\quad -x_1^*(k)\exp\left\{a(k)-b(k)x_1^*(k)-\frac{c(k)x_1^*(k)x_2^*(k)}{a_1(k)x_2^*(k)+x_1^*(k)}\right\},\\
u_2(k+1) &= x_2(k)\exp\left\{\frac{f(k)x_1(k)}{m(k)x_2(k)+x_1(k)}-d(k)\right\}\\
&\quad -x_2^*(k)\exp\left\{\frac{f(k)x_1^*(k)}{a_1(k)x_2^*(k)+x_1^*(k)}-d(k)\right\}
\end{aligned} \tag{3.3}$$

which, by Taylor formula, can be rewritten as

$$\begin{aligned}
u_1(k+1) &= \exp\left\{a(k)-b(k)x_1^*(k)-\frac{c(k)x_1^*(k)x_2^*(k)}{m(k)x_2^*(k)+x_1^*(k)}\right\}\\
&\quad\times\left(\left(1-b(k)x_1^*(k)+\frac{c(k)x_1^*(k)x_2^*(k)}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}\right)u_1(k)\right.\\
&\quad\left.-\frac{c(k)x_1^*(k)}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}x_1^*(k)u_2(k)+f_1\left(k,u(k)\right)\right),\\
u_2(k+1) &= \exp\left\{\frac{f(k)x_1^*(k)}{m(k)x_2^*(k)+x_1^*(k)}-d(k)\right\}\\
&\quad\times\left(\left(1-\frac{f(k)m(k)x_1^*(k)x_2^*(k)}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}\right)u_2(k)\right.\\
&\quad\left.+\frac{f(k)m(k)\left(x_2^*(k)\right)^2}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}u_1(k)+f_2\left(k,u(k)\right)\right),
\end{aligned} \tag{3.4}$$

where $|f_i(k,u)|/\|u\|$ converges, uniformly with respect to $k\in\mathbb{N}$, to zero as $\|u\|\to 0$. In view of system (1.1), it follows from (3.4) that

$$\begin{aligned}
u_1(k+1) &= x_1^*(k+1)\left(\left(1-b(k)x_1^*(k)+\frac{c(k)x_1^*(k)x_2^*(k)}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}\right)\frac{u_1(k)}{x_1^*(k)}\right.\\
&\quad\left.-\frac{c(k)x_1^*(k)}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}u_2(k)+\frac{f_1\left(k,u(k)\right)}{x_1^*(k)}\right),\\
u_2(k+1) &= x_2^*(k+1)\left(\left(1-\frac{f(k)m(k)x_1^*(k)x_2^*(k)}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}\right)\frac{u_2(k)}{x_2^*(k)}\right.\\
&\quad\left.+\frac{f(k)m(k)x_2^*(k)}{\left(m(k)x_2^*(k)+x_1^*(k)\right)^2}u_1(k)+\frac{f_2\left(k,u(k)\right)}{x_2^*(k)}\right),
\end{aligned} \tag{3.5}$$

where $|f_i(k,u)|/\|u\|$ converges, uniformly with respect to $k \in \mathbb{N}$, to zero as $\|u\| \to 0$. We define the function V by

$$V(u(k)) = n_1 \left| \frac{u_1(k)}{x_1^*(k)} \right| + n_2 \left| \frac{u_2(k)}{x_2^*(k)} \right|, \tag{3.6}$$

where n_j are positive constants given in (i). Calculating the difference of V along the solution of system (3.5) and using (ii), we obtain

$$\begin{aligned} \Delta V \le & -\left(n_1 b(k) - n_1 \frac{c(k)}{4m(k)D_1} - n_2 \frac{f(k)}{4D_1} \right) x_1^*(k) \left| \frac{u_1(k)}{x_1^*(k)} \right| \\ & -\left(n_2 \frac{f(k)D_1}{\left(m(k)B_2 + B_1\right)^2} - n_1 \frac{c(k)}{4m(k)D_2} \right) x_2^*(k) \left| \frac{u_2(k)}{x_2^*(k)} \right| \\ & + n_1 \frac{|f_1(k,u(k))|}{x_1^*(k)} + n_2 \frac{|f_1(k,u(k))|}{x_2^*(k)}, \quad \text{for large } k. \end{aligned} \tag{3.7}$$

Since $|f_i(k,u)|/\|u\|$ converges uniformly to zero as $\|u\| \to 0$, it follows from condition (i) and Theorem 2.3 that there is a positive constant γ such that if k is sufficiently large and $\|u(k)\| < \gamma$,

$$\Delta V \le -\frac{\nu \|u(k)\|}{2}. \tag{3.8}$$

By [1], we see that the trivial solution of (3.5) is uniformly asymptotically stable, and so is the solution $\{x_1^*(k), x_2^*(k)\}$ of (1.1). Note that the positive solution $\{x_1(k), x_2(k)\}$ is chosen in an arbitrary way. Proceeding exactly as in [11], we conclude that the positive solution $\{x_1^*(k), x_2^*(k)\}$ of (1.1) is globally stable. The proof is complete □

Acknowledgments

The first author was supported by the NSF of Gansu Province of China, the NSF of Bureau of Education of Gansu Province of China (0416B-08), the Key Research and Development Program for Outstanding Groups of Lanzhou University of Technology, and the Development Program for Outstanding Young Teachers in Lanzhou University of Technology. The second author was supported by the NNSF of China (10171040), the NSF of Gansu Province of China (ZS011-A25-007-Z), and the Teaching and Research Award Program for Outstanding Young Teachers in Higher Education Institutions of the Ministry of Education of China.

References

[1] R. P. Agarwal, *Difference Equations and Inequalities. Theory, Methods, and Applications*, 2nd ed., Monographs and Textbooks in Pure and Applied Mathematics, vol. 228, Marcel Dekker, New York, 2000.

[2] R. P. Agarwal and P. J. Y. Wong, *Advanced Topics in Difference Equations*, Mathematics and Its Applications, vol. 404, Kluwer Academic, Dordrecht, 1997.

[3] L. S. Chen, Z. Y. Lu, and W. D. Wang, *The effect of delays on the permanence for Lotka-Volterra systems*, Appl. Math. Lett. **8** (1995), no. 4, 71–73.
[4] Y. M. Chen and Z. Zhou, *Stable periodic solution of a discrete periodic Lotka-Volterra competition system*, J. Math. Anal. Appl. **277** (2003), no. 1, 358–366.
[5] M. Fan and K. Wang, *Periodic solutions of a discrete time nonautonomous ratio-dependent predator-prey system*, Math. Comput. Modelling **35** (2002), no. 9-10, 951–961.
[6] H. I. Freedman and S. G. Ruan, *Uniform persistence in functional-differential equations*, J. Differential Equations **115** (1995), no. 1, 173–192.
[7] H.-F. Huo, W.-T. Li, and S. S. Cheng, *Periodic solutions of two-species diffusion models with continuous time delays*, Demonstratio Math. **35** (2002), no. 2, 433–446.
[8] Y. Kuang, *Delay Differential Equations with Applications in Population Dynamics*, Mathematics in Science and Engineering, vol. 191, Academic Press, Massachusetts, 1993.
[9] Z. Y. Lu and Y. Takeuchi, *Permanence and global attractivity for competitive Lotka-Volterra systems with delay*, Nonlinear Anal. **22** (1994), no. 7, 847–856.
[10] Y. Muroya, *Persistence and global stability for discrete models of nonautonomous Lotka-Volterra type*, J. Math. Anal. Appl. **273** (2002), no. 2, 492–511.
[11] W. Wendi and L. Zhengyi, *Global stability of discrete models of Lotka-Volterra type*, Nonlinear Anal. Ser. B: Real World Applications **35** (1999), no. 8, 1019–1030.
[12] W. Wendi and M. Zhien, *Harmless delays for uniform persistence*, J. Math. Anal. Appl. **158** (1991), no. 1, 256–268.

Hai-Feng Huo: Department of Applied Mathematics, Lanzhou University of Technology, Lanzhou, Gansu 730050, China
E-mail address: hfhuo@lut.cn

Wan-Tong Li: Department of Mathematics, Lanzhou University, Lanzhou, Gansu 730000, China
E-mail address: wtli@lzu.edu.cn

Mathematical Problems in Engineering

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at http://www.hindawi.com/journals/mpe/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/ according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru