

Research Article

Composition Operators from the Weighted Bergman Space to the n th Weighted Spaces on the Unit Disc

Stevo Stević

Mathematical Institute of the Serbian Academy of Sciences, Knez Mihailova 36/III, 11000 Beograd, Serbia

Correspondence should be addressed to Stevo Stević, sstevic@ptt.rs

Received 23 May 2009; Revised 27 August 2009; Accepted 4 September 2009

Recommended by Leonid Berezansky

The boundedness of the composition operator from the weighted Bergman space to the recently introduced by the author, the n th weighted space on the unit disc, is characterized. Moreover, the norm of the operator in terms of the inducing function and weights is estimated.

Copyright © 2009 Stevo Stević. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let \mathbb{D} be the open unit disc in the complex plane \mathbb{C} , $dm(z)$ the Lebesgue area measure on \mathbb{D} , $dm_\alpha(z) = (1 - |z|^2)^\alpha dm(z)$, $\alpha > -1$, and $H(\mathbb{D})$ the space of all analytic functions on the unit disc.

The weighted Bergman space $A_\alpha^p(\mathbb{D})$, where $p > 0$ and $\alpha > -1$, consists of all $f \in H(\mathbb{D})$ such that

$$\|f\|_{A_\alpha^p(\mathbb{D})}^p = (\alpha + 1) \int_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\alpha dm(z) < \infty. \quad (1.1)$$

With this norm, $A_\alpha^p(\mathbb{D})$ is a Banach space when $p \geq 1$, while for $p \in (0, 1)$, it is a Fréchet space with the translation invariant metric

$$d(f, g) = \|f - g\|_{A_\alpha^p(\mathbb{D})}^p, \quad f, g \in A_\alpha^p(\mathbb{D}). \quad (1.2)$$

Let $\mu(z)$ be a positive continuous function on a set $X \subset \mathbb{C}$ (*weight*) and $n \in \mathbb{N}_0$ be fixed. The n th *weighted-type space* on X , denoted by $\mathcal{W}_\mu^{(n)}(X)$, consists of all $f \in H(X)$ such that

$$b_{\mathcal{W}_\mu^{(n)}(X)}(f) := \sup_{z \in X} \mu(z) \left| f^{(n)}(z) \right| < \infty. \quad (1.3)$$

For $n = 0$, the space becomes the weighted-type space $H_\mu^\infty(X)$, for $n = 1$ the Bloch-type space $\mathcal{B}_\mu(X)$, and for $n = 2$ the Zygmund-type space $\mathcal{Z}_\mu(X)$.

For $n \in \mathbb{N}$, the quantity $b_{\mathcal{W}_\mu^{(n)}(X)}(f)$ is a seminorm on the n th weighted-type space $\mathcal{W}_\mu^{(n)}(X)$ and a norm on $\mathcal{W}_\mu^{(n)}(X)/\mathbb{P}_{n-1}$, where \mathbb{P}_{n-1} is the set of all polynomials whose degrees are less than or equal to $n - 1$. A natural norm on the n th weighted-type space can be introduced as follows:

$$\|f\|_{\mathcal{W}_\mu^{(n)}(X)} = \sum_{j=0}^{n-1} \left| f^{(j)}(a) \right| + b_{\mathcal{W}_\mu^{(n)}(X)}(f), \quad (1.4)$$

where a is an element in X . With this norm, the n th weighted-type space becomes a Banach space.

For $X = \mathbb{D}$ is obtained the space $\mathcal{W}_\mu^{(n)}(\mathbb{D})$, on which a norm is introduced as follows:

$$\|f\|_{\mathcal{W}_\mu^{(n)}(\mathbb{D})} := \sum_{j=0}^{n-1} \left| f^{(j)}(0) \right| + \sup_{z \in \mathbb{D}} \mu(z) \left| f^{(n)}(z) \right|. \quad (1.5)$$

Some information on Zygmund-type spaces on the unit disc and some operators on them can be found, for example, in [1–6], for the case of the upper half-plane, see [7, 8], while some information in the setting of the unit ball can be found, for example, in [9–13]. This considerable interest in Zygmund-type spaces motivated us to introduce the n th weighted-type space (see [8]).

Assume φ is a holomorphic self-map of \mathbb{D} . The composition operator induced by φ is defined on $H(\mathbb{D})$ by

$$(C_\varphi f)(z) = f(\varphi(z)). \quad (1.6)$$

A typical problem is to provide function theoretic characterizations when φ induces bounded or compact composition operators between two given spaces of holomorphic functions. Some classical results on composition and weighted composition operators can be found, for example, in [14], while some recent results can be found in [1, 5, 7, 15–34] (see also related references therein).

Here we characterize the boundedness of the composition operator from the weighted Bergman space to the n th weighted space on the unit disc when $n \in \mathbb{N}$. The case $n = 0$ was previously treated in [16, 22, 24, 31, 35]. Hence we will not consider this case here. See also [36] for some good results on weighted composition operators between weighted-type spaces. The case $n = 1$ was treated, for example, in [26, 32]. For some other results on weighted composition operators which map a space into a weighted or a Bloch-type space, see, for example, [15, 17–21, 23, 25, 33, 34].

Let X and Y be topological vector spaces whose topologies are given by translation-invariant metrics d_X and d_Y , respectively, and $T : X \rightarrow Y$ be a linear operator. It is said that T is *metrically bounded* if there exists a positive constant K such that

$$d_Y(Tf, 0) \leq Kd_X(f, 0) \quad (1.7)$$

for all $f \in X$. When X and Y are Banach spaces, the metrically boundedness coincides with the usual definition of bounded operators between Banach spaces.

If Y is a Banach space, then the quantity $\|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow Y}$ is defined as follows:

$$\|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow Y} := \sup_{\|f\|_{A_\alpha^p(\mathbb{D})} \leq 1} \|C_\varphi f\|_Y. \quad (1.8)$$

It is easy to see that this quantity is finite if and only if the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow Y$ is metrically bounded. For the case $p \geq 1$ this is the standard definition of the norm of the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow Y$, between two Banach spaces. If we say that an operator is bounded, it means that it is metrically bounded.

Throughout this paper, constants are denoted by C , they are positive and may differ from one occurrence to the other. The notation $a \asymp b$ means that there is a positive constant C such that $a \leq Cb$. Moreover, if both $a \asymp b$ and $b \asymp a$ hold, then one says that $a \approx b$.

2. Auxiliary Results

Here, we quote several auxiliary results. The first lemma is a direct consequence of a well-known estimate in [37, Proposition 1.4.10]. Hence, we omit its proof.

Lemma 2.1. *Assume $p > 0$, $\alpha > -1$, $n \in \mathbb{N}_0$, and $w \in \mathbb{D}$. Then the function*

$$g_{w,n}(z) = \frac{(1 - |w|^2)^{n+(\alpha+2)/p}}{(1 - \bar{w}z)^{n+2((\alpha+2)/p)}}, \quad (2.1)$$

belongs to $A_\alpha^p(\mathbb{D})$. Moreover, $\sup_{w \in \mathbb{D}} \|g_{w,n}\|_{A_\alpha^p} < \infty$.

The next lemma is folklore and was essentially proved in [38]. We will sketch a proof of it for the completeness and the benefit of the reader.

Lemma 2.2. *Assume $p > 0$, $\alpha > -1$, $n \in \mathbb{N}_0$, and $z \in \mathbb{D}$. Then there is a positive constant C independent of f such that*

$$|f^{(n)}(z)| \leq C \frac{\|f\|_{A_\alpha^p(\mathbb{D})}}{(1 - |z|)^{n+(\alpha+2)/p}}. \quad (2.2)$$

Proof. By the subharmonicity of the function $|f^{(n)}(z)|^p$, $p > 0$, applied on the disk:

$$D\left(z, \frac{1-|z|}{2}\right) = \left\{ z \in \mathbb{C} \mid |z-w| < \frac{1-|z|}{2} \right\}, \quad (2.3)$$

and since

$$1-|w| \asymp 1-|z|, \quad w \in D\left(z, \frac{1-|z|}{2}\right), \quad (2.4)$$

we have that

$$\left|f^{(n)}(z)\right|^p \leq \frac{C}{(1-|z|)^{2+\alpha+pn}} \int_{D(z, (1-|z|)/2)} \left|f^{(n)}(w)\right|^p dm_{\alpha+pn}(w). \quad (2.5)$$

From (2.5) and in light of the following well-known asymptotic relation [38]:

$$\int_{\mathbb{D}} |f(z)|^p (1-|z|^2)^\alpha dm(z) \asymp \sum_{j=0}^{n-1} |f^{(j)}(0)| + \int_{\mathbb{D}} |f^{(n)}(z)|^p (1-|z|^2)^{\alpha+np} dm(z), \quad (2.6)$$

the lemma easily follows. \square

Lemma 2.3. Assume $a > 0$ and

$$D_n(a) = \begin{vmatrix} 1 & 1 & \cdots & 1 \\ a & a+1 & \cdots & a+n-1 \\ a(a+1) & (a+1)(a+2) & \cdots & (a+n-1)(a+n) \\ & & \cdots & \\ \prod_{j=0}^{n-2} (a+j) & \prod_{j=0}^{n-2} (a+j+1) & \cdots & \prod_{j=0}^{n-2} (a+j+n-1) \end{vmatrix}. \quad (2.7)$$

Then $D_n = \prod_{j=1}^{n-1} j!$.

Proof. By using elementary transformations, we have

$$D_n(a) = \begin{vmatrix} 1 & 0 & \cdots & 0 \\ a & 1 & \cdots & 1 \\ a(a+1) & 2(a+1) & \cdots & 2(a+n-1) \\ & & \cdots & \\ \prod_{j=0}^{n-2} (a+j) & (n-1) \prod_{j=0}^{n-3} (a+j+1) & \cdots & (n-1) \prod_{j=0}^{n-3} (a+j+n-1) \end{vmatrix}, \quad (2.8)$$

from which it follows that

$$D_n(a) = (n-1)!D_{n-1}(a+1), \quad (2.9)$$

which along with the fact $D_2(a+n-2) = 1$ implies the lemma. \square

We will also need the classical Faà di Bruno's formula

$$(f \circ \varphi)^{(n)}(z) = \sum \frac{n!}{k_1! \cdots k_n!} f^{(k)}(\varphi(z)) \prod_{j=1}^n \left(\frac{\varphi^{(j)}(z)}{j!} \right)^{k_j}, \quad (2.10)$$

where $k = k_1 + k_2 + \cdots + k_n$ and the sum is over all nonnegative integers k_1, k_2, \dots, k_n satisfying $k_1 + 2k_2 + \cdots + nk_n = n$. For a nice exposition related to this formula see, for example, [39].

By using Bell polynomials $B_{n,k}(x_1, \dots, x_{n-k+1})$, (2.10) can be written in the following form:

$$(f \circ \varphi)^{(n)}(z) = \sum_{k=0}^n f^{(k)}(\varphi(z)) B_{n,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(n-k+1)}(z)). \quad (2.11)$$

Remark 2.4. Since $B_{n,0}(x_1, \dots, x_{n+1}) = 0$ the summation in (2.11) is from 1 to k . Moreover, since $B_{n,1}(x_1, \dots, x_n) = x_n$ and $B_{n,n}(x_1) = x_1^n$, (2.11) can be written in the following form:

$$\begin{aligned} (f \circ \varphi)^{(n)}(z) &= f'(\varphi(z))\varphi^{(n)}(z) + \sum_{k=2}^{n-1} f^{(k)}(\varphi(z)) B_{n,k}(\varphi'(z), \dots, \varphi^{(n-k+1)}(z)) \\ &\quad + f^{(n)}(\varphi(z))(\varphi'(z))^n. \end{aligned} \quad (2.12)$$

3. Main Result

Here, we formulate and prove our main result.

Theorem 3.1. *Assume $p > 0$, $\alpha > -1$, $n \in \mathbb{N}$, μ is a weight on \mathbb{D} and φ is a holomorphic self-map of \mathbb{D} . Then $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})$ is bounded if and only if*

$$I_k := \sup_{z \in \mathbb{D}} \frac{\mu(z) \left| \sum (n! / (k_1! \cdots k_n!)) \prod_{j=1}^n (\varphi^{(j)}(z) / j!)^{k_j} \right|}{(1 - |\varphi(z)|^2)^{k+(\alpha+2)/p}} < \infty, \quad k = 1, \dots, n, \quad (3.1)$$

where for each fixed $k \in \{1, \dots, n\}$, the sum is over all nonnegative integers k_1, k_2, \dots, k_n such that $k = k_1 + k_2 + \cdots + k_n$ and $k_1 + 2k_2 + \cdots + nk_n = n$.

Moreover, if the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D}) / \mathbb{P}_{n-1}$ is bounded, then

$$\|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D}) / \mathbb{P}_{n-1}} \asymp \sum_{k=1}^n I_k. \quad (3.2)$$

Remark 3.2. Note that by (2.11) we see that the conditions in (3.1) can be written in the following form:

$$I_k = \sup_{z \in \mathbb{D}} \frac{\mu(z) |B_{n,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(n-k+1)}(z))|}{(1 - |\varphi(z)|^2)^{k+(\alpha+2)/p}} < \infty, \quad k = 1, \dots, n. \quad (3.3)$$

Proof. First assume that conditions in (3.1) hold. By formula (2.10) and Lemma 2.2 we have

$$\begin{aligned} \|C_\varphi f\|_{\mathcal{W}_\mu^{(n)}(\mathbb{D})} &= \sum_{j=0}^{n-1} \left| (f \circ \varphi)^{(j)}(0) \right| + \sup_{z \in \mathbb{D}} \mu(z) \left| (C_\varphi f)^{(n)}(z) \right| \\ &= \sum_{j=0}^{n-1} \left| \sum_{l=0}^j \frac{j!}{l_1! \dots l_j!} f^{(l)}(\varphi(0)) \prod_{s=1}^j \left(\frac{\varphi^{(s)}(0)}{s!} \right)^{l_s} \right| \\ &\quad + \sup_{z \in \mathbb{D}} \mu(z) \left| \sum_{k=1}^n \frac{n!}{k_1! \dots k_n!} f^{(k)}(\varphi(z)) \prod_{j=1}^n \left(\frac{\varphi^{(j)}(z)}{j!} \right)^{k_j} \right| \\ &\leq \sum_{j=0}^{n-1} \sum_{l=0}^j \left| f^{(l)}(\varphi(0)) \right| \left| \sum_{s=1}^j \frac{j!}{l_1! \dots l_j!} \prod_{s=1}^j \left(\frac{\varphi^{(s)}(0)}{s!} \right)^{l_s} \right| \\ &\quad + C \|f\|_{A_\alpha^p(\mathbb{D})} \sum_{k=1}^n \sup_{z \in \mathbb{D}} \frac{\mu(z) \left| \sum (n! / (k_1! \dots k_n!)) \prod_{j=1}^n (\varphi^{(j)}(z) / j!)^{k_j} \right|}{(1 - |\varphi(z)|^2)^{k+(\alpha+2)/p}}. \end{aligned} \quad (3.4)$$

From this, (2.2) with $z = \varphi(0)$, and by conditions in (3.1), it follows that the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})$ is bounded. Moreover, if we consider the space $\mathcal{W}_\mu^{(n)}(\mathbb{D}) / \mathbb{P}_{n-1}$, we have that

$$\|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D}) / \mathbb{P}_{n-1}} \leq C \sum_{k=1}^n \sup_{z \in \mathbb{D}} \frac{\mu(z) \left| \sum (n! / (k_1! \dots k_n!)) \prod_{j=1}^n (\varphi^{(j)}(z) / j!)^{k_j} \right|}{(1 - |\varphi(z)|^2)^{k+(\alpha+2)/p}}. \quad (3.5)$$

Now assume that the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})$ is bounded. For a fixed $w \in \mathbb{D}$, and constants c_1, \dots, c_n , set

$$g_w(z) = \sum_{j=1}^n \frac{c_j}{n-2+j+2((\alpha+2)/p)} \frac{(1 - |w|^2)^{n-2+j+(\alpha+2)/p}}{(1 - \bar{w}z)^{n-2+j+2((\alpha+2)/p)}}. \quad (3.6)$$

Applying Lemma 2.1 we see that $g_w \in A_\alpha^p(\mathbb{D})$ for every $w \in \mathbb{D}$. Moreover, we have that

$$\sup_{w \in \mathbb{D}} \|g_w\|_{A_\alpha^p(\mathbb{D})} \leq C. \quad (3.7)$$

Now we show that for each $l \in \{1, \dots, n\}$, there are constants c_1, c_2, \dots, c_n , such that

$$g_w^{(l)}(w) = \frac{\bar{w}^l}{(1 - |w|^2)^{l+(\alpha+2)/p}}, \quad g_w^{(m)}(w) = 0, \quad m \in \{1, \dots, n\} \setminus \{l\}. \quad (3.8)$$

Indeed, by differentiating function g_w , for each $l \in \{1, \dots, n\}$, the system in (3.8) becomes

$$\begin{aligned} c_1 + c_2 + \dots + c_n &= 0, \\ \left(n + 2\frac{\alpha+2}{p}\right)c_1 + \left(n+1 + 2\frac{\alpha+2}{p}\right)c_2 + \dots + \left(2n-1 + 2\frac{\alpha+2}{p}\right)c_n &= 0, \\ &\vdots \\ \prod_{j=0}^{l-2} \left(n+j + 2\frac{\alpha+2}{p}\right)c_1 + \prod_{j=0}^{l-2} \left(n+1+j + 2\frac{\alpha+2}{p}\right)c_2 + \dots + \prod_{j=0}^{l-2} \left(2n-1+j + 2\frac{\alpha+2}{p}\right)c_n &= 1, \\ &\vdots \\ \prod_{j=0}^{n-2} \left(n+j + 2\frac{\alpha+2}{p}\right)c_1 + \prod_{j=0}^{n-2} \left(n+1+j + 2\frac{\alpha+2}{p}\right)c_2 + \dots + \prod_{j=0}^{n-2} \left(2n-1+j + 2\frac{\alpha+2}{p}\right)c_n &= 0. \end{aligned} \quad (3.9)$$

By using Lemma 2.3 with $a = n + 2(2 + \alpha)/p > 0$, we obtain that the determinant of system (3.9) is different from zero from which the claim follows.

Now for each $k \in \{1, \dots, n\}$, we choose the corresponding family of functions which satisfy (3.8) and denote it by $g_{w,k}$.

For each $k \in \{1, \dots, n\}$, the boundedness of the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})$ along with (2.10) and (3.7) implies that for each $\varphi(w) \neq 0$:

$$\begin{aligned} &\frac{\mu(w) |\varphi(w)|^k \left| \sum (n! / (k_1! \dots k_n!)) \prod_{j=1}^n (\varphi^{(j)}(w) / j!)^{k_j} \right|}{(1 - |\varphi(w)|^2)^{k+(\alpha+2)/p}} \\ &\leq \sup_{w \in \mathbb{D}} \|C_\varphi(g_{\varphi(w),k})\|_{\mathcal{W}_\mu^{(n)}(\mathbb{D})} \leq C \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})}, \end{aligned} \quad (3.10)$$

where (for each fixed $k \in \{1, \dots, n\}$) the sum is over all nonnegative integers k_1, k_2, \dots, k_n such that $k = k_1 + k_2 + \dots + k_n$ and $k_1 + 2k_2 + \dots + nk_n = n$.

From (3.10), it follows that for each $k \in \{1, \dots, n\}$,

$$\sup_{|\varphi(z)| > 1/2} \frac{\mu(z) \left| \sum (n! / (k_1! \dots k_n!)) \prod_{j=1}^n (\varphi^{(j)}(z) / j!)^{k_j} \right|}{(1 - |\varphi(z)|^2)^{k+(\alpha+2)/p}} \leq C \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})}. \quad (3.11)$$

Now we use consecutively the test functions

$$h_k(z) = z^k \in A_\alpha^p(\mathbb{D}), \quad k = 1, \dots, n, \quad (3.12)$$

in order to deal with the case $|\varphi(z)| \leq 1/2$. Note that

$$\|h_k\|_{A_\alpha^p(\mathbb{D})} \leq 1, \quad \text{for each } k \in \mathbb{N}. \quad (3.13)$$

By applying (2.11) to the function $f(z) = h_1(z)$, we get

$$(h_1 \circ \varphi)^{(n)}(z) = h_1'(\varphi(z))B_{n,1}(\varphi'(z), \dots, \varphi^{(n)}(z)) = B_{n,1}(\varphi'(z), \dots, \varphi^{(n)}(z)), \quad (3.14)$$

which along with the boundedness of the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})$ and (3.13) implies that

$$\sup_{z \in \mathbb{B}} \mu(z) \left| B_{n,1}(\varphi'(z), \dots, \varphi^{(n)}(z)) \right| \leq \|C_\varphi\|_{\mathcal{W}_\mu^{(n)}(\mathbb{D})} \leq \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})}, \quad (3.15)$$

or equivalently $\varphi \in \mathcal{W}_\mu^{(n)}(\mathbb{D})$ (see Remark 2.4).

Further, by applying formula (2.11) to the function $f(z) = h_2(z)$, we get

$$\begin{aligned} (h_2 \circ \varphi)^{(n)}(z) &= h_2'(\varphi(z))B_{n,1}(\varphi'(z), \dots, \varphi^{(n)}(z)) \\ &\quad + h_2''(\varphi(z))B_{n,2}(\varphi'(z), \dots, \varphi^{(n-1)}(z)). \end{aligned} \quad (3.16)$$

From the boundedness of $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})$ and (3.13), we get

$$\sup_{z \in \mathbb{B}} \mu(z) \left| (h_2 \circ \varphi)^{(n)}(z) \right| \leq \|C_\varphi(z^2)\|_{\mathcal{W}_\mu^{(n)}(\mathbb{D})} \leq \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})}. \quad (3.17)$$

From (3.16) and (3.17), and by using the triangle inequality it follows that

$$\begin{aligned} &2 \sup_{z \in \mathbb{B}} \mu(z) \left| B_{n,2}(\varphi'(z), \dots, \varphi^{(n-1)}(z)) \right| \\ &\leq \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})} + 2 \sup_{z \in \mathbb{B}} \mu(z) \left| \varphi(z) B_{n,1}(\varphi'(z), \dots, \varphi^{(n)}(z)) \right|. \end{aligned} \quad (3.18)$$

Using the fact $\sup_{z \in \mathbb{D}} |\varphi(z)| \leq 1$ and applying inequality (3.15) in (3.18) we get

$$\sup_{z \in \mathbb{B}} \mu(z) \left| B_{n,2}(\varphi'(z), \dots, \varphi^{(n-1)}(z)) \right| \leq \frac{3}{2} \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{W}_\mu^{(n)}(\mathbb{D})}. \quad (3.19)$$

Assume that we have proved the following inequalities:

$$\sup_{z \in \mathbb{B}} \mu(z) \left| B_{n,j} \left(\varphi'(z), \dots, \varphi^{(n-j+1)}(z) \right) \right| \leq C \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{K}_\mu^{(n)}(\mathbb{D})}, \quad (3.20)$$

for $j \in \{1, \dots, k-1\}$ and a $k \leq n$.

Applying formula (2.11) to the function $f(z) = h_k(z)$, $k \in \{1, \dots, n\}$, we have that

$$\begin{aligned} (h_k \circ \varphi)^{(n)}(z) &= \sum_{j=1}^k h_k^{(j)}(\varphi(z)) B_{n,j} \left(\varphi'(z), \dots, \varphi^{(n-j+1)}(z) \right) \\ &= \sum_{j=1}^k k(k-1) \cdots (k-j+1) (\varphi(z))^{k-j} B_{n,j} \left(\varphi'(z), \dots, \varphi^{(n-j+1)}(z) \right). \end{aligned} \quad (3.21)$$

From this, by using the boundedness of the operator $C_\varphi : A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{K}_\mu^{(n)}(\mathbb{D})$, the boundedness of function φ , the triangle inequality, noticing that the coefficient at $B_{n,k}(\varphi'(z), \dots, \varphi^{(n-k+1)}(z))$ is independent of z (it is equal $k!$), and finally using hypothesis (3.20), we easily obtain

$$\sup_{z \in \mathbb{B}} \mu(z) \left| B_{n,k} \left(\varphi'(z), \dots, \varphi^{(n-k+1)}(z) \right) \right| \leq C \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{K}_\mu^{(n)}(\mathbb{D})}. \quad (3.22)$$

Hence, by induction, we get that (3.22) holds for each $k \in \{1, \dots, n\}$.

From (3.22) and bearing in mind Remark 2.4, for each fixed $k \in \{1, \dots, n\}$, we have that

$$\begin{aligned} &\sup_{|\varphi(z)| \leq 1/2} \frac{\mu(z) \left| \sum (n! / (k_1! \cdots k_n!)) \prod_{j=1}^n (\varphi^{(j)}(z) / j!)^{k_j} \right|}{(1 - |\varphi(z)|^2)^{k+(\alpha+2)/p}} \\ &\leq \sup_{z \in \mathbb{B}} \mu(z) \left| B_{n,k} \left(\varphi'(z), \dots, \varphi^{(n-k+1)}(z) \right) \right| \leq C \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{K}_\mu^{(n)}(\mathbb{D})}, \end{aligned} \quad (3.23)$$

where as usual for a fixed $k \in \{1, \dots, n\}$, the sum is over all nonnegative integers k_1, k_2, \dots, k_n such that $k = k_1 + k_2 + \dots + k_n$ and $k_1 + 2k_2 + \dots + nk_n = n$.

Hence from (3.11) and (3.23), we get

$$\sum_{k=1}^n I_k \leq C \|C_\varphi\|_{A_\alpha^p(\mathbb{D}) \rightarrow \mathcal{K}_\mu^{(n)}(\mathbb{D})}. \quad (3.24)$$

From (3.5) and (3.24), we obtain asymptotic relation (3.2). □

Acknowledgment

The author would like to express his sincere thanks to the referees for numerous comments which improved the presentation of this paper.

References

- [1] B. R. Choe, H. Koo, and W. Smith, "Composition operators on small spaces," *Integral Equations and Operator Theory*, vol. 56, no. 3, pp. 357–380, 2006.
- [2] S. Li and S. Stević, "Volterra-type operators on Zygmund spaces," *Journal of Inequalities and Applications*, vol. 2007, Article ID 32124, 10 pages, 2007.
- [3] S. Li and S. Stević, "Generalized composition operators on Zygmund spaces and Bloch type spaces," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1282–1295, 2008.
- [4] S. Li and S. Stević, "Products of Volterra type operator and composition operator from H^∞ and Bloch spaces to Zygmund spaces," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 1, pp. 40–52, 2008.
- [5] S. Li and S. Stević, "Weighted composition operators from Zygmund spaces into Bloch spaces," *Applied Mathematics and Computation*, vol. 206, no. 2, pp. 825–831, 2008.
- [6] X. Zhu, "Volterra type operators from logarithmic Bloch spaces to Zygmund type spaces," *International Journal of Modern Mathematics*, vol. 3, no. 3, pp. 327–336, 2008.
- [7] S. Stević, "Composition operators from the Hardy space to the Zygmund-type space on the upper half-plane," *Abstract and Applied Analysis*, vol. 2009, Article ID 161528, 8 pages, 2009.
- [8] S. Stević, "Composition operators from the Hardy space to Zygmund-type spaces on the upper half-plane and the unit disk," *Journal of Computational Analysis and Applications*, vol. 12, 2010, (to appear).
- [9] S. Li and S. Stević, "Cesàro-type operators on some spaces of analytic functions on the unit ball," *Applied Mathematics and Computation*, vol. 208, no. 2, pp. 378–388, 2009.
- [10] S. Li and S. Stević, "Integral-type operators from Bloch-type spaces to Zygmund-type spaces," *Applied Mathematics and Computation*, vol. 215, no. 2, pp. 464–473, 2009.
- [11] S. Stević, "On an integral operator from the Zygmund space to the Bloch-type space on the unit ball," *Glasgow Mathematical Journal*, vol. 51, no. 2, pp. 275–287, 2009.
- [12] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, vol. 226 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2005.
- [13] X. Zhu, "Extended Cesàro operators from H^∞ to Zygmund type spaces in the unit ball," *Journal of Computational Analysis and Applications*, vol. 11, no. 2, pp. 356–363, 2009.
- [14] C. C. Cowen and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, Fla, USA, 1995.
- [15] X. Fu and X. Zhu, "Weighted composition operators on some weighted spaces in the unit ball," *Abstract and Applied Analysis*, vol. 2008, Article ID 605807, 8 pages, 2008.
- [16] D. Gu, "Weighted composition operators from generalized weighted Bergman spaces to weighted-type spaces," *Journal of Inequalities and Applications*, vol. 2008, Article ID 619525, 14 pages, 2008.
- [17] S. Li and S. Stević, "Weighted composition operators from α -Bloch space to H^∞ on the polydisc," *Numerical Functional Analysis and Optimization*, vol. 28, no. 7-8, pp. 911–925, 2007.
- [18] S. Li and S. Stević, "Weighted composition operators from H^∞ to the Bloch space on the polydisc," *Abstract and Applied Analysis*, vol. 2007, Article ID 48478, 13 pages, 2007.
- [19] S. Li and S. Stević, "Weighted composition operators between H^∞ and α -Bloch spaces in the unit ball," *Taiwanese Journal of Mathematics*, vol. 12, no. 7, pp. 1625–1639, 2008.
- [20] S. Ohno, K. Stroethoff, and R. Zhao, "Weighted composition operators between Bloch-type spaces," *The Rocky Mountain Journal of Mathematics*, vol. 33, no. 1, pp. 191–215, 2003.
- [21] S. Stević, "Composition operators between H^∞ and α -Bloch spaces on the polydisc," *Zeitschrift für Analysis und ihre Anwendungen*, vol. 25, no. 4, pp. 457–466, 2006.
- [22] S. Stević, "Weighted composition operators between mixed norm spaces and H^∞_α spaces in the unit ball," *Journal of Inequalities and Applications*, vol. 2007, Article ID 28629, 9 pages, 2007.
- [23] S. Stević, "Essential norms of weighted composition operators from the α -Bloch space to a weighted-type space on the unit ball," *Abstract and Applied Analysis*, vol. 2008, Article ID 279691, 11 pages, 2008.
- [24] S. Stević, "Norms of some operators from Bergman spaces to weighted and Bloch-type spaces," *Utilitas Mathematica*, vol. 76, pp. 59–64, 2008.

- [25] S. Stević, "Norm of weighted composition operators from Bloch space to H_μ^∞ on the unit ball," *Ars Combinatoria*, vol. 88, pp. 125–127, 2008.
- [26] S. Stević, "Weighted composition operators from mixed norm spaces into weighted Bloch spaces," *Journal of Computational Analysis and Applications*, vol. 11, no. 1, pp. 70–80, 2009.
- [27] S.-I. Ueki, "Composition operators on the Privalov spaces of the unit ball of \mathbb{C}^n ," *Journal of the Korean Mathematical Society*, vol. 42, no. 1, pp. 111–127, 2005.
- [28] S.-I. Ueki, "Weighted composition operators on the Bargmann-Fock space," *International Journal of Modern Mathematics*, vol. 3, no. 3, pp. 231–243, 2008.
- [29] S. I. Ueki, "Weighted composition operators on some function spaces of entire functions," to appear in *Bulletin of the Belgian Mathematical Society. Simon Stevin*.
- [30] S.-I. Ueki and L. Luo, "Compact weighted composition operators and multiplication operators between Hardy spaces," *Abstract and Applied Analysis*, vol. 2008, Article ID 196498, 12 pages, 2008.
- [31] E. Wolf, "Weighted composition operators between weighted Bergman spaces and weighted Banach spaces of holomorphic functions," *Revista Matemática Complutense*, vol. 21, no. 2, pp. 475–480, 2008.
- [32] E. Wolf, "Weighted composition operators between weighted Bergman spaces and weighted Bloch type spaces," *Journal of Computational Analysis and Applications*, vol. 11, no. 2, pp. 317–321, 2009.
- [33] W. Yang, "Weighted composition operators from Bloch-type spaces to weighted-type spaces," to appear in *Ars Combinatoria*.
- [34] X. Zhu, "Weighted composition operators from $F(p, q, s)$ spaces to H_μ^∞ spaces," *Abstract and Applied Analysis*, vol. 2009, Article ID 290978, 14 pages, 2009.
- [35] X. Zhu, "Weighted composition operators between H^∞ and Bergman type spaces," *Communications of the Korean Mathematical Society*, vol. 21, no. 4, pp. 719–727, 2006.
- [36] A. Montes-Rodríguez, "Weighted composition operators on weighted Banach spaces of analytic functions," *Journal of the London Mathematical Society*, vol. 61, no. 3, pp. 872–884, 2000.
- [37] W. Rudin, *Function Theory in the Unit Ball of \mathbb{C}^n* , vol. 241 of *Fundamental Principles of Mathematical Science*, Springer, New York, NY, USA, 1980.
- [38] T. M. Flett, "The dual of an inequality of Hardy and Littlewood and some related inequalities," *Journal of Mathematical Analysis and Applications*, vol. 38, pp. 746–765, 1972.
- [39] Wikipedia, http://en.wikipedia.org/wiki/Fa%C3%A0_di_Bruno%E2%80%99s_formula.