Hindawi Publishing Corporation
Differential Equations and Nonlinear Mechanics
Volume 2007, Article ID 19685, 9 pages
doi:10.1155/2007/19685

*Research Article*

# Global Existence and Asymptotic Behavior of Solutions for a Class of Nonlinear Degenerate Wave Equations

Yaojun Ye

Received 20 December 2006; Accepted 10 April 2007

Recommended by Ramon Quintanilla

This paper studies the existence of global solutions to the initial-boundary value problem for some nonlinear degenerate wave equations by means of compactness method and the potential well idea. Meanwhile, we investigate the decay estimate of the energy of the global solutions to this problem by using a difference inequality.

Copyright © 2007 Yaojun Ye. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

We are concerned with the following nonlinear degenerate wave equation:

$$u_{tt} - \operatorname{div}\left(|Du|^{p-2}Du\right) + u_t = |u|^{m-2}u, \quad (x,t) \in \Omega \times [0,+\infty), \tag{1.1}$$

with the initial-boundary value conditions

$$u(x,0) = u_0(x), \quad u_t(x,0) = u_1(x), \quad x \in \Omega; \quad u(x,t)\big|_{\partial\Omega\times[0,+\infty)} = 0, \tag{1.2}$$

where $\Omega$ is a bounded open domain in $\mathbb{R}^n$ with a smooth boundary $\partial\Omega$, $m \geq 2$ is a nonnegative real number, and $\operatorname{div}(|Du|^{p-2}Du)$ is a divergence operator (degenerate Laplace operator) with $p > 2$ and $Du = (D_1u, D_2u, \ldots, D_nu)$, $D_i = \partial/\partial x_i$, $i = 1,2,\ldots,n$.

When $p = 2$, (1.1) is converted into the form

$$u_{tt} - \triangle u + u_t = |u|^{m-2}u. \tag{1.3}$$

The global existence, the decay property of weak solutions, and the blow up of solutions to the initial-boundary value problem for the semilinear wave equations related to (1.2)-(1.3), under suitable assumptive conditions, have been investigated by many people

through various approaches [1–4]. However, little attention is paid to problem (1.1)-(1.2). Because the divergence operator $\operatorname{div}(|Du|^{p-2}Du)$ is a nonlinear operator, the reasonable proof and computation are greatly different from the Laplace operator; thus, the investigation of problem (1.1)-(1.2) becomes more complicated. In this paper, on the one hand, by a Galerkin approximation scheme [5], as well as combining it with the potential well method, we prove the global existence of solutions to problem (1.1)-(1.2). On the other hand, we obtain the asymptotic behavior of the global solutions to this problem by using a difference inequality.

For simplicity of notation, hereafter we denote by $\|\cdot\|_p$ the space $L^p(\Omega)$ norm. $\|\cdot\|$ denotes $L^2(\Omega)$ norm and we write equivalent norm $\|\bigtriangledown\cdot\|_p$ instead of $W_0^{1,p}(\Omega)$ norm $\|\cdot\|_{W_0^{1,p}(\Omega)}$, $C$ denotes various positive constants depending on the known constants and may be different at each appearance.

We define the functionals

$$J(u)=\frac{1}{p}\|\nabla u\|_p^p-\frac{1}{m}\|u\|_m^m,\quad K(u)=\|\nabla u\|_p^p-\|u\|_m^m,\quad u\in W_0^{1,p}(\Omega),\tag{1.4}$$

and according to [6] we put

$$d=\inf\left\{\sup_{\lambda\geq 0}J(\lambda u):u\in W_0^{1,p}(\Omega)/\{0\}\right\}.\tag{1.5}$$

Then, for problem (1.1)-(1.2) we are able to define the stable sets as follows:

$$W=\{u:u\in W_0^{1,p}(\Omega),\ K(u)>0,\ J(u)<d\}\cup\{0\}.\tag{1.6}$$

We denote the total energy related to (1.1) by

$$E(t)=E(u(t))=\frac{1}{2}||u_t||^2+\frac{1}{p}\|\nabla u\|_p^p-\frac{1}{m}\|u\|_m^m=\frac{1}{2}||u_t||^2+J(u(t)),\quad t\geq 0,\tag{1.7}$$

and $E(0)=(1/2)\|u_1\|^2+J(u_0)$ is the total energy of the initial data.

## 2. Some lemmas

We list up some useful lemmas here for the following discussion.

Lemma 2.1. *Let $a\geq 0$, $b\geq 0$ and $1/p+1/q=1$ for $1<p,q<+\infty$, then one has the inequality*

$$ab\leq\delta a^p+C(\delta)b^q,\tag{2.1}$$

*where $\delta>0$ is an arbitrary constant and $C(\delta)>0$ is a positive constant depending on $\delta$.*

Lemma 2.2. *Let $u\in W_0^{1,p}(\Omega)$, then $u\in L^q(\Omega)$ and the inequality $\|u\|_q\leq C\|u\|_{W_0^{1,p}(\Omega)}$ holds, provided that (i) $1\leq q\leq np/(n-p)$ if $1<p<n$; (ii) $1\leq q<+\infty$ if $1\leq n\leq p$.*

Lemma 2.3. *Assume that $u\in W_0^{1,p}(\Omega)$ and (i) $p<m<np/(n-p)$ for $2<p<n$; (ii) $p<m<+\infty$ for $n\leq p$; then $d$ is a positive real number.*

*Proof.* By Lemma 2.2, we have $\|u\|_m \le C\|\nabla u\|_p$. Since

$$J(\lambda u) = \frac{\lambda^p}{p}\|\nabla u\|_p^p - \frac{\lambda^m}{m}\|u\|_m^m, \tag{2.2}$$

we get

$$\frac{d}{d\lambda}J(\lambda u) = \lambda^{p-1}\|\nabla u\|_p^p - \lambda^{m-1}\|u\|_m^m. \tag{2.3}$$

Let $(d/d\lambda)J(\lambda u) = 0$, which implies that

$$\lambda_1 = \left(\frac{\|\nabla u\|_p^p}{\|u\|_m^m}\right)^{1/(m-p)}. \tag{2.4}$$

An elementary calculation shows that

$$\frac{d^2}{d\lambda^2}J(\lambda u)\Big|_{\lambda=\lambda_1} < 0. \tag{2.5}$$

So, we have

$$\begin{aligned}\sup_{\lambda\ge 0} J(\lambda u) = J(\lambda_1 u) &= \frac{\lambda_1^p}{p}\|\nabla u\|_p^p - \frac{\lambda_1^m}{m}\|u\|_m^m \\ &= \frac{m-p}{mp}\left(\frac{\|\nabla u\|_p}{\|u\|_m}\right)^{mp/(m-p)} \ge \frac{m-p}{mp}C^{mp/(p-m)} > 0.\end{aligned} \tag{2.6}$$

Therefore

$$d = \inf\left\{\sup_{\lambda\ge 0} J(\lambda u) : u \in W_0^{1,p}(\Omega)/\{0\}\right\} > 0. \tag{2.7}$$

This completes the proof of Lemma 2.3. □

Lemma 2.4. *Provided that (i) $p < m < np/(n-p)$ for $2 < p < n$; (ii) $p < m < +\infty$ for $n \le p$, then $d$ is a finite real number and the set $W$ is bounded in $W_0^{1,p}(\Omega)$.*

*Proof.* From the proof of Lemma 2.3 and the definition of $d$, we have for any $u \in W_0^{1,p}(\Omega)$ that

$$\begin{aligned}d \le \sup_{\lambda\ge 0} J(\lambda u) = J(\lambda_1 u) &= \frac{\lambda_1^p}{p}\|\nabla u\|_p^p - \frac{\lambda_1^m}{m}\|u\|_m^m \\ &= \frac{m-p}{mp}\left(\frac{\|\nabla u\|_p}{\|u\|_m}\right)^{mp/(m-p)} = \frac{m-p}{mp}\lambda_1^p\|\nabla u\|_p^p < +\infty.\end{aligned} \tag{2.8}$$

So $d$ is a finite real number.

Setting $u \in W$, then $\|\nabla u\|_p^p - \|u\|_m^m \geq 0$. Consequently,

$$d > J(u) = \frac{1}{p}\|\nabla u\|_p^p - \frac{1}{m}\|u\|_m^m \geq \frac{m-p}{mp}\|\nabla u\|_p^p, \tag{2.9}$$

which yields

$$\|\nabla u\|_p^p \leq \frac{mp}{m-p}d < +\infty. \tag{2.10}$$

As a result, $u \in W_0^{1,p}(\Omega)$ and

$$W \subset \left\{u : u \in W_0^{1,p}(\Omega),\ \|\nabla u\|_p^p \leq \frac{mp}{m-p}d\right\}. \tag{2.11}$$

Thus the stable set $W$ is bounded in $W_0^{1,p}(\Omega)$. □

Lemma 2.5 [7]. *Suppose that $\phi(t)$ is a nonincreasing nonnegative function on $[0,+\infty)$ and satisfies*

$$\phi(t)^{1+\alpha} \leq k\{\phi(t) - \phi(t+1)\} \tag{2.12}$$

*for some constants $\alpha > 0$ and $k > 0$. Then $\phi(t)$ has the decay property*

$$\phi(t) \leq \left\{\phi(0)^{-\alpha} + \alpha k^{-1}[t-1]^+\right\}^{-1/\alpha}, \quad t > 0, \tag{2.13}$$

*where $[t-1]^+ = \max\{t-1, 0\}$.*

*Proof.* Setting $\psi(t) = \phi(t)^{-\alpha}$, we see from (2.13) that

$$\begin{aligned}\psi(t+1) - \psi(t) &= \int_0^1 \frac{d}{d\theta}\{\theta\phi(t+1) + (1-\theta)\phi(t)\}^{-\alpha}d\theta \\ &= \alpha(\phi(t) - \phi(t+1))\int_0^1 \{\theta\phi(t+1) + (1-\theta)\phi(t)\}^{-\alpha-1}d\theta \\ &\geq \alpha k^{-1}.\end{aligned} \tag{2.14}$$

Then we get

$$\psi(t) \geq \psi(0) + \alpha k^{-1}t \tag{2.15}$$

and the desired estimate (2.13). □

## 3. The global existence

Theorem 3.1. *Given that $p \leq m \leq np/(n-p)$, $p < n$, and $p < m < +\infty$, $n \leq p$, if $u_0 \in W$, $u_1 \in L^2(\Omega)$ and the initial data energy $E(0) < d$, then problem (1.1)-(1.2) admits a global solution $u(x,t)$ such that $u(x,t) \in W$ and*

$$u(x,t) \in L^\infty\left(0,T;W_0^{1,p}(\Omega)\right), \qquad u_t(x,t) \in L^\infty\left(0,T;L^2(\Omega)\right). \tag{3.1}$$

*Proof.* Let $r$ be an integer for which $H_0^r(\Omega) \hookrightarrow W_0^{1,p}(\Omega)$ is continuous. Then the eigenfunctions of $-\Delta^r \omega_j = \alpha_j \omega_j$ in $H_0^r(\Omega)$ yield a Gaerkin basis for both $H_0^r(\Omega) \subset W_0^{1,p}(\Omega)$ and $L^2(\Omega)$. We seek approximate solutions $u_N(t)$ to the problem (1.1)-(1.2) of the form

$$u_N(t) = \sum_{j=1}^{N} g_{jN}(t)\omega_j, \quad N = 1,2,\ldots, \tag{3.2}$$

where the coefficients $g_{jN}(t)$ satisfy $g_{jN}(t) = (u_N(t), \omega_j)$ with

$$(u_N''(t), \omega_j) + (\operatorname{div}(|Du_N|^{p-2}Du_N), \omega_j) + (u_N'(t), \omega_j) = (|u_N|^{m-2}u_N, \omega_j), \tag{3.3}$$

$$u_N(0) = u_{0N}, \quad u_N'(0) = u_{1N}, \quad 1 \le j \le N. \tag{3.4}$$

Here $(u,v) = \int_\Omega u(x)v(x)dx$. Since $C_0^\infty(\Omega)$ is dense in $W_0^{1,p}(\Omega)$ and $L^2(\Omega)$, we choose $u_{0N}$, $u_{1N} \in C_0^\infty(\Omega)$ such that $u_N(0) = u_{0N} \to u_0(x)$ in $W_0^{1,p}(\Omega)$ and $u_N'(0) = u_{1N} \to u_1(x)$ in $L^2(\Omega)$ as $N \to \infty$.

We observe that (3.3) is a system of ordinary differential equations in the variable $t$ and has a local solution $u_N(t)$ in an interval $[0, t_m)$ by the existence theorem. In the next step, we obtain the a priori estimates for the solution $u_N(t)$ so that it can be extended to the whole interval $[0,T]$ according to the extension theorem.

Multiplying (3.3) by $g_{jN}'(t)$ and summing over $j$ from 1 to $N$, and then integrating over $[0,t]$; we get

$$\frac{1}{2}||u_N'(t)||^2 + J(u_N(t)) + \int_0^t ||u_N'(\tau)||^2 d\tau = \frac{1}{2}||u_{1N}||^2 + J(u_{0N}). \tag{3.5}$$

By using formula (3.5), we can obtain

$$u_N(t) \in W, \quad t \in [0, t_m). \tag{3.6}$$

In fact, suppose that (3.6) is false and let $t_1$ be the smallest time for $u_N(t_1) \notin W$. Then, by means of the continuity of $u_N(t)$, we see $u_N(t_1) \in \partial W$. From the definition of $W$ and the continuity of $J(u(t))$ and $K(u(t))$ in $t$, we have either

$$J(u_N(t_1)) = d, \tag{3.7}$$

or

$$K(u_N(t_1)) = 0. \tag{3.8}$$

By (3.5) together with the condition $E(u(0)) < d$, we have

$$J(u_N(t_1)) \le \frac{1}{2}||u_{1N}||^2 + J(u_{0N}) = E(u_N(0)) < d. \tag{3.9}$$

So, case (3.7) is impossible.

Assume that (3.8) holds, then we obtain

$$\frac{d}{d\lambda}J(\lambda u_N(t_1)) = \lambda^{p-1}(1-\lambda^{m-p})||\nabla u_N(t_1)||_p^p. \tag{3.10}$$

Consequently,

$$\sup_{\lambda\geq 0} J(\lambda u_N(t_1)) = J(\lambda u_N(t_1))\,|_{\lambda=1} = J(u_N(t_1)) < d, \tag{3.11}$$

which contradicts the definition of $d$. Therefore, case (3.8) is impossible as well. Thus, we verify that $u_N(t) \in W$, $t \in [0, t_m)$.

From (3.5) and (3.6), we have

$$\frac{1}{2}||u_N'||^2 + \frac{m-p}{mp}||\nabla u_N||_p^p + \int_0^t ||u_N'(\tau)||^2 d\tau \leq \frac{1}{2}||u_{1N}||^2 + d \leq C. \tag{3.12}$$

With this estimate, we can extend the approximate solutions $u_N(t)$ to the interval $[0, T]$ and we have

$$\{u_N\} \quad \text{is bounded in } L^\infty(0,T;W_0^{1,p}(\Omega)), \tag{3.13}$$

$$\{u_N'\} \quad \text{is bounded in } L^\infty(0,T;L^2(\Omega)), \tag{3.14}$$

$$\{u_N'\} \quad \text{is bounded in } L^2(0,T;L^2(\Omega)), \tag{3.15}$$

$$\operatorname{div}(|Du_N|^{p-2}Du_N) \quad \text{is bounded in } L^\infty(0,T;W_0^{-1,p/(p-1)}(\Omega)), \tag{3.16}$$

$$|u_N|^{m-2}u_N \quad \text{is bounded in } L^\infty(0,T;L^{m/(m-1)}(\Omega)). \tag{3.17}$$

Since our Galerkin basis was taken in the Hilbert space $H^r(\Omega) \subset W_0^{1,p}(\Omega)$, we can use the standard projection argument as described in [5]. Then from the approximate equation (3.3) and the estimates (3.13)–(3.17), we get

$$\{u_N''\} \quad \text{is bounded in } L^2(0,T;W^{-1,p/(p-1)}(\Omega)). \tag{3.18}$$

Now from (3.13)–(3.17) and the standard arguments of the approximate solutions, we conclude that after the extraction of suitable subsequence $\{u_\mu\}$ from $\{u_N\}$ if necessary, we have the following:

$$\{u_\mu\} \longrightarrow u \quad \text{weakly star in } L^\infty(0,T;W_0^{1,p}(\Omega)), \tag{3.19}$$

$$\{u_\mu'\} \longrightarrow u' \quad \text{weakly star in } L^\infty(0,T;L^2(\Omega)), \tag{3.20}$$

$$\{u_\mu'\} \longrightarrow u' \quad \text{weakly in } L^2(0,T;L^2(\Omega)), \tag{3.21}$$

$$\operatorname{div}(|Du_\mu|^{p-2}Du_\mu) \longrightarrow \chi_1 \quad \text{weakly star in } L^\infty(0,T;W_0^{-1,p/(p-1)}(\Omega)), \tag{3.22}$$

$$|u_\mu|^{m-2}u_\mu \longrightarrow \chi_2 \quad \text{weakly star in } L^\infty(0,T;L^{m/(m-1)}(\Omega)). \tag{3.23}$$

By applying the Lions-Aubin compactness lemma [5], we get that from (3.13) and (3.14),

$$\{u_\mu\} \longrightarrow u \quad \text{strongly in } L^2(0,T;L^2(\Omega)). \tag{3.24}$$

We receive that from (3.15) and (3.18)

$$\{u'_\mu\} \longrightarrow u' \quad \text{strongly in } L^2(0,T;L^2(\Omega)). \tag{3.25}$$

Using (3.13) and (3.24), we see that

$$\int_0^T \int_\Omega \left| \left| u_\mu \right|^{m-2} u_\mu \right|^{m/(m-1)} dx\, dt = \int_0^T ||u_\mu||_m^m dt \le C \int_0^T ||u_\mu||_{W_0^{1,p}}^m dt \le C, \tag{3.26}$$

and $|u_\mu|^{m-2}u_\mu \to |u|^{m-2}u$ almost everywhere in $(0,T)\times\Omega$. Therefore from [5, Lemma 1.3], we infer that

$$\left| u_\mu \right|^{m-2} u_\mu \longrightarrow |u|^{m-2}u \quad \text{weakly in } L^{m/(m-1)}(0,T;L^{m/(m-1)}(\Omega)). \tag{3.27}$$

We have from (3.23) and (3.27) that $\chi_2 = |u|^{m-2}u$. Finally, since we have the strong convergence (3.25), we can use a standard monotonicity argument as done by Lions in [5] or by Ye in [8] to show that $\chi_1 = \operatorname{div}(|Du|^{p-2}Du)$.

Multiplying both sides of (3.3) by $g(t) \in C^2[0,T]$ and letting $\mu = N \to \infty$, we get that $u(x,t)$ is a global solution of problem (1.1)-(1.2). This ends the proof of Theorem 3.1. □

## 4. The asymptotic behavior

Theorem 4.1. *Under the hypotheses of Theorem 3.1, the global solution $u(x,t)$ in $W$ of problem (1.1)-(1.2) on $[0,+\infty)$ has the following decay property:*

$$E(t) \le E(0)\left\{1 + CE(0)I_0^{-2}[t-1]^+\right\}^{-1}, \quad t \in (0,+\infty), \tag{4.1}$$

*where $I_0$ is some positive constant depending only on $u_0$ and $u_1$.*

*Proof.* Multiplying (1.1) by $u_t$ and integrating over $[t,t+1]\times\Omega$, $t>0$, we have

$$\int_t^{t+1} ||u_t(s)||^2 ds = E(t) - E(t+1) \equiv D(t)^2. \tag{4.2}$$

Thus, there exist $t_1 \in [t, t+1/4]$, $t_2 \in [t+3/4, t+1]$ such that

$$||u_t(t_i)|| \le 2D(t), \quad i = 1,2. \tag{4.3}$$

On the other hand, we multiply (1.1) by $u(t,x)$ and integrate over $[t_1,t_2]\times\Omega$, which yields

$$\begin{aligned}\int_{t_1}^{t_2} K(u(s))\,ds &= \int_{t_1}^{t_2} ||u_t(s)||^2 ds \\ &\quad + (u_t(t_1),u(t_1)) - (u_t(t_2),u(t_2)) - \int_{t_1}^{t_2} (u_t(s),u(s))\,ds \\ &\le D(t)^2 + 5D(t)\sup_{t\le s\le t+1}||u(s)||.\end{aligned} \tag{4.4}$$

To estimate $\|u(t)\|$, we multiply (1.1) by $u(t,x)$ and integrate over $[0,t]\times\Omega$ to obtain

$$\frac{1}{2}||u(t)||^2 + \int_0^t K(u(s))\,ds = \frac{1}{2}||u_0||^2 + (u_1,u_0) + \int_0^t ||u_t(s)||^2 ds - (u_t(t),u(t)). \tag{4.5}$$

Since $K(u(t))>0$, we derive that from Lemma 2.1,

$$\begin{aligned}||u(t)||^2 &\le ||u_0||^2 + 2(u_1,u_0) + 2\|u\|||u_t|| + 2\int_0^t ||u_t(s)||^2 ds \\ &\le ||u_0||^2 + 2(u_1,u_0) + 2||u_t(t)||^2 + \frac{1}{2}||u(t)||^2 + 2\int_0^t ||u_t(s)||^2 ds.\end{aligned} \tag{4.6}$$

From (4.2), we get

$$\int_0^t ||u_t(s)||^2 ds = E(0) - E(t) < E(0), \tag{4.7}$$

and hence we have from (4.6) that

$$||u(t)||^2 \le 2\{||u_0||^2 + 2(u_1,u_0) + 6E(0)\} \equiv I_0^2. \tag{4.8}$$

It follows from (4.4) and (4.8) that

$$\int_{t_1}^{t_2} K(u(s))\,ds \le D(t)^2 + 5I_0 D(t). \tag{4.9}$$

Now, it follows from (4.2) and (4.9) that

$$\begin{aligned}E(t_2) &\le 2\int_{t_1}^{t_2} E(s)ds \le C\{D(t)^2 + I_0 D(t)\}, \\ E(t_1) &= E(t_2) + \int_{t_1}^{t_2} ||u_t(s)||^2 ds \le E(t_2) + \int_t^{t+1} ||u_t(s)||^2 ds \\ &\le C\{D(t)^2 + I_0 D(t)\} + D(t)^2 \le C I_0 D(t),\end{aligned} \tag{4.10}$$

which implies by (4.2) that

$$\sup_{t\le s\le t+1} E(s)^2 \le C I_0^2 D(t)^2 = C I_0^2\{E(t) - E(t+1)\}. \tag{4.11}$$

Thus, applying Lemma 2.5 to (4.11) and using the fact that $E(t) \le E(0) < d$, we derive the decay estimate

$$E(t) \le \{E(0)^{-1} + CI_0^{-2}[t-1]^+\}^{-1} = E(0)\{1 + CE(0)I_0^{-2}[t-1]^+\}^{-1}. \tag{4.12}$$

This completes the proof of Theorem 4.1. □

**Acknowledgments**

This project is supported by NSF of China (Grant no. 10441002) and Henan Province (Grant no. 200510466011).

**References**

[1] M. Aassila, "Global existence and global nonexistence of solutions to a wave equation with nonlinear damping and source terms," *Asymptotic Analysis*, vol. 30, no. 3-4, pp. 301–311, 2002.
[2] V. Georgiev and G. Todorova, "Existence of a solution of the wave equation with nonlinear damping and source terms," *Journal of Differential Equations*, vol. 109, no. 2, pp. 295–308, 1994.
[3] M. Nakao and K. Ono, "Existence of global solutions to the Cauchy problem for the semilinear dissipative wave equations," *Mathematische Zeitschrift*, vol. 214, no. 2, pp. 325–342, 1993.
[4] G. Todorova, "Stable and unstable sets for the Cauchy problem for a nonlinear wave equation with nonlinear damping and source terms," *Journal of Mathematical Analysis and Applications*, vol. 239, no. 2, pp. 213–226, 1999.
[5] J.-L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites Non Linéaires*, Dunod, Paris, France, 1969.
[6] D. H. Sattinger, "On global solution of nonlinear hyperbolic equations," *Archive for Rational Mechanics and Analysis*, vol. 30, pp. 148–172, 1968.
[7] M. Nakao, "A difference inequality and its application to nonlinear evolution equations," *Journal of the Mathematical Society of Japan*, vol. 30, no. 4, pp. 747–762, 1978.
[8] Y. Ye, "Existence of global solutions for some nonlinear hyperbolic equation with a nonlinear dissipative term," *Journal of Zhengzhou University. Natural Science Edition*, vol. 29, no. 3, pp. 18–23, 1997.

Yaojun Ye: Department of Information and Computational Science, Henan Agricultural University, Zhengzhou 450002, China
*Email address*: yeyaojun2002@yahoo.com.cn

Mathematical Problems in Engineering

# Special Issue on Time-Dependent Billiards

## Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at http://www.hindawi.com/journals/mpe/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/ according to the following timetable:

| | |
|---|---|
| Manuscript Due | December 1, 2008 |
| First Round of Reviews | March 1, 2009 |
| Publication Date | June 1, 2009 |

**Guest Editors**

**Edson Denis Leonel,** Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

**Alexander Loskutov,** Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru