A NOTE ON WELL-POSED NULL AND FIXED POINT PROBLEMS

SIMEON REICH AND ALEXANDER J. ZASLAVSKI

Received 16 October 2004

We establish generic well-posedness of certain null and fixed point problems for ordered Banach space-valued continuous mappings.

The notion of well-posedness is of great importance in many areas of mathematics and its applications. In this note, we consider two complete metric spaces of continuous mappings and establish generic well-posedness of certain null and fixed point problems (Theorems 1 and 2, resp.). Our results are a consequence of the variational principle established in [2]. For other recent results concerning the well-posedness of fixed point problems, see [1, 3].

Let $(X, \|\cdot\|, \geq)$ be a Banach space ordered by a closed convex cone $X_+ = \{x \in X : x \geq 0\}$ such that $\|x\| \leq \|y\|$ for each pair of points $x, y \in X_+$ satisfying $x \leq y$. Let (K, ρ) be a complete metric space. Denote by $\mathfrak{M}$ the set of all continuous mappings $A : K \to X$. We equip the set $\mathfrak{M}$ with the uniformity determined by the following base:

$$E(\epsilon) = \{(A, B) \in \mathfrak{M} \times \mathfrak{M} : \|Ax - Bx\| \leq \epsilon \ \forall x \in K\}, \tag{1}$$

where $\epsilon > 0$. It is not difficult to see that this uniform space is metrizable (by a metric d) and complete.

Denote by $\mathfrak{M}_p$ the set of all $A \in \mathfrak{M}$ such that

$$\begin{aligned} Ax &\in X_+ \quad \forall x \in K, \\ \inf\{\|Ax\| : x \in K\} &= 0. \end{aligned} \tag{2}$$

It is not difficult to see that $\mathfrak{M}_p$ is a closed subset of $(\mathfrak{M}, d)$.

We can now state and prove our first result.

THEOREM 1. *There exists an everywhere dense G_δ subset $\mathcal{F} \subset \mathfrak{M}_p$ such that for each $A \in \mathcal{F}$, the following properties hold.*

(1) There is a unique $\bar{x} \in K$ such that $A\bar{x} = 0$.

(2) For any $\epsilon > 0$, there exist $\delta > 0$ and a neighborhood U of A in $\mathfrak{M}_p$ such that if $B \in U$ and if $x \in K$ satisfies $\|Bx\| \leq \delta$, then $\rho(x, \bar{x}) \leq \epsilon$.

Copyright © 2005 Hindawi Publishing Corporation
Fixed Point Theory and Applications 2005:2 (2005) 207–211
DOI: 10.1155/FPTA.2005.207

Proof. We obtain this theorem as a realization of the variational principle established in [2, Theorem 2.1] with $f_A(x) = \|Ax\|$, $x \in K$. In order to prove our theorem by using this variational principle, we need to prove the following assertion.

(A) For each $A \in \mathfrak{M}_p$ and each $\epsilon > 0$, there are $\bar{A} \in \mathfrak{M}_p$, $\delta > 0$, $\bar{x} \in K$, and a neighborhood W of $\bar{A}$ in $\mathfrak{M}_p$ such that

$$(A, \bar{A}) \in E(\epsilon), \tag{3}$$

and if $B \in W$ and $z \in K$ satisfy $\|Bz\| \le \delta$, then

$$\rho(z, \bar{x}) \le \epsilon. \tag{4}$$

Let $A \in \mathfrak{M}_p$ and $\epsilon > 0$. Choose $\bar{u} \in X_+$ such that

$$\|\bar{u}\| = \frac{\epsilon}{4}, \tag{5}$$

and $\bar{x} \in K$ such that

$$\|A\bar{x}\| \le \frac{\epsilon}{8}. \tag{6}$$

Since A is continuous, there is a positive number r such that

$$r < \min\left\{1, \frac{\epsilon}{16}\right\}, \tag{7}$$

$$\|Ax - A\bar{x}\| \le \frac{\epsilon}{8} \quad \text{for each } x \in K \text{ satisfying } \rho(x, \bar{x}) \le 4r. \tag{8}$$

By Urysohn's theorem, there is a continuous function $\phi : K \to [0,1]$ such that

$$\phi(x) = 1 \quad \text{for each } x \in K \text{ satisfying } \rho(x, \bar{x}) \le r, \tag{9}$$

$$\phi(x) = 0 \quad \text{for each } x \in K \text{ satisfying } \rho(x, \bar{x}) \ge 2r. \tag{10}$$

Define

$$\bar{A}x = \big(1 - \phi(x)\big)(Ax + \bar{u}), \quad x \in K. \tag{11}$$

It is clear that $\bar{A} : K \to X$ is continuous. Now (9), (10), and (11) imply that

$$\bar{A}x = 0 \quad \text{for each } x \in K \text{ satisfying } \rho(x, \bar{x}) \le r, \tag{12}$$

$$\bar{A}x \ge \bar{u} \quad \text{for each } x \in K \text{ satisfying } \rho(x, \bar{x}) \ge 2r. \tag{13}$$

It is not difficult to see that $\bar{A} \in \mathfrak{M}_p$. We claim that $(A, \bar{A}) \in E(\epsilon)$.

Let $x \in K$. There are two cases: either

$$\rho(x,\bar{x}) \geq 2r \tag{14}$$

or

$$\rho(x,\bar{x}) < 2r. \tag{15}$$

Assume first that (14) holds. Then it follows from (14), (10), (11), and (5) that

$$\|Ax - \bar{A}x\| = \|\bar{u}\| = \frac{\epsilon}{4}. \tag{16}$$

Now assume that (15) holds. Then by (15), (11), and (5),

$$\begin{aligned} \|\bar{A}x - Ax\| &= \big\|\big(1-\phi(x)\big)(Ax+\bar{u}) - Ax\big\| \\ &\leq \|\bar{u}\| + \|Ax\| \leq \frac{\epsilon}{4} + \|Ax\|. \end{aligned} \tag{17}$$

It follows from this inequality, (15), (8), and (6) that

$$\|\bar{A}x - Ax\| \leq \frac{\epsilon}{4} + \|Ax\| < \frac{\epsilon}{2}. \tag{18}$$

Therefore, in both cases, $\|\bar{A}x - Ax\| \leq \epsilon/2$. Since this inequality holds for any $x \in K$, we conclude that

$$(A,\bar{A}) \in E(\epsilon). \tag{19}$$

Consider now an open neighborhood U of $\bar{A}$ in $\mathfrak{M}_p$ such that

$$U \subset \left\{B \in \mathfrak{M}_p : (\bar{A},B) \in E\left(\frac{\epsilon}{16}\right)\right\}. \tag{20}$$

Let

$$B \in U, \qquad z \in K, \tag{21}$$

$$\|Bz\| \leq \frac{\epsilon}{16}. \tag{22}$$

Relations (22), (21), (20), and (1) imply that

$$\|\bar{A}z\| \leq \|Bz\| + \|\bar{A}z - Bz\| \leq \frac{\epsilon}{16} + \frac{\epsilon}{16}. \tag{23}$$

We claim that

$$\rho(z,\bar{x}) \leq \epsilon. \tag{24}$$

We assume the converse. Then by (7),

$$\rho(z,\bar{x}) > \epsilon \geq 2r. \tag{25}$$

When combined with (13), this implies that

$$\bar{A}z \geq \bar{u}. \tag{26}$$

It follows from this inequality, the monotonicity of the norm, (21), (20), (1), and (5) that

$$\begin{aligned} \|Bz\| &\geq \|\bar{A}z\| - \frac{\epsilon}{16} \geq \|\bar{u}\| - \frac{\epsilon}{16} \\ &= \frac{\epsilon}{4} - \frac{\epsilon}{16} = \frac{3\epsilon}{16}. \end{aligned} \tag{27}$$

This, however, contradicts (22). The contradiction we have reached proves (24) and Theorem 1 itself. □

Now assume that the set K is a subset of X and

$$\rho(x,y) = \|x-y\|, \quad x,y \in K. \tag{28}$$

Denote by $\mathfrak{M}_n$ the set of all mappings $A \in \mathfrak{M}$ such that

$$\begin{aligned} Ax &\geq x \quad \forall x \in K, \\ \inf\{\|Ax - x\| : x \in K\} &= 0. \end{aligned} \tag{29}$$

Clearly, $\mathfrak{M}_n$ is a closed subset of $(\mathfrak{M}, d)$. Define a map $J : \mathfrak{M}_n \to \mathfrak{M}_p$ by

$$J(A)x = Ax - x \quad \forall x \in K \tag{30}$$

and all $A \in \mathfrak{M}_n$. Clearly, there exists $J^{-1} : \mathfrak{M}_p \to \mathfrak{M}_n$, and both J and its inverse J^{-1} are continuous. Therefore Theorem 1 implies the following result regarding the generic well-posedness of the fixed point problem for $A \in \mathfrak{M}_n$.

Theorem 2. *There exists an everywhere dense G_δ subset $\mathfrak{F} \subset \mathfrak{M}_n$ such that for each $A \in \mathfrak{F}$, the following properties hold.*

(1) There is a unique $\bar{x} \in K$ such that $A\bar{x} = \bar{x}$.

(2) For any $\epsilon > 0$, there exist $\delta > 0$ and a neighborhood U of A in $\mathfrak{M}_n$ such that if $B \in U$ and if $x \in K$ satisfies $\|Bx - x\| \leq \delta$, then $\|x - \bar{x}\| \leq \epsilon$.

Acknowledgments

The work of the first author was partially supported by the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities (Grant 592/00), by the Fund for the Promotion of Research at the Technion, and by the Technion VPR Fund.

References

[1] F. S. De Blasi and J. Myjak, *Sur la porosité de l'ensemble des contractions sans point fixe* [*On the porosity of the set of contractions without fixed points*], C. R. Acad. Sci. Paris Sér. I Math. **308** (1989), no. 2, 51–54 (French).

[2] A. D. Ioffe and A. J. Zaslavski, *Variational principles and well-posedness in optimization and calculus of variations*, SIAM J. Control Optim. **38** (2000), no. 2, 566–581.

[3] S. Reich and A. J. Zaslavski, *Well-posedness of fixed point problems*, Far East J. Math. Sci. (FJMS), (2001), Special Volume (Functional Analysis and Its Applications), Part III, 393–401.

Simeon Reich: Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, 2-12-1 O-okayama, Meguro-ku, Tokyo 152-8552, Japan

E-mail address: sreich@tx.technion.ac.il

Alexander J. Zaslavski: Department of Mathematics, Technion – Israel Institute of Technology, 32000 Haifa, Israel

E-mail address: ajzasl@tx.technion.ac.il

Mathematical Problems in Engineering

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at http://www.hindawi.com/journals/mpe/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/ according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru