

WEAK AND STRONG CONVERGENCE THEOREMS FOR NONEXPANSIVE SEMIGROUPS IN BANACH SPACES

SACHIKO ATSUSHIBA AND WATARU TAKAHASHI

Received 24 February 2005

We introduce an implicit iterative process for a nonexpansive semigroup and then we prove a weak convergence theorem for the nonexpansive semigroup in a uniformly convex Banach space which satisfies Opial's condition. Further, we discuss the strong convergence of the implicit iterative process.

1. Introduction

Let C be a closed convex subset of a Hilbert space and let T be a nonexpansive mapping from C into itself. For each $t \in (0, 1)$, the contraction mapping T_t of C into itself defined by

$$T_t x = tu + (1 - t)Tx \quad (1.1)$$

for every $x \in C$, has a unique fixed point x_t , where u is an element of C . Browder [4] proved that $\{x_t\}$ converges strongly to a fixed point of T as $t \rightarrow 0$ in a Hilbert space. Motivated by Browder's theorem [4], Takahashi and Ueda [20] proved the strong convergence of the following iterative process in a uniformly convex Banach space with a uniformly Gâteaux differentiable norm (see also [14]):

$$x_k = \frac{1}{k}x + \left(1 - \frac{1}{k}\right)Tx_k \quad (1.2)$$

for every $k = 1, 2, 3, \dots$, where $x \in C$. On the other hand, Xu and Ori [21] studied the following implicit iterative process for finite nonexpansive mappings $T_1, T_2, \dots, T_r$ in a Hilbert space: $x_0 = x \in C$ and

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n)T_n x_n \quad (1.3)$$

for every $n = 1, 2, \dots$, where $\{\alpha_n\}$ is a sequence in $(0, 1)$ and $T_n = T_{n+r}$. And they proved a weak convergence of the iterative process defined by (1.3) in a Hilbert space. Sun et al. [17] studied the iterations defined by (1.3) and proved the strong convergence of the iterations in a uniformly convex Banach space, requiring one mapping T_i in the family to be semi compact.

In this paper, using the idea of [17, 21], we introduce an implicit iterative process for a nonexpansive semigroup and then prove a weak convergence theorem for the nonexpansive semigroup in a uniformly convex Banach space which satisfies Opial's condition. Further, we discuss the strong convergence of the implicit iterative process (see also [1, 2, 7, 12, 13]).

2. Preliminaries and notations

Throughout this paper, we denote by $\mathbb{N}$ and $\mathbb{Z}^+$ the set of all positive integers and the set of all nonnegative integers, respectively. Let E be a real Banach space. We denote by B_r the set $\{x \in E : \|x\| \leq r\}$. A Banach space E is said to be *strictly convex* if $\|x + y\|/2 < 1$ for each $x, y \in B_1$ with $x \neq y$, and it is said to be *uniformly convex* if for each $\varepsilon > 0$, there exists $\delta > 0$ such that $\|x + y\|/2 \leq 1 - \delta$ for each $x, y \in B_1$ with $\|x - y\| \geq \varepsilon$. It is well-known that a uniformly convex Banach space is reflexive and strictly convex (see [19]). Let C be a closed subset of a Banach space and let T be a mapping from C into itself. We denote by $F(T)$ and $F_\varepsilon(T)$ for $\varepsilon > 0$, the sets $\{x \in C : x = Tx\}$ and $\{x \in C : \|x - Tx\| \leq \varepsilon\}$, respectively.

A mapping T of C into itself is said to be *compact* if T is continuous and maps bounded sets into relatively compact sets. A mapping T of C into itself is said to be *demicompact* at $\xi \in C$ if for any bounded sequence $\{y_n\}$ in C such that $y_n - Ty_n \rightarrow \xi$ as $n \rightarrow \infty$, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ and $y \in C$ such that $y_{n_k} \rightarrow y$ as $k \rightarrow \infty$ and $y - Ty = \xi$. In particular, a continuous mapping T is *demicompact at 0* if for any bounded sequence $\{y_n\}$ in C such that $y_n - Ty_n \rightarrow 0$ as $n \rightarrow \infty$, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ and $y \in C$ such that $y_{n_k} \rightarrow y$ as $k \rightarrow \infty$. T is also said to be *semicompact* if T is continuous and *demicompact at 0* (e.g., see [21]). T is said to be *demicompact on C* if T is demicompact for each $y \in C$. If T is compact on C , then T is demicompact on C . For examples of demicompact mappings, see [1, 2, 12, 13]. We also denote by I the identity mapping. A mapping T of C into itself is said to be *nonexpansive* if $\|Tx - Ty\| \leq \|x - y\|$ for every $x, y \in C$. We denote by $N(C)$ the set of all nonexpansive mappings from C into itself. We know from [5] that if C is a nonempty closed convex subset of a strictly convex Banach space, then $F(T)$ is convex for each $T \in N(C)$ with $F(T) \neq \emptyset$. The following are crucial to prove our results (see [5, 6]).

PROPOSITION 2.1 (Browder). *Let C be a nonempty bounded closed convex subset of a uniformly convex Banach space and let T be a nonexpansive mapping from C into itself. Let $\{x_n\}$ be a sequence in C such that it converges weakly to an element x of C and $\{x_n - Tx_n\}$ converges strongly to 0. Then x is a fixed point of T .*

PROPOSITION 2.2 (Bruck). *Let E be a uniformly convex Banach space and let C be a nonempty closed convex subset of E . For any $\varepsilon > 0$, there exists $\delta > 0$ such that for any nonexpansive mapping T of C into itself with $F(T) \neq \emptyset$,*

$$\overline{\text{co}}F_\delta(T) \subset F_\varepsilon(T). \quad (2.1)$$

Let E^* be the dual space of a Banach space E . The value of $x^* \in E^*$ at $x \in E$ will be denoted by $\langle x, x^* \rangle$. We say that a Banach space E satisfies *Opial's condition* [11] if for each

sequence $\{x_n\}$ in E which converges weakly to x ,

$$\lim_{n \rightarrow \infty} \|x_n - x\| < \lim_{n \rightarrow \infty} \|x_n - y\| \quad (2.2)$$

for each $y \in E$ with $y \neq x$. Since if the duality mapping $x \mapsto \{x^* \in E^* : \langle x, x^* \rangle = \|x\|^2 = \|x^*\|^2\}$ from E into E^* is single-valued and weakly sequentially continuous, then E satisfies Opial's condition. Each Hilbert space and the sequence spaces ℓ^p with $1 < p < \infty$ satisfy Opial's condition (see [8, 11]). Though an L^p -space with $p \neq 2$ does not usually satisfy Opial's condition, each separable Banach space can be equivalently renormed so that it satisfies Opial's condition (see [11, 22]).

Let S be a semigroup. Let $B(S)$ be the Banach space of all bounded real-valued functions on S with supremum norm. For $s \in S$ and $f \in B(S)$, we define an element $l_s f$ in $B(S)$ by $(l_s f)(t) = f(st)$ for each $t \in S$. Let X be a subspace of $B(S)$ containing 1. An element μ in X^* is said to be a *mean* on X if $\|\mu\| = \mu(1) = 1$. We often write $\mu_t(f(t))$ instead of $\mu(f)$ for $\mu \in X^*$ and $f \in X$. Let X be l_s -invariant, that is, $l_s(X) \subset X$ for each $s \in S$. A mean μ on X is said to be *left invariant* if $\mu(l_s f) = \mu(f)$ for each $s \in S$ and $f \in X$. A sequence $\{\mu_n\}$ of means on X is said to be *strongly left regular* if $\|\mu_n - l_s^* \mu_n\| \rightarrow 0$ for each $s \in S$, where l_s^* is the adjoint operator of l_s . In the case when S is commutative, a strongly left regular sequence is said to be *strongly regular* [9, 10]. Let E be a Banach space, let X be a subspace of $B(S)$ containing 1 and let μ be a mean on X . Let f be a mapping from S into E such that $\{f(t) : t \in S\}$ is contained in a weakly compact convex subset of E and the mapping $t \mapsto \langle f(t), x^* \rangle$ is in X for each $x^* \in E^*$. We know from [9, 18] that there exists a unique element $x_0 \in E$ such that $\langle x_0, x^* \rangle = \mu_t \langle f(t), x^* \rangle$ for all $x^* \in E^*$. Following [9], we denote such x_0 by $\int f(t) d\mu(t)$. Let C be a nonempty closed convex subset of a Banach space E . A family $\mathcal{S} = \{T(t) : t \in S\}$ is said to be a *nonexpansive semigroup* on C if it satisfies the following:

- (1) for each $t \in S$, $T(t)$ is a nonexpansive mapping from C into itself;
- (2) $T(ts) = T(t)T(s)$ for each $t, s \in S$.

We denote by $F(\mathcal{S})$ the set of common fixed points of $\mathcal{S}$, that is, $\bigcap_{t \in S} F(T(t))$. Let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that for each $x \in C$, $\{T(t)x : t \in S\}$ is contained in a weakly compact convex subset of C . Let X be a subspace of $B(S)$ with $1 \in X$ such that the mapping $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$, and let μ be a mean on X . Following [15], we also write $T_\mu x$ instead of $\int T(t)x d\mu(t)$ for $x \in C$. We remark that T_μ is nonexpansive on C and $T_\mu x = x$ for each $x \in F(\mathcal{S})$; for more details, see [19].

We write $x_n \rightarrow x$ (or $\lim_{n \rightarrow \infty} x_n = x$) to indicate that the sequence $\{x_n\}$ of vectors converges strongly to x . Similarly, we write $x_n \rightharpoonup x$ (or $w\text{-}\lim_{n \rightarrow \infty} x_n = x$) will symbolize weak convergence. For any element z and any set A , we denote the distance between z and A by $d(z, A) = \inf\{\|z - y\| : y \in A\}$.

3. Weak convergence theorem

Throughout the rest of this paper, we assume that S is a semigroup. Let C be a nonempty weakly compact convex subset of a Banach space E and let $\mathcal{S} = \{T(s) : s \in S\}$ be

a nonexpansive semigroup of C . We consider the following iterative procedure (see [21]): $x_0 = x \in C$ and

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{\mu_n} x_n \quad (3.1)$$

for every $n \in \mathbb{N}$, where $\{\alpha_n\}$ is a sequence in $(0, 1)$.

LEMMA 3.1. *Let C be a nonempty weakly compact convex subset of a Banach space E and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let X be a subspace of $B(S)$ with $1 \in X$ such that the function $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$. Let $\{\mu_n\}$ be a sequence of means on S and let $\{\alpha_n\}$ be a sequence of real numbers such that $0 < \alpha_n < 1$ for every $n \in \mathbb{N}$. Let $x \in C$ and let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{\mu_n} x_n \quad (3.2)$$

for every $n \in \mathbb{N}$. Then, $\|x_{n+1} - w\| \leq \|x_n - w\|$ and $\lim_{n \rightarrow \infty} \|x_n - w\|$ exists for each $w \in F(\mathcal{S})$.

Proof. Let $w \in F(\mathcal{S})$. By the definition of $\{x_n\}$, we obtain that

$$\begin{aligned} \|x_n - w\| &= \|\alpha_n(x_{n-1} - w) + (1 - \alpha_n)(T_{\mu_n} x_n - w)\| \\ &\leq \alpha_n \|x_{n-1} - w\| + (1 - \alpha_n) \|T_{\mu_n} x_n - w\| \\ &\leq \alpha_n \|x_{n-1} - w\| + (1 - \alpha_n) \|x_n - w\| \end{aligned} \quad (3.3)$$

and hence

$$\alpha_n \|x_n - w\| \leq \alpha_n \|x_{n-1} - w\|. \quad (3.4)$$

It follows from $\alpha_n \neq 0$ that $\{\|x_n - w\|\}$ is a nonincreasing sequence. Hence, it follows that $\lim_{n \rightarrow \infty} \|x_n - w\|$ exists. $\square$

The following lemma was proved by Shioji and Takahashi [16] (see also [3, 9]).

LEMMA 3.2 (Shioji and Takahashi). *Let C be a nonempty closed convex subset of a uniformly convex Banach space E and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C . Let X be a subspace of $B(S)$ with $1 \in X$ such that it is l_s -invariant for each $s \in S$, and the function $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$. Let $\{\mu_n\}$ be a sequence of means on S which is strongly left regular. For each $r > 0$ and $t \in S$,*

$$\lim_{n \rightarrow \infty} \sup_{y \in C \cap B_r} \|T_{\mu_n} y - T(t) T_{\mu_n} y\| = 0. \quad (3.5)$$

The following lemma is crucial in the proofs of the main theorems.

LEMMA 3.3. *Let C be a nonempty closed convex subset of a uniformly convex Banach space E and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let X be a subspace of $B(S)$ with $1 \in X$ such that it is l_s -invariant for each $s \in S$, and the function $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$. Let $\{\mu_n\}$ be a sequence of means on S*

which is strongly left regular and let $\{\alpha_n\}$ be a sequence of real numbers such that $0 < \alpha_n < 1$ for every $n \in \mathbb{N}$ and $\sum_{n=1}^{\infty} (1 - \alpha_n) = \infty$. Let $x \in C$ and let $\{x_n\}$ be the sequence defined by $x_0 = x$ and

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{\mu_n} x_n \quad (3.6)$$

for every $n \in \mathbb{N}$. Then, for each $t \in S$,

$$\lim_{n \rightarrow \infty} \|x_n - T(t)x_n\| = 0. \quad (3.7)$$

Proof. For $x \in C$ and $w \in F(\mathcal{S})$, put $r = \|x - w\|$ and set $D = \{u \in E : \|u - w\| \leq r\} \cap C$. Then, D is a nonempty bounded closed convex subset of C which is $T(s)$ -invariant for each $s \in S$ and contains $x_0 = x$. So, without loss of generality, we may assume that C is bounded. Fix $\varepsilon > 0$, $t \in S$ and set $M_0 = \sup\{\|z\| : z \in C\}$. Then, from Proposition 2.2, there exists $\delta > 0$ such that

$$\overline{CO}F_\delta(T(t)) \subset F_\varepsilon(T(t)). \quad (3.8)$$

From Lemma 3.2 there exists $l \in \mathbb{N}$ such that

$$\|T_{\mu_i} y - T(t)T_{\mu_i} y\| < \delta \quad (3.9)$$

for every $i \geq l$ and $y \in C$. We have, for each $k \in \mathbb{N}$,

$$\begin{aligned} x_{l+k} &= \alpha_{l+k} x_{l+k-1} + (1 - \alpha_{l+k}) T_{\mu_{l+k}} x_{l+k} \\ &= \alpha_{l+k} \{ \alpha_{l+k-1} x_{l+k-2} + (1 - \alpha_{l+k-1}) T_{\mu_{l+k-1}} x_{l+k-1} \} + (1 - \alpha_{l+k}) T_{\mu_{l+k}} x_{l+k} \\ &\quad \vdots \\ &= \left(\prod_{i=l}^{l+k} \alpha_i \right) x_{l-1} + \sum_{j=l}^{l+k-1} \left\{ \left(\prod_{i=j+1}^{l+k} \alpha_i \right) (1 - \alpha_j) T_{\mu_j} x_j \right\} + (1 - \alpha_{l+k}) T_{\mu_{l+k}} x_{l+k}. \end{aligned} \quad (3.10)$$

Put

$$y_k = \frac{1}{1 - \prod_{i=l}^{l+k} \alpha_i} \left\{ \sum_{j=l}^{l+k-1} \left\{ \left(\prod_{i=j+1}^{l+k} \alpha_i \right) (1 - \alpha_j) T_{\mu_j} x_j \right\} + (1 - \alpha_{l+k}) T_{\mu_{l+k}} x_{l+k} \right\}. \quad (3.11)$$

From

$$\sum_{j=l}^{l+k-1} \left\{ \left(\prod_{i=j+1}^{l+k} \alpha_i \right) (1 - \alpha_j) \right\} + (1 - \alpha_{l+k}) = 1 - \prod_{i=l}^{l+k} \alpha_i, \quad (3.12)$$

we obtain $y_k \in \text{co}(\{T_{\mu_i}x_i\}_{i=l}^{l+k})$ and

$$x_{l+k} = \left(\prod_{i=l}^{l+k} \alpha_i \right) x_{l-1} + \left(1 - \prod_{i=l}^{l+k} \alpha_i \right) y_k. \quad (3.13)$$

From (3.9), we know that for every $k \in \mathbb{N}$, $T_{\mu_i}x_i \in F_\delta(T(t))$ for $i = l, l+1, \dots, l+k$. So, it follows from (3.8) that $y_k \in \text{co}F_\delta(T(t)) \subset F_\varepsilon(T(t))$ for every $k \in \mathbb{N}$. We know from Abel-Dini theorem that $\sum_{i=l}^{\infty} (1 - \alpha_i) = \infty$ implies $\prod_{i=l}^{\infty} \alpha_i = 0$. Then, there exists $m \in \mathbb{N}$ such that $\prod_{i=l}^{l+k} \alpha_i < \varepsilon/(2M_0)$ for every $k \geq m$. From (3.13), we obtain

$$\|x_{l+k} - y_k\| = \left(\prod_{i=l}^{l+k} \alpha_i \right) \|x_{l-1} - y_k\| < \frac{\varepsilon}{2M_0} \cdot 2M_0 = \varepsilon \quad (3.14)$$

for every $k \geq m$. Hence,

$$\begin{aligned} \|T(t)x_{l+k} - x_{l+k}\| &\leq \|T(t)x_{l+k} - T(t)y_k\| + \|T(t)y_k - y_k\| + \|y_k - x_{l+k}\| \\ &\leq 2\|x_{l+k} - y_k\| + \|T(t)y_k - y_k\| \leq 2\varepsilon + \varepsilon = 3\varepsilon \end{aligned} \quad (3.15)$$

for every $k \geq m$. Since $\varepsilon > 0$ is arbitrary, we get $\lim_{n \rightarrow \infty} \|T(t)x_n - x_n\| = 0$ for each $t \in S$. $\square$

Now, we prove a weak convergence theorem for a nonexpansive semigroup in a Banach space.

THEOREM 3.4. *Let C be a nonempty closed convex subset of a uniformly convex Banach space E which satisfies Opial's condition and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let X be a subspace of $B(S)$ with $1 \in X$ such that it is l_s -invariant for each $s \in S$, and the function $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$. Let $\{\mu_n\}$ be a sequence of means on S which is strongly left regular and let $\{\alpha_n\}$ be a sequence of real numbers such that $0 < \alpha_n < 1$ for every $n \in \mathbb{N}$ and $\sum_{n=1}^{\infty} (1 - \alpha_n) = \infty$. Let $x \in C$ and let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{\mu_n} x_n \quad (3.16)$$

for every $n \in \mathbb{N}$. Then, $\{x_n\}$ converges weakly to an element of $F(\mathcal{S})$.

Proof. Since E is reflexive and $\{x_n\}$ is bounded, $\{x_n\}$ must contain a subsequence of $\{x_n\}$ which converges weakly to a point in C . Let $\{x_{n_i}\}$ and $\{x_{n_j}\}$ be two subsequences of $\{x_n\}$ which converge weakly to y and z , respectively. From Lemma 3.3 and Proposition 2.1, we know $y, z \in F(\mathcal{S})$. We will show $y = z$. Suppose $y \neq z$. Then from Lemma 3.1 and Opial's condition, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - y\| &= \lim_{i \rightarrow \infty} \|x_{n_i} - y\| < \lim_{i \rightarrow \infty} \|x_{n_i} - z\| \\ &= \lim_{n \rightarrow \infty} \|x_n - z\| = \lim_{j \rightarrow \infty} \|x_{n_j} - z\| \\ &< \lim_{j \rightarrow \infty} \|x_{n_j} - y\| = \lim_{j \rightarrow \infty} \|x_n - y\|. \end{aligned} \quad (3.17)$$

This is a contradiction. Hence $\{x_n\}$ converges weakly to an element of $F(\mathcal{S})$. $\square$

4. Strong convergence theorems

In this section, we discuss the strong convergence of the iterates defined by (3.1). Now, we can prove a strong convergence theorem for a nonexpansive semigroup in a Banach space (see also [2]).

THEOREM 4.1. *Let C be a nonempty closed convex subset of a uniformly convex Banach space E and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let X be a subspace of $B(S)$ with $1 \in X$ such that it is l_s -invariant for each $s \in S$, and the function $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$. Let $\{\mu_n\}$ be a sequence of means on S which is strongly left regular and let $\{\alpha_n\}$ be a sequence of real numbers such that $0 < \alpha_n < 1$ for every $n \in \mathbb{N}$ and $\sum_{n=1}^{\infty} (1 - \alpha_n) = \infty$. Let $x \in C$ and let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{\mu_n} x_n \quad (4.1)$$

for every $n \in \mathbb{N}$. If there exists some $T(s) \in \mathcal{S}$ which is semicompact, then $\{x_n\}$ converges strongly to an element of $F(\mathcal{S})$.

Proof. Since the nonexpansive mapping $T(s)$ is semicompact, there exist a subsequence $\{x_{n_j}\}$ of $\{x_n\}$ and $y \in C$ such that $x_{n_j} \rightarrow y$ as $j \rightarrow \infty$. By Lemma 3.3, we have that

$$0 = \lim_{j \rightarrow \infty} \|x_{n_j} - T(t)x_{n_j}\| = \|y - T(t)y\| \quad (4.2)$$

for each $t \in S$ and hence $y \in F(\mathcal{S})$. Then, it follows from Lemma 3.1 that

$$\lim_{n \rightarrow \infty} \|x_n - y\| = \lim_{j \rightarrow \infty} \|x_{n_j} - y\| = 0. \quad (4.3)$$

Therefore, $\{x_n\}$ converges strongly to an element of $F(\mathcal{S})$. □

Next, we give a necessary and sufficient condition for the strong convergence of the iterates.

THEOREM 4.2. *Let C be a nonempty weakly compact convex subset of a Banach space E and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let X be a subspace of $B(S)$ with $1 \in X$ such that the function $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$. Let $\{\mu_n\}$ be a sequence of means on S and let $\{\alpha_n\}$ be a sequence of real numbers such that $0 < \alpha_n < 1$ for every $n \in \mathbb{N}$. Let $x \in C$ and let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{\mu_n} x_n \quad (4.4)$$

for every $n \in \mathbb{N}$. Then, $\{x_n\}$ converges strongly to a common fixed point of $\mathcal{S}$ if and only if $\lim_{n \rightarrow \infty} d(x_n, F(\mathcal{S})) = 0$.

Proof. The necessity is obvious. So, we will prove the sufficiency. Assume

$$\lim_{n \rightarrow \infty} d(x_n, F(\mathcal{S})) = 0. \quad (4.5)$$

By Lemma 3.1, we have

$$\|x_{n+1} - w\| \leq \|x_n - w\| \quad (4.6)$$

for each $w \in F(\mathcal{S})$. Taking the infimum over $w \in F(\mathcal{S})$,

$$d(x_{n+1}, F(\mathcal{S})) \leq d(x_n, F(\mathcal{S})) \quad (4.7)$$

and hence the sequence $\{d(x_n, F(\mathcal{S}))\}$ is nonincreasing. So, from $\lim_{n \rightarrow \infty} d(x_n, F(\mathcal{S})) = 0$, we obtain that

$$\lim_{n \rightarrow \infty} d(x_n, F(\mathcal{S})) = 0. \quad (4.8)$$

We will show that $\{x_n\}$ is a Cauchy sequence. Let $\varepsilon > 0$. There exists a positive integer N such that for each $n \geq N$, $d(x_n, F(\mathcal{S})) < \varepsilon/2$. For any $l, k \geq N$ and $w \in F(\mathcal{S})$, we obtain

$$\|x_l - w\| \leq \|x_N - w\|, \quad \|x_k - w\| \leq \|x_N - w\| \quad (4.9)$$

by Lemma 3.1. So, we obtain $\|x_l - x_k\| \leq \|x_l - w\| + \|w - x_k\| \leq 2\|x_N - w\|$ and hence

$$\|x_l - x_k\| \leq 2 \inf \{\|x_N - y\| : y \in F(\mathcal{S})\} = 2d(x_N, F(\mathcal{S})) < \varepsilon \quad (4.10)$$

for every $l, k \geq N$. This implies that $\{x_n\}$ is a Cauchy sequence. Since C is a closed subset of E , $\{x_n\}$ converges strongly to $z_0 \in C$. Further, since $F(\mathcal{S})$ is a closed subset of C , (4.8) implies that $z_0 \in F(\mathcal{S})$. Thus, we have that $\{x_n\}$ converges strongly to a common fixed point of $\mathcal{S}$. $\square$

THEOREM 4.3. *Let C be a nonempty closed convex subset of a uniformly convex Banach space E and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let X be a subspace of $B(S)$ with $1 \in X$ such that it is l_s -invariant for each $s \in S$, and the function $t \mapsto \langle T(t)x, x^* \rangle$ is in X for each $x \in C$ and $x^* \in E^*$. Let $\{\mu_n\}$ be a sequence of means on S which is strongly left regular and let $\{\alpha_n\}$ be a sequence of real numbers such that $0 < \alpha_n < 1$ for every $n \in \mathbb{N}$ and $\sum_{n=1}^{\infty} (1 - \alpha_n) = \infty$. Assume that there exist $s \in S$ and $k > 0$ such that*

$$\|(I - T(s))z\| \geq kd(z, F(\mathcal{S})) \quad (4.11)$$

for every $z \in C$. Let $x \in C$ and let $\{x_n\}$ be the sequence defined by $x_0 = x$ and

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{\mu_n} x_n \quad (4.12)$$

for every $n \in \mathbb{N}$. Then, $\{x_n\}$ converges strongly to an element of $F(\mathcal{F})$.

Proof. From Lemma 3.3, we obtain that $\|(I - T(s))x_n\| \rightarrow 0$ as $n \rightarrow \infty$. Then, it follows from (4.11) that

$$\lim_{n \rightarrow \infty} kd(x_n, F(\mathcal{F})) = 0 \quad (4.13)$$

for some $k > 0$. Therefore, we can conclude that $\{x_n\}$ converges strongly to an element of $F(\mathcal{F})$ by Theorem 4.2. $\square$

5. Deduced theorems from main results

Throughout this section, we assume that C is a nonempty closed convex subset of a uniformly convex Banach space E , x is an element of C , and $\{\alpha_n\}$ is a sequence of real numbers such that $0 < \alpha_n < 1$ for each $n \in \mathbb{N}$ and $\sum_{n=1}^{\infty} (1 - \alpha_n) = \infty$. As direct consequences of Theorems 3.4 and 4.1, we can show some convergence theorems.

THEOREM 5.1. *Let T be a nonexpansive mapping from C into itself such that $F(T) \neq \emptyset$. Let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) \frac{1}{n+1} \sum_{i=0}^n T^i x_n \quad (5.1)$$

for every $n \in \mathbb{N}$. If E satisfies Opial's condition, then $\{x_n\}$ converges weakly to a fixed point of T , and if T is semicompact, then $\{x_n\}$ converges strongly to a fixed point of T .

THEOREM 5.2. *Let T be as in Theorem 5.1. Let $\{s_n\}$ be a sequence of positive real numbers with $s_n \uparrow 1$. Let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) \sum_{i=0}^{\infty} s_n^i T^i x_n \quad (5.2)$$

for every $n \in \mathbb{N}$. If E satisfies Opial's condition, then $\{x_n\}$ converges weakly to a fixed point of T , and if T is semicompact, then $\{x_n\}$ converges strongly to a fixed point of T .

THEOREM 5.3. *Let T be as in Theorem 5.1. Let $\{q_{n,m} : n, m \in \mathbb{Z}^+\}$ be a sequence of real numbers such that $q_{n,m} \geq 0$, $\sum_{m=0}^{\infty} q_{n,m} = 1$ for every $n \in \mathbb{Z}^+$ and $\lim_{n \rightarrow \infty} \sum_{m=0}^{\infty} |q_{n,m+1} - q_{n,m}| = 0$. Let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) \sum_{m=0}^{\infty} q_{n,m} T^m x_n \quad (5.3)$$

for every $n \in \mathbb{N}$. If E satisfies Opial's condition, then $\{x_n\}$ converges weakly to a fixed point of T , and if T is semicompact, then $\{x_n\}$ converges strongly to a fixed point of T .

THEOREM 5.4. *Let T and U be commutative nonexpansive mappings from C into itself such that $F(T) \cap F(U) \neq \emptyset$. Let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) \frac{1}{(n+1)^2} \sum_{i,j=0}^n T^i U^j x_n \quad (5.4)$$

for every $n \in \mathbb{N}$. If E satisfies Opial's condition, then $\{x_n\}$ converges weakly to a common fixed point of T and U , and if either T or U is semicompact, then $\{x_n\}$ converges strongly to a common fixed point of T and U .

Let C be a closed convex subset of a Banach space E and let $\mathcal{S} = \{T(t) : t \in [0, \infty)\}$ be a family of nonexpansive mappings of C into itself. Then, $\mathcal{S}$ is called a one-parameter nonexpansive semigroup on C if it satisfies the following conditions: $T(0) = I$, $T(t+s) = T(t)T(s)$ for all $t, s \in [0, \infty)$ and $T(t)x$ is continuous in $t \in [0, \infty)$ for each $x \in C$.

THEOREM 5.5. *Let $\mathcal{S} = \{T(t) : t \in [0, \infty)\}$ be a one-parameter nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let $\{s_n\}$ be a sequence of positive real numbers with $s_n \rightarrow \infty$. Let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) \frac{1}{s_n} \int_0^{s_n} T(t)x_n dt \quad (5.5)$$

for every $n \in \mathbb{N}$. If E satisfies Opial's condition, then $\{x_n\}$ converges weakly to a common fixed point of $\mathcal{S}$, and if there exists some $T(s) \in \mathcal{S}$ which is semicompact, then $\{x_n\}$ converges strongly to a common fixed point of $\mathcal{S}$.

THEOREM 5.6. *Let $\mathcal{S}$ be as in Theorem 5.5. Let $\{r_n\}$ be a sequence of positive real numbers with $r_n \rightarrow 0$. Let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) r_n \int_0^\infty e^{-r_n t} T(t)x_n dt \quad (5.6)$$

for every $n \in \mathbb{N}$. If E satisfies Opial's condition, then $\{x_n\}$ converges weakly to a common fixed point of $\mathcal{S}$, and if there exists some $T(s) \in \mathcal{S}$ which is semicompact, then $\{x_n\}$ converges strongly to a common fixed point of $\mathcal{S}$.

THEOREM 5.7. *Let $\mathcal{S}$ be as in Theorem 5.5. Let $\{q_n\}$ be a sequence of continuous functions from $[0, \infty)$ into $[0, \infty)$ such that $\int_0^\infty q_n(t)dt = 1$ for every $n \in \mathbb{N}$, $\lim_{n \rightarrow \infty} q_n(t) = 0$ for $t \geq 0$ and $\lim_{n \rightarrow \infty} \int_0^\infty |q_n(t+s) - q_n(t)|dt = 0$ for all $s \geq 0$. Let $\{x_n\}$ be the sequence defined by $x_0 = x$ and*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) \int_0^\infty q_n(t)T(t)x_n dt \quad (5.7)$$

for every $n \in \mathbb{N}$. If E satisfies Opial's condition, then $\{x_n\}$ converges weakly to a common fixed point of $\mathcal{S}$, and if there exists some $T(s) \in \mathcal{S}$ which is semicompact, then $\{x_n\}$ converges strongly to a common fixed point of $\mathcal{S}$.

Acknowledgments

This research was supported by Grant-in-Aid for Young Scientists (B), the Ministry of Education, Culture, Sports, Science and Technology, Japan, and Grant-in-Aid for Scientific Research, Japan Society for the Promotion of Science.

References

- [1] S. Atsushiba, *Strong convergence theorems for finite quasi-nonexpansive mappings*, Comm. Appl. Nonlinear Anal. **9** (2002), no. 3, 57–68.
- [2] ———, *Strong convergence theorems for finite nonexpansive mappings in Banach spaces*, Proc. 3rd International Conference on Nonlinear Analysis and Convex Analysis (W. Takahashi and T. Tanaka, eds.), Yokohama Publishers, Yokohama, 2004, pp. 9–16.
- [3] S. Atsushiba, N. Shioji, and W. Takahashi, *Approximating common fixed points by the Mann iteration procedure in Banach spaces*, J. Nonlinear Convex Anal. **1** (2000), no. 3, 351–361.
- [4] F. E. Browder, *Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces*, Arch. Ration. Mech. Anal. **24** (1967), 82–90.
- [5] ———, *Nonlinear operators and nonlinear equations of evolution in Banach spaces*, Nonlinear Functional Analysis (Proc. Sympos. Pure Math., Vol. XVIII, Part 2, Chicago, Ill, 1968), American Mathematical Society, Rhode Island, 1976, pp. 1–308.
- [6] R. E. Bruck, *A simple proof of the mean ergodic theorem for nonlinear contractions in Banach spaces*, Israel J. Math. **32** (1979), no. 2-3, 107–116.
- [7] M. K. Ghosh and L. Debnath, *Convergence of Ishikawa iterates of quasi-nonexpansive mappings*, J. Math. Anal. Appl. **207** (1997), no. 1, 96–103.
- [8] J.-P. Gossez and E. Lami Dozo, *Some geometric properties related to the fixed point theory for nonexpansive mappings*, Pacific J. Math. **40** (1972), 565–573.
- [9] N. Hirano, K. Kido, and W. Takahashi, *Nonexpansive retractions and nonlinear ergodic theorems in Banach spaces*, Nonlinear Anal. **12** (1988), no. 11, 1269–1281.
- [10] G. G. Lorentz, *A contribution to the theory of divergent sequences*, Acta Math. **80** (1948), 167–190.
- [11] Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
- [12] W. V. Petryshyn, *Construction of fixed points of demicompact mappings in Hilbert space*, J. Math. Anal. Appl. **14** (1966), no. 2, 276–284.
- [13] W. V. Petryshyn and T. E. Williamson Jr., *Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings*, J. Math. Anal. Appl. **43** (1973), no. 2, 459–497.
- [14] S. Reich, *Strong convergence theorems for resolvents of accretive operators in Banach spaces*, J. Math. Anal. Appl. **75** (1980), no. 1, 287–292.
- [15] G. Rodé, *An ergodic theorem for semigroups of nonexpansive mappings in a Hilbert space*, J. Math. Anal. Appl. **85** (1982), no. 1, 172–178.
- [16] N. Shioji and W. Takahashi, *Strong convergence theorems for asymptotically nonexpansive semigroups in Banach spaces*, J. Nonlinear Convex Anal. **1** (2000), no. 1, 73–87.
- [17] Z.-H. Sun, C. He, and Y.-Q. Ni, *Strong convergence of an implicit iteration process for nonexpansive mappings in Banach spaces*, Nonlinear Funct. Anal. Appl. **8** (2003), no. 4, 595–602.
- [18] W. Takahashi, *A nonlinear ergodic theorem for an amenable semigroup of nonexpansive mappings in a Hilbert space*, Proc. Amer. Math. Soc. **81** (1981), no. 2, 253–256.
- [19] ———, *Nonlinear Functional Analysis*, Yokohama Publishers, Yokohama, 2000.
- [20] W. Takahashi and Y. Ueda, *On Reich's strong convergence theorems for resolvents of accretive operators*, J. Math. Anal. Appl. **104** (1984), no. 2, 546–553.

- [21] H.-K. Xu and R. G. Ori, *An implicit iteration process for nonexpansive mappings*, Numer. Funct. Anal. Optim. **22** (2001), no. 5-6, 767–773.
- [22] D. van Dulst, *Equivalent norms and the fixed point property for nonexpansive mappings*, J. London Math. Soc. (2) **25** (1982), no. 1, 139–144.

Sachiko Atsushiba: Department of Mathematics, Shibaura Institute of Technology, Fukasaku, Minuma-ku, Saitama-City, Saitama 337-8570, Japan

E-mail address: atusiba@sic.shibaura-it.ac.jp

Wataru Takahashi: Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, O-okayama, Meguro-ku, Tokyo 152-8552, Japan

E-mail address: wataru@is.titech.ac.jp

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskkklai@cityu.edu.hk