# APPROXIMATING COMMON FIXED POINTS OF TWO ASYMPTOTICALLY QUASI-NONEXPANSIVE MAPPINGS IN BANACH SPACES

NASEER SHAHZAD AND ANIEFIOK UDOMENE

*Received 21 April 2005; Revised 13 July 2005; Accepted 18 July 2005*

Suppose $K$ is a nonempty closed convex subset of a real Banach space $E$. Let $S, T : K \to K$ be two asymptotically quasi-nonexpansive maps with sequences $\{u_n\}, \{v_n\} \subset [0, \infty)$ such that $\sum_{n=1}^{\infty} u_n < \infty$ and $\sum_{n=1}^{\infty} v_n < \infty$, and $F = F(S) \cap F(T) := \{x \in K : Sx = Tx = x\} \neq \emptyset$. Suppose $\{x_n\}$ is generated iteratively by $x_1 \in K$, $x_{n+1} = (1 - \alpha_n)x_n + \alpha_n S^n[(1 - \beta_n)x_n + \beta_n T^n x_n]$, $n \geq 1$, where $\{\alpha_n\}$ and $\{\beta_n\}$ are real sequences in $[0, 1]$. It is proved that (a) $\{x_n\}$ converges strongly to some $x^* \in F$ if and only if $\liminf_{n\to\infty} d(x_n, F) = 0$; (b) if $X$ is uniformly convex and if either $T$ or $S$ is compact, then $\{x_n\}$ converges strongly to some $x^* \in F$. Furthermore, if $X$ is uniformly convex, either $T$ or $S$ is compact and $\{x_n\}$ is generated by $x_1 \in K$, $x_{n+1} = \alpha_n x_n + \beta_n S^n[\alpha'_n x_n + \beta'_n T^n x_n + \gamma'_n z'_n] + \gamma_n z_n$, $n \geq 1$, where $\{z_n\}$, $\{z'_n\}$ are bounded, $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\alpha'_n\}$, $\{\beta'_n\}$, $\{\gamma'_n\}$ are real sequences in $[0, 1]$ such that $\alpha_n + \beta_n + \gamma_n = 1 = \alpha'_n + \beta'_n + \gamma'_n$ and $\{\gamma_n\}$, $\{\gamma'_n\}$ are summable; it is established that the sequence $\{x_n\}$ (with error member terms) converges strongly to some $x^* \in F$.

Copyright © 2006 N. Shahzad and A. Udomene. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let $K$ be a nonempty subset of a real normed linear space $E$. Let $T$ be a self-mapping of $K$. Then $T$ is said to be *asymptotically nonexpansive* with sequence $\{v_n\} \subset [0, \infty)$ if $\lim_{n\to\infty} v_n = 0$ and

$$||T^n x - T^n y|| \leq (1 + v_n)\|x - y\| \tag{1.1}$$

for all $x, y \in K$ and $n \geq 1$; and is said to be *asymptotically quasi-nonexpansive* with sequence $\{v_n\} \subset [0, \infty)$ if $F(T) := \{x \in K : Tx = x\} \neq \emptyset$, $\lim_{n\to\infty} v_n = 0$ and

$$||T^n x - x^*|| \leq (1 + v_n)\|x - x^*\| \tag{1.2}$$

Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2006, Article ID 18909, Pages 1–10
DOI 10.1155/FPTA/2006/18909

for all $x \in K$, $x^* \in F(T)$ and $n \geq 1$. The mapping $T$ is called *nonexpansive* if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in K$; and is called *quasi-nonexpansive* if $F(T) \neq \emptyset$ and $\|Tx - x^*\| \leq \|x - x^*\|$ for all $x \in K$ and $x^* \in F(T)$. It is therefore clear that a nonexpansive mapping with a nonempty fixed point set is quasi-nonexpansive and an asymptotically nonexpansive mapping with a nonempty fixed point set is asymptotically quasi-nonexpansive. The converses do not hold in general. The mapping $T$ is called *uniformly* $(L, \gamma)$*-Lipschitzian* if there exists a constant $L > 0$ and $\gamma > 0$ such that

$$\|T^n x - T^n y\| \leq L\|x - y\|^{\gamma} \tag{1.3}$$

for all $x, y \in K$ and $n \geq 1$.

The class of asymptotically nonexpansive maps was introduced by Goebel and Kirk [3] as an important generalization of the class of nonexpansive maps. They established that if $K$ is a nonempty closed convex bounded subset of a uniformly convex Banach space $E$ and $T$ is an asymptotically nonexpansive self-mapping of $K$, then $T$ has a fixed point. In [4], they extended this result to the broader class of uniformly $(L, 1)$-Lipschitzian mappings with $L < \lambda$, where $\lambda$ is sufficiently near 1.

Iterative techniques for approximating fixed points of nonexpansive mappings and their generalizations (asymptotically nonexpansive mappings, etc.) have been studied by a number of authors (see, e.g., [1, 12–15] and references cited therein), using the *Mann iteration method* (see, e.g., [7]) or the *Ishikawa-type iteration method* (see, e.g., [5]).

In 1973, Petryshyn and Williamson [8] established a necessary and sufficient condition for a Mann iterative sequence to converge strongly to a fixed point of a quasi-nonexpansive mapping. Subsequently, Ghosh and Debnath [2] extended Petryshyn and Williamson's results and obtained some necessary and sufficient conditions for an Ishikawa-type iterative sequence to converge to a fixed point of a quasi-nonexpansive mapping. Recently, in [9, 10], Qihou extended the results of Ghosh and Debnath to the more general asymptotically quasi-nonexpansive mappings. More precisely, he obtained the following result.

Theorem 1.1 [9, Theorem 1, page 2]. *Let $K$ be a nonempty closed convex subset of a Banach space $E$ and $T : K \to K$ an asymptotically quasi-nonexpansive mapping with sequence $\{v_n\} \subset [0, \infty)$ such that $\sum_{n=1}^{\infty} v_n < \infty$, and $F(T) \neq \emptyset$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be real sequences in $[0, 1]$. Then the sequence $\{x_n\}$ generated from arbitrary $x_1 \in K$ by*

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n[(1 - \beta_n)x_n + \beta_n T^n x_n], \quad n \geq 1, \tag{1.4}$$

*converges strongly to some fixed point of $T$ if and only if* $\liminf_{n\to\infty} d(x_n, F(T)) = 0$, *here $d(y, C)$ denotes the distance of $y$ to a set $C$, that is, $d(y, C) = \inf\{d(y, x) : x \in C\}$.*

Furthermore, in [11], Qihou also established sufficient conditions for the strong convergence of the Ishikawa-type iterative sequences with error member for a uniformly $(L, \gamma)$-Lipschitzian asymptotically nonexpansive self-mapping of a nonempty *compact convex* subset of a uniformly convex Banach space. In [6], Khan and Takahashi studied

the problem of approximating common fixed points of two asymptotically nonexpansive mappings and obtained the following result.

Theorem 1.2 [6, Theorem 2, page 147]. *Let $E$ be a uniformly convex Banach space and $K$ a nonempty compact convex subset of $E$. Let $S, T : K \to K$ be two asymptotically nonexpansive mappings with sequence $\{k_n - 1\} \subset [0, \infty)$ such that $\sum_{n=1}^{\infty}(k_n - 1) < \infty$, and $F(S) \cap F(T) \neq \emptyset$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be real sequences in $[\epsilon, 1-\epsilon]$ for some $\epsilon \in (0,1)$. Then the sequence $\{x_n\}$ generated from arbitrary $x_1 \in K$ by*

$$x_{n+1} = (1-\alpha_n)x_n + \alpha_n S^n[(1-\beta_n)x_n + \beta_n T^n x_n], \quad n \geq 1 \tag{1.5}$$

*converges strongly to some common fixed point of $S$ and $T$.*

The purpose of this paper is to establish:

(i) necessary and sufficient conditions for the convergence of the Ishikawa-type iterative sequences involving two asymptotically quasi-nonexpansive mappings to a common fixed point of the mappings defined on a nonempty *closed convex* subset of a Banach space, and

(ii) a sufficient condition for the convergence of the Ishikawa-type iterative sequences involving two *uniformly continuous* asymptotically quasi-nonexpansive mappings to a common fixed point of the mappings defined on a nonempty *closed convex* subset of a uniformly convex Banach space. Further, we establish, as corollaries, the cases with error member terms. Our results are significant generalizations of the corresponding results of Ghosh and Debnath [2], Petryshyn and Williamson [8], Qihou [9–11], and of Khan and Takahashi [6].

## 2. Preliminaries

In what follows, we will make use of the following lemmas.

Lemma 2.1 (see, e.g., [13]). *Let $E$ be a uniformly convex Banach space and $\{\alpha_n\}$ a sequence in $[\epsilon, 1-\epsilon]$ for some $\epsilon \in (0,1)$. Suppose $\{x_n\}$ and $\{y_n\}$ are sequences in $E$ such that $\limsup_{n\to\infty}\|x_n\| \leq r$, $\limsup_{n\to\infty}\|y_n\| \leq r$, and $\limsup_{n\to\infty}\|\alpha_n x_n + (1-\alpha_n)y_n\| = r$ hold for some $r \geq 0$. Then $\lim_{n\to\infty}\|x_n - y_n\| = 0$.*

Lemma 2.2 (see, e.g., [16]). *Let $p > 1$ and $R > 1$ be two fixed numbers and $E$ a Banach space. Then $E$ is uniformly convex if and only if there exists a continuous, strictly increasing, and convex function $g : [0,\infty) \to [0,\infty)$ with $g(0) = 0$ such that $\|\lambda x + (1-\lambda)y\|^p \leq \lambda\|x\|^p + (1-\lambda)\|y\|^p - W_p(\lambda)g(\|x-y\|)$ for all $x, y \in B_R(0) = \{x \in E : \|x\| \leq R\}$, and $\lambda \in [0,1]$, where $W_p(\lambda) = \lambda(1-\lambda)^p + \lambda^p(1-\lambda)$.*

Lemma 2.3 (see, e.g., [14]). *Let $\{\lambda_n\}$ and $\{\sigma_n\}$ be sequences of nonnegative real numbers such that $\lambda_{n+1} \leq \lambda_n + \sigma_n$, $\forall n \geq 1$ and $\sum_{n=1}^{\infty}\sigma_n < \infty$. Then $\lim_{n\to\infty}\lambda_n$ exists. Moreover, if there exists a subsequence $\{\lambda_{n_j}\}$ of $\{\lambda_n\}$ such that $\lambda_{n_j} \to 0$ as $j \to \infty$, then $\lambda_n \to 0$ as $n \to \infty$.*

## 3. Main results

Let $K$ be a nonempty closed convex subset of a real Banach space $E$. Let $S, T : K \to K$ be two asymptotically quasi-nonexpansive mappings. The following iteration scheme is studied:

$$x_{n+1} = (1-\alpha_n)x_n + \alpha_n S^n[(1-\beta_n)x_n + \beta_n T^n x_n], \tag{3.1}$$

with $x_1 \in K$, $n \geq 1$, where $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences in $[0,1]$.

Theorem 3.1. *Let $E$ be a real Banach space and $K$ a nonempty closed convex subset of $E$. Let $S, T : K \to K$ be two asymptotically quasi-nonexpansive mappings with sequences $\{u_n\}, \{v_n\} \subset [0,\infty)$ such that $\sum_{n=1}^{\infty} u_n < \infty$ and $\sum_{n=1}^{\infty} v_n < \infty$, and $F = F(S) \cap F(T) := \{x \in K : Sx = Tx = x\} \neq \emptyset$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $[0,1]$. Starting from arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ by the recursion (3.1). Then*

(1) *$\|x_{n+1} - x^*\| \leq (1+b_n)\|x_n - x^*\|$ for all $n \geq 1$, $x^* \in F$, and for some sequence $\{b_n\}$ of numbers with $\sum_{n=1}^{\infty} b_n < \infty$.*

(2) *There exists a constant $M > 0$ such that $\|x_{n+m} - x^*\| \leq M\|x_n - x^*\|$ for all $n, m \geq 1$ and $x^* \in F$.*

*Proof.* (1) Let $x^* \in F$ and $y_n = (1-\beta_n)x_n + \beta_n T^n x_n$. Then

$$\begin{aligned} ||x_{n+1} - x^*|| &= ||(1-\alpha_n)x_n + \alpha_n S^n y_n - x^*|| \\ &\leq (1-\alpha_n)||x_n - x^*|| + \alpha_n(1+u_n)||y_n - x^*||, \\ ||y_n - x^*|| &= ||(1-\beta_n)x_n + \beta_n T^n x_n - x^*|| \\ &\leq (1-\beta_n)||x_n - x^*|| + \beta_n(1+v_n)||x_n - x^*|| \\ &\leq (1+v_n)||x_n - x^*||. \end{aligned} \tag{3.2}$$

Using (3.2), we obtain

$$\begin{aligned} ||x_{n+1} - x^*|| &\leq [1 + \alpha_n(u_n + v_n) + \alpha_n u_n v_n]||x_n - x^*|| \\ &\leq [1 + u_n + v_n + u_n v_n]||x_n - x^*|| \\ &\leq (1+b_n)||x_n - x^*||, \end{aligned} \tag{3.3}$$

where $b_n = u_n + v_n + u_n v_n$ with $\sum_{n=1}^{\infty} b_n < \infty$.

(2) Notice that for any $n, m \geq 1$

$$\begin{aligned} ||x_{n+m} - x^*|| &\leq (1+b_{n+m-1})||x_{n+m-1} - x^*|| \\ &\leq \exp(b_{n+m-1})||x_{n+m-1} - x^*|| \\ &\leq \cdots \leq \exp\left(\sum_{k=n}^{n+m-1} b_k\right)||x_n - x^*||. \end{aligned} \tag{3.4}$$

Let $M = \exp(\sum_{k=1}^{\infty} b_k)$. Then $0 < M < \infty$ and

$$||x_{n+m} - x^*|| \leq M||x_n - x^*||. \tag{3.5}$$

□

Theorem 3.2. *Let $E$ be a real Banach space and $K$ a nonempty closed convex subset of $E$. Let $S, T : K \to K$ be two asymptotically quasi-nonexpansive mappings ($S$ and $T$ need not be continuous) with sequences $\{u_n\}, \{v_n\} \subset [0, \infty)$ such that $\sum_{n=1}^{\infty} u_n < \infty$ and $\sum_{n=1}^{\infty} v_n < \infty$, and $F = F(S) \cap F(T) := \{x \in K : Sx = Tx = x\} \neq \emptyset$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $[0, 1]$. From arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ by the recursion (3.1). Then $\{x_n\}$ converges strongly to some common fixed point of $S$ and $T$ if and only if $\liminf_{n\to\infty} d(x_n, F) = 0$.*

*Proof.* It suffices that we only prove the sufficiency. By Theorem 3.1, we have $\|x_{n+1} - x^*\| \leq (1 + b_n)\|x_n - x^*\|$ for all $n \geq 1$ and $x^* \in F$. Therefore, $d(x_{n+1}, F) \leq (1 + b_n)d(x_n, F)$. Since $\sum_{n=1}^{\infty} b_n < \infty$ and $\liminf_{n\to\infty} d(x_n, F) = 0$, it follows by Lemma 2.3 that $\lim_{n\to\infty} d(x_n, F) = 0$. Next we will show that $\{x_n\}$ is a Cauchy sequence. Since $\lim_{n\to\infty} d(x_n, F) = 0$, given any $\epsilon > 0$, there exists a natural number $n_0$ such that $d(x_n, F) < \epsilon/4M$ for all $n \geq n_0$. Here $M > 0$ is the constant in Theorem 3.1(2). So we can find $w^* \in F$ such that $\|x_{n_0} - w^*\| \leq \epsilon/3M$. Using Theorem 3.1(2), we have for all $n \geq n_0$ and $m \geq 1$ that

$$\begin{aligned} ||x_{n+m} - x_n|| &\leq ||x_{n+m} - w^*|| + ||x_n - w^*|| \\ &\leq M||x_{n_0} - w^*|| + M||x_{n_0} - w^*|| \\ &= 2M||x_{n_0} - w^*|| < \epsilon. \end{aligned} \tag{3.6}$$

This implies that $\{x_n\}$ is a Cauchy sequence and so is convergent, since $X$ is complete. Let $\lim_{n\to\infty} x_n = y^*$. Then $y^* \in K$. It remains to show that $y^* \in F$. Let $\hat{\epsilon} > 0$ be given. Then there exists a natural number $n_1$ such that $\|x_n - y^*\| < \hat{\epsilon}/2\max\{2 + u_1, 2 + v_1\}$ for all $n \geq n_1$. Since $\lim_{n\to\infty} d(x_n, F) = 0$, there exists a natural number $n_2 \geq n_1$ such that for all $n \geq n_2$ we have $d(x_n, F) < \hat{\epsilon}/3\max\{2 + u_1, 2 + v_1\}$ and in particular we have $d(x_{n_2}, F) < \hat{\epsilon}/3\max\{2 + u_1, 2 + v_1\}$. Therefore, there exists $z^* \in F$ such that $\|x_{n_2} - z^*\| \leq \hat{\epsilon}/2\max\{2 + u_1, 2 + v_1\}$. Consequently we have

$$\begin{aligned} \|Sy^* - y^*\| &= ||Sy^* - z^* + z^* - x_{n_2} + x_{n_2} - y^*|| \\ &\leq \|Sy^* - z^*\| + ||z^* - x_{n_2}|| + ||x_{n_2} - y^*|| \\ &\leq (1 + u_1)\|y^* - z^*\| + ||z^* - x_{n_2}|| + ||x_{n_2} - y^*|| \\ &\leq (2 + u_1)||y^* - x_{n_2}|| + (2 + u_1)||x_{n_2} - z^*|| \\ &< (2 + u_1)\frac{\hat{\epsilon}}{2\max\{2 + u_1, 2 + v_1\}} + (2 + u_1)\frac{\hat{\epsilon}}{2\max\{2 + u_1, 2 + v_1\}} \\ &\leq \hat{\epsilon}. \end{aligned} \tag{3.7}$$

This implies that $y^* \in F(S)$. Similarly, $y^* \in F(T)$. Hence $y^* \in F$. This completes the proof. □

Theorem 3.3. *Let $E$ be a real uniformly convex Banach space and $K$ a nonempty closed convex subset of $E$. Let $S, T : K \to K$ be two uniformly continuous asymptotically quasi-nonexpansive mappings with sequences $\{u_n\}, \{v_n\} \subset [0, \infty)$ such that $\sum_{n=1}^{\infty} u_n < \infty$ and $\sum_{n=1}^{\infty} v_n < \infty$, and $F = F(S) \cap F(T) := \{x \in K : Sx = Tx = x\} \neq \emptyset$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $[\epsilon, 1-\epsilon]$ for some $\epsilon \in (0,1)$. From arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ by the recursion (3.1). Then*

$$\lim_{n\to\infty} ||x_n - T^n x_n|| = 0 = \lim_{n\to\infty} ||x_n - S^n x_n||. \tag{3.8}$$

*Proof.* Let $x^* \in F$. Then, by Theorem 3.1(1) and Lemma 2.3, $\lim_{n\to\infty} \|x_n - x^*\|$ exists. Let $\lim_{n\to\infty} \|x_n - x^*\| = r$. If $r = 0$, then by the continuity of $S$ and $T$ the conclusion follows. Now suppose $r > 0$. We claim

$$\lim_{n\to\infty} ||S^n x_n - x_n|| = 0 = \lim_{n\to\infty} ||T^n x_n - x_n||. \tag{3.9}$$

Set $y_n = (1-\beta_n)x_n + \beta_n T^n x_n$. Since $\{x_n\}$ is bounded, there exists $R > 0$ such that $x_n - x^*, y_n - x^* \in B_R(0)$ for all $n \geq 1$. Using Lemma 2.2, we have that

$$\begin{aligned} ||y_n - x^*||^2 &= ||(1-\beta_n)x_n + \beta_n T^n x_n - x^*||^2 \\ &\leq \beta_n ||T^n x_n - x^*||^2 + (1-\beta_n)||x_n - x^*||^2 - W_2(\beta_n) g(||T^n x_n - x_n||) \\ &\leq \beta_n (1+v_n)^2 ||x_n - x^*||^2 + (1-\beta_n)||x_n - x^*||^2 \leq (1+v_n)^2 ||x_n - x^*||^2. \end{aligned} \tag{3.10}$$

From Lemma 2.2, it follows that

$$\begin{aligned} ||x_{n+1} - x^*||^2 &= ||(1-\alpha_n)x_n + \alpha_n S^n y_n - x^*||^2 \\ &\leq (1-\alpha_n)||x_n - x^*||^2 + \alpha_n (1+u_n)^2 ||y_n - x^*||^2 \\ &\quad - W_2(\alpha_n) g(||S^n y_n - x_n||) \\ &\leq (1-\alpha_n)||x_n - x^*||^2 + \alpha_n (1+u_n)^2 (1+v_n)^2 ||x_n - x^*||^2 \\ &\quad - W_2(\alpha_n) g(||S^n y_n - x_n||) \\ &\leq ||x_n - x^*||^2 + c_n R^2 - W_2(\alpha_n) g(||S^n y_n - x_n||), \end{aligned} \tag{3.11}$$

where $c_n = (1-\epsilon)[2(u_n + v_n) + (u_n^2 + 4u_n v_n + v_n^2) + 2(u_n v_n^2 + u_n^2 v_n) + v_n^2 v_n^2]$. Observe that $W_2(\alpha_n) \geq \epsilon^2$ and $\sum_{n=1}^{\infty} c_n < \infty$. Now (3.11) implies that

$$\epsilon^2 \sum_{n=1}^{\infty} g(||S^n y_n - x_n||) < ||x_1 - x^*||^2 + R^2 \sum_{n=1}^{\infty} c_n < \infty. \tag{3.12}$$

Therefore, we have $\lim_{n\to\infty} g(\|S^n y_n - x_n\|) = 0$. Since $g$ is strictly increasing and continuous at 0, it follows that

$$\lim_{n\to\infty} ||S^n y_n - x_n|| = 0. \tag{3.13}$$

Since $S$ is asymptotically quasi-nonexpansive, we can get that

$$||x_n - x^*|| \leq ||x_n - S^n y_n|| + (1+u_n)||y_n - x^*||, \tag{3.14}$$

from which we deduce that $r \leq \liminf_{n\to\infty} \|y_n - x^*\|$. On the other hand, we have

$$\begin{aligned} ||y_n - x^*|| &\leq ||(1-\beta_n)x_n + \beta_n T^n x_n - x^*|| \\ &= ||(1-\beta_n)(x_n - x^*) + \beta_n (T^n x_n - x^*)|| \\ &\leq (1-\beta_n)||x_n - x^*|| + (1+\nu_n)\beta_n ||x_n - x^*|| \\ &= ||x_n - x^*|| + \beta_n \nu_n ||x_n - x^*|| \leq (1+\nu_n)||x_n - x^*||, \end{aligned} \tag{3.15}$$

which implies $\limsup_{n\to\infty} \|y_n - x^*\| \leq r$. Therefore, $\lim_{n\to\infty} \|y_n - x^*\| = r$ and so

$$\lim_{n\to\infty} ||\beta_n (T^n x_n - x^*) + (1-\beta_n)(x_n - x^*)|| = r. \tag{3.16}$$

Since $\limsup_{n\to\infty} \|T^n x_n - x^*\| \leq r$, it follows from Lemma 2.1 that

$$\lim_{n\to\infty} ||T^n x_n - x_n|| = 0. \tag{3.17}$$

Also, we have

$$||S^n x_n - x_n|| \leq ||S^n x_n - S^n y_n|| + ||S^n y_n - x_n||. \tag{3.18}$$

Since $S$ is uniformly continuous and $\|x_n - y_n\| \to 0$ as $n \to \infty$, it follows from (3.18) that $\lim_{n\to\infty} \|S^n x_n - x_n\| = 0$. This completes the proof. □

Theorem 3.4. *Let $E$ be a real uniformly convex Banach space and $K$ a nonempty closed convex subset of $E$. Let $S, T : K \to K$ be two uniformly continuous asymptotically quasi-nonexpansive mappings with sequences $\{u_n\}, \{v_n\} \subset [0, \infty)$ such that $\sum_{n=1}^{\infty} u_n < \infty$ and $\sum_{n=1}^{\infty} v_n < \infty$, and $F = F(S) \cap F(T) := \{x \in K : Sx = Tx = x\} \neq \emptyset$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $[\epsilon, 1-\epsilon]$ for some $\epsilon \in (0,1)$. From arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ by the recursion (3.1). Assume, in addition, that either $T$ or $S$ is compact. Then $\{x_n\}$ converges strongly to some common fixed point of $S$ and $T$.*

*Proof.* By Theorem 3.3, we have

$$\lim_{n\to\infty} ||S^n x_n - x_n|| = 0 = \lim_{n\to\infty} ||T^n x_n - x_n|| \tag{3.19}$$

and also

$$\lim_{n\to\infty} ||x_n - S^n y_n|| = 0. \tag{3.20}$$

If $T$ is compact, then there exists a subsequence $\{T^{n_k} x_{n_k}\}$ of $\{T^n x_n\}$ such that $T^{n_k} x_{n_k} \to x^*$ as $k \to \infty$ for some $x^* \in K$ and so $T^{n_k+1} x_{n_k} \to Tx^*$ as $k \to \infty$. From (3.19), we have $x_{n_k} \to x^*$ as $k \to \infty$. Also $S^{n_k} y_{n_k} \to x^*$ as $k \to \infty$ by (3.20). Since $\|x_{n_k+1} - x_{n_k}\| \leq \|x_{n_k} - S^{n_k} y_{n_k}\|$,

it follows that $x_{n_k+1} \to x^*$ as $k \to \infty$. Again, from (3.20), we have $S^{n_k+1}y_{n_k+1} \to x^*$. Next we show that $x^* \in F$. Notice that

$$\begin{aligned} ||x^* - Tx^*|| \leq{}& ||x^* - x_{n_k+1}|| + ||x_{n_k+1} - T^{n_k+1}x_{n_k+1}|| \\ &+ ||T^{n_k+1}x_{n_k+1} - T^{n_k+1}x_{n_k}|| + ||T^{n_k+1}x_{n_k} - Tx^*||. \end{aligned} \tag{3.21}$$

Since $T$ is uniformly continuous, taking the limit as $k \to \infty$ and using (3.19), we obtain that $x^* = Tx^*$ and so $x^* \in F(T)$. Notice also that

$$\begin{aligned} ||x^* - Sx^*|| \leq{}& ||x^* - x_{n_k+1}|| + ||x_{n_k+1} - S^{n_k+1}x_{n_k+1}|| \\ &+ ||S^{n_k+1}x_{n_k+1} - S^{n_k+1}x_{n_k}|| + ||S^{n_k+1}x_{n_k} - Sx^*||. \end{aligned} \tag{3.22}$$

Letting $k \to \infty$, we also have that $x^* = Sx^*$ and so $x^* \in F(S)$. Thus $x^* \in F$. Hence, by Lemma 2.3, $x_n \to x^* \in F$ since $\lim_{n\to\infty} \|x_n - x^*\|$ exists. If $S$ is compact, then essentially the same arguments as above give the conclusion. This completes the proof. □

Corollary 3.5. *Let $E$ be a real uniformly convex Banach space and $K$ a nonempty compact convex subset of $E$. Let $S, T : K \to K$ be two continuous asymptotically quasi-nonexpansive mappings with sequences $\{u_n\}, \{v_n\} \subset [0,\infty)$ such that $\sum_{n=1}^{\infty} u_n < \infty$ and $\sum_{n=1}^{\infty} v_n < \infty$, and $F = F(S) \cap F(T) := \{x \in K : Sx = Tx = x\} \neq \emptyset$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $[\epsilon, 1-\epsilon]$ for some $\epsilon \in (0,1)$. From arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ by the recursion (3.1). Then $\{x_n\}$ converges strongly to some common fixed point of $S$ and $T$.*

Corollary 3.6. *Let $E$ be a real uniformly convex Banach space and $K$ a nonempty compact convex subset of $E$. Let $T : K \to K$ be a continuous asymptotically quasi-nonexpansive mapping with sequence $\{v_n\} \subset [0,\infty)$ such that $\sum_{n=1}^{\infty} v_n < \infty$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $[\epsilon, 1-\epsilon]$ for some $\epsilon \in (0,1)$. From arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ by the recursion*

$$x_{n+1} = (1-\alpha_n)x_n + \alpha_n T^n\left[(1-\beta_n)x_n + \beta_n T^n x_n\right] \tag{3.23}$$

*with $n \geq 1$. Then $\{x_n\}$ converges strongly to some fixed point of $T$.*

*Remarks.* (1) Corollary 3.5 extends Theorem 1.2 to the more general class of mappings considered in this paper. It is worth noting that Theorem 1.2 is proved for two asymptotically nonexpansive mappings having the same sequence $\{u_n\}$ (here $u_n = k_n - 1$). However, in our results $S$ and $T$ have separate sequences $\{u_n\}$ and $\{v_n\}$, respectively.

(2) Theorem 3.2 contains as special cases Theorem 1.1, the main result of Qihou [9], together with [9, Corollaries 1 and 2], which are themselves extensions of the results of Ghosh and Debnath [2] and Petryshyn and Williamson [8].

(3) Theorem 3.7 and Corollary 3.8 below are easily provable since the sequences $\{\gamma_n\}$, $\{\gamma_n'\}$ in $[0,1]$ are assumed summable and the sequences $\{z_n\}$, $\{z_n'\}$ in $K$ are bounded. Usually, once a convergence result has been established for an iteration scheme *without errors*, such as (1.4) or (3.1), it is not always difficult to establish the corresponding result for the case *with errors* such as the main theorem of [11] or Theorem 3.7 and Corollary 3.8 below, once $\{\gamma_n\}$, $\{\gamma_n'\}$ are assumed summable and the sequences of error terms are bounded.

Theorem 3.7. *Let $E$, $K$, $S$, $T$ and $F$ be as in Theorem 3.4. Let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\alpha'_n\}$, $\{\beta'_n\}$, and $\{\gamma'_n\}$ be sequences in $[0,1]$ with $\alpha_n+\beta_n+\gamma_n=\alpha'_n+\beta'_n+\gamma'_n=1$ for all $n\geq 1$. From arbitrary $x_1\in K$, define the sequence $\{x_n\}$ by*

$$\begin{aligned} x_{n+1} &= \alpha_n x_n+\beta_n S^n y_n+\gamma_n z_n,\\ y_n &= \alpha'_n x_n+\beta'_n T^n x_n+\gamma'_n z'_n, \end{aligned} \tag{3.24}$$

*where $\{z_n\}$ and $\{z'_n\}$ are bounded sequences in $K$. Suppose* (i) for some $\epsilon\in(0,1)$, $\beta_n+\gamma_n\in[\epsilon,1-\epsilon]$ and $\beta'_n+\gamma'_n\in[\epsilon,1-\epsilon]$ for all $n\geq 1$, and (ii) $\sum_{n=1}^{\infty}\gamma_n<\infty$, $\sum_{n=1}^{\infty}\gamma'_n<\infty$. Then $\{x_n\}$ converges strongly to some common fixed point of $S$ and $T$.

Corollary 3.8. *Let $E$, $K$, $S$, $T$ and $F$ be as in Corollary 3.5. Let $\{x_n\}$ be defined as in Theorem 3.7 and let the sequences $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\alpha'_n\}$, $\{\beta'_n\}$, and $\{\gamma'_n\}$ satisfy the same conditions as in Theorem 3.7. Then $\{x_n\}$ converges strongly to some common fixed point of $S$ and $T$.*

*Remarks.* (4) Corollary 3.8 extends the results of Qihou [11] to the more general class of *continuous* asymptotically quasi-nonexpansive mappings on a compact convex subset of a uniformly convex Banach space.

(5) In Theorems 3.3, 3.4 and Corollaries 3.5, 3.6, the prototypes for the sequence $\{\alpha_n\}$ and $\{\beta_n\}$ are $\alpha_n=1/2=\beta_n$ for all $n\geq 1$. In this case $\epsilon=1/4$ satisfies the conditions given therein.

(6) In Theorem 3.7 and Corollary 3.8, the prototypes for the sequences $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\alpha'_n\}$, $\{\beta'_n\}$, and $\{\gamma'_n\}$ are $\alpha_n=3/4-1/(n+1)^2=\alpha'_n$; $\beta_n=1/4=\beta'_n$; $\gamma_n=1/(n+1)^2=\gamma'_n$ for all $n\geq 1$. In this case, $\epsilon=1/4$ satisfies the conditions given therein.

(7) Theorems 3.1, 3.2, 3.3, 3.4, 3.7 and Corollaries 3.5, 3.6, 3.8 remain true for the subclass of asymptotically nonexpansive mappings.

**Acknowledgments**

The authors thank the referees for their useful comments. This work was done while the authors were visiting the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy. They would like to thank the Centre for hospitality and financial support.

**References**

[1] C. E. Chidume, *Iterative algorithms for nonexpansive mappings and some of their generalizations*, Nonlinear Analysis and Applications: to V. Lakshmikantham on His 80th Birthday. Vol. 1, 2 (R. P. Agarwal, et al., eds.), Kluwer Academic, Dordrecht, 2003, pp. 383–429.

[2] M. K. Ghosh and L. Debnath, *Convergence of Ishikawa iterates of quasi-nonexpansive mappings*, Journal of Mathematical Analysis and Applications **207** (1997), no. 1, 96–103.

[3] K. Goebel and W. A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proceedings of the American Mathematical Society **35** (1972), 171–174.

[4] ———, *A fixed point theorem for transformations whose iterates have uniform Lipschitz constant*, Polska Akademia Nauk. Instytut Matematyczny. Studia Mathematica **47** (1973), 135–140.

[5] S. Ishikawa, *Fixed points by a new iteration method*, Proceedings of the American Mathematical Society **44** (1974), 147–150.

[6] S. H. Khan and W. Takahashi, *Approximating common fixed points of two asymptotically nonexpansive mappings*, Scientiae Mathematicae Japonicae **53** (2001), no. 1, 143–148.
[7] W. R. Mann, *Mean value methods in iteration*, Proceedings of the American Mathematical Society **4** (1953), 506–510.
[8] W. V. Petryshyn and T. E. Williamson Jr., *Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings*, Journal of Mathematical Analysis and Applications **43** (1973), 459–497.
[9] L. Qihou, *Iterative sequences for asymptotically quasi-nonexpansive mappings*, Journal of Mathematical Analysis and Applications **259** (2001), no. 1, 1–7.
[10] ———, *Iterative sequences for asymptotically quasi-nonexpansive mappings with error member*, Journal of Mathematical Analysis and Applications **259** (2001), no. 1, 18–24.
[11] ———, *Iteration sequences for asymptotically quasi-nonexpansive mapping with an error member of uniform convex Banach space*, Journal of Mathematical Analysis and Applications **266** (2002), no. 2, 468–471.
[12] B. E. Rhoades, *Fixed point iterations for certain nonlinear mappings*, Journal of Mathematical Analysis and Applications **183** (1994), no. 1, 118–120.
[13] J. Schu, *Weak and strong convergence to fixed points of asymptotically nonexpansive mappings*, Bulletin of the Australian Mathematical Society **43** (1991), no. 1, 153–159.
[14] K.-K. Tan and H. K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, Journal of Mathematical Analysis and Applications **178** (1993), no. 2, 301–308.
[15] ———, *Fixed point iteration processes for asymptotically nonexpansive mappings*, Proceedings of the American Mathematical Society **122** (1994), no. 3, 733–739.
[16] H. K. Xu, *Inequalities in Banach spaces with applications*, Nonlinear Analysis. Theory, Methods & Applications. An International Multidisciplinary Journal. Series A: Theory and Methods **16** (1991), no. 12, 1127–1138.

Naseer Shahzad: Department of Mathematics, King Abdul Aziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia
*E-mail address*: nshahzad@kaau.edu.sa

Aniefiok Udomene: Department of Mathematics/Statistics/Computer Science, University of Port Harcourt, PMB 5323, Port Harcourt, Nigeria
*E-mail address*: epsilon_ani@yahoo.com

Mathematical Problems in Engineering

# Special Issue on
# Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

## Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at http://www.hindawi.com/journals/mpe/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/ according to the following timetable:

| | |
|---|---|
| Manuscript Due | December 1, 2008 |
| First Round of Reviews | March 1, 2009 |
| Publication Date | June 1, 2009 |

**Guest Editors**

**José Roberto Castilho Piqueira,** Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

**Elbert E. Neher Macau,** Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São Josè dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

**Celso Grebogi,** Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk