NOTE ON SOME RESULTS FOR ASYMPTOTICALLY PSEUDOCONTRACTIVE MAPPINGS AND ASYMPTOTICALLY NONEXPANSIVE MAPPINGS

YUCHAO TANG AND LIWEI LIU

Received 26 May 2006; Revised 5 August 2006; Accepted 13 August 2006

We discuss convergence theorems of modified Ishikawa and Mann iterative sequences with errors for asymptotically pseudocontractive and asymptotically nonexpansive mappings in Banach spaces, and the boundedness of the domain and range can be dropped, generalizing theorems of Chang.

Copyright © 2006 Y. Tang and L. Liu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction and preliminaries

Throughout this paper, we assume that E is a real Banach space, E^* is the topological dual space of E, $\langle\cdot,\cdot\rangle$ is the dual between E and E^*, $D(T)$ and $F(T)$ denote the domain of T and the set of all fixed points of T, respectively, and $J : E \to 2^{E^*}$ is the normalized duality mapping defined by

$$J(x) = \{f \in E^* : \langle x, f\rangle = \|x\| \cdot \|f\|,\ \|f\| = \|x\|\}, \quad x \in E. \tag{1.1}$$

Definition 1.1. Let $T : D(T) \subset E \to E$ be a mapping.

(1) T is said to be asymptotically nonexpansive if there exists a sequence $\{k_n\}$ in $(0,\infty)$ with $\lim_{n\to\infty} k_n = 1$ such that

$$||T^n x - T^n y|| \le k_n \|x - y\| \tag{1.2}$$

for all $x, y \in D(T)$ and $n = 1, 2, \ldots$.

(2) T is said to be asymptotically pseudocontractive if there exists a sequence $\{k_n\}$ in $(0,\infty)$ with $\lim_{n\to\infty} k_n = 1$, and for any $x, y \in D(T)$ there exists $j(x-y) \in J(x-y)$ such that

$$\langle T^n x - T^n y,\ j(x-y)\rangle \le k_n \|x - y\|^2 \tag{1.3}$$

for all $n = 1, 2, \ldots$.

Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2006, Article ID 24978, Pages 1–7
DOI 10.1155/FPTA/2006/24978

(3) T is said to be uniformly L-Lipschitzian if there exists $L > 0$ such that

$$||T^n x - T^n y|| \leq L\|x - y\| \tag{1.4}$$

for all $x, y \in D(T)$ and $n = 1, 2, \ldots$.

The following proposition follows from Definition 1.1 immediately.

Proposition 1.2. (1) *If $T : D(T) \subset E \to E$ is nonexpansive, then T is an asymptocially nonexpansive mapping with a constant sequence* $\{1\}$.

(2) *If $T : D(T) \subset E \to E$ is asymptotically nonexpansive, then T is a uniformly L-Lipschitzian, where $L = \sup_{n\geq 1}\{k_n\}$ and asymptotically pseudocontractive mapping.*

Definition 1.3. (1) Let $T : D(T) \subset E \to E$ be a mapping, let $D(T)$ be a nonempty convex subset of E, let $x_0 \in D(T)$ be a given point, and let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, and $\{\delta_n\}$ be four sequences in $[0, 1]$. Then the sequence $\{x_n\}$ defined by

$$\begin{aligned} x_{n+1} &= (1 - \alpha_n - \gamma_n)x_n + \alpha_n T^n y_n + \gamma_n u_n, \\ y_n &= (1 - \beta_n - \delta_n)x_n + \beta_n T^n x_n + \delta_n v_n, \end{aligned} \quad \forall n \geq 0, \tag{1.5}$$

is called the modified Ishikawa iterative sequence with errors of T, where $\{u_n\}$ and $\{v_n\}$ are two bounded sequences in $D(T)$.

(2) In (1.5) if $\beta_n = 0$ and $\delta_n = 0$, $n = 0, 1, 2, \ldots$, then $y_n = x_n$. The sequence $\{x_n\}$ defined by

$$x_{n+1} = (1 - \alpha_n - \gamma_n)x_n + \alpha_n T^n x_n + \gamma_n u_n, \quad \forall n \geq 0, \tag{1.6}$$

is called the modified Mann iterative sequence with errors of T.

In this paper, we discuss convergence theorems of modified Ishikawa and Mann iterative sequences with errors for asymptotically pseudocontractive and asymptotically nonexpansive mappings in Banach spaces, and the boundedness of the domain and range can be dropped, generalizing theorems of Chang [1].

Lemma 1.4 [4]. *Let $\{A_n\}$, $\{B_n\}$, and $\{C_n\}$ be sequences of nonnegative real numbers satisfying the inequality*

$$A_{n+1} \leq (1 + B_n)A_n + C_n, \quad \forall n \geq 0. \tag{1.7}$$

If $\sum_{n=0}^{\infty} B_n < +\infty$ and $\sum_{n=0}^{\infty} C_n < +\infty$, then $\lim_{n\to\infty} A_n$ exists.

2. Main results

Lemma 2.1. *Let E be an arbitrary real Banach space, let D be a nonempty closed convex subset of E, let $T : D \to D$ be a uniformly L-Lipschitzian asymptotically pseudocontractive mapping with a sequence $\{k_n\} \subset [1, \infty)$, $\sum_{n=0}^{\infty}(k_n - 1) < +\infty$. Let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, and*

$\{\delta_n\}$ *be four sequences in* $[0,1]$ *satisfying the following conditions:*

(i) $\alpha_n + \gamma_n \leq 1$, $\beta_n + \delta_n \leq 1$;
(ii) $\beta_n \leq \alpha_n$, $\delta_n \leq \gamma_n$, *for all* $n \geq 0$;
(iii) $\sum_{n=0}^{\infty} \gamma_n < +\infty$;
(iv) $\sum_{n=0}^{\infty} \alpha_n^2 < +\infty$.

Let $x_0 \in D$ be any given point and let $\{x_n\}$ and $\{y_n\}$ be the modified Ishikawa iterative sequence with errors defined by (1.5). If $F(T) \neq \emptyset$, then for any given $q \in F(T)$, $\lim_{n\to\infty} \|x_n - q\|$ exists.

Proof. Set $M = \max\{\sup_{n\geq 0} \|u_n - q\|, \sup_{n\geq 0} \|v_n - q\|\}$.

Since T is asymptotically pseudocontractive, for all $x, y \in D$, there exists $j(x-y) \in J(x-y)$ such that

$$\langle T^n x - T^n y,\ j(x-y)\rangle \leq k_n \|x-y\|^2. \tag{2.1}$$

Then from inequality (2.1), we obtain

$$\langle (k_n I - T^n)x - (k_n I - T^n)y,\ j(x-y)\rangle = k_n\|x-y\|^2 - \langle T^n x - T^n y,\ j(x-y)\rangle \geq 0, \tag{2.2}$$

and it follows from Kato [2] that

$$\|x-y\| \leq \|x - y + \lambda[(k_n I - T^n)x - (k_n I - T^n)y]\|, \quad \forall x, y \in D,\ \lambda > 0. \tag{2.3}$$

Set $a_n := \alpha_n + \gamma_n$. Then from the recursive formula (1.5), we have $x_{n+1} = (1-a_n)x_n + a_n T^n y_n - \gamma_n(T^n y_n - u_n)$. It follows that

$$\begin{aligned} x_n = {} & (1+a_n)x_{n+1} + a_n(k_n I - T^n)x_{n+1} - a_n k_n x_n + a_n^2(1+k_n)(x_n - T^n y_n) \\ & + a_n(T^n x_{n+1} - T^n y_n) + \gamma_n[1 + a_n(1+k_n)](T^n y_n - u_n). \end{aligned} \tag{2.4}$$

Observe that

$$q = (1+a_n)q + a_n(k_n I - T^n)q - a_n k_n q. \tag{2.5}$$

So that

$$\begin{aligned} x_n - q = {} & (1+a_n)(x_{n+1} - q) + a_n[(k_n I - T^n)x_{n+1} - (k_n I - T^n)q] - a_n k_n (x_n - q) \\ & + a_n^2(1+k_n)(x_n - T^n y_n) + a_n(T^n x_{n+1} - T^n y_n) + \gamma_n[1 + a_n(1+k_n)](T^n y_n - u_n). \end{aligned} \tag{2.6}$$

Hence

$$\begin{aligned}||x_n - q|| &\geq (1+a_n)\left|\left|x_{n+1} - q + \frac{a_n}{1+a_n}[(k_nI - T^n)x_{n+1} - (k_nI - T^n)q]\right|\right| - a_nk_n||x_n - q|| \\ &\quad - a_n^2(1+k_n)||x_n - T^ny_n|| - a_n||T^nx_{n+1} - T^ny_n|| \\ &\quad - \gamma_n[1 + a_n(1+k_n)]||T^ny_n - u_n|| \\ &\geq (1+a_n)||x_{n+1} - q|| - a_nk_n||x_n - q|| - a_n^2(1+k_n)||x_n - T^ny_n|| \\ &\quad - a_n||T^nx_{n+1} - T^ny_n|| - \gamma_n[1 + a_n(1+k_n)]||T^ny_n - u_n||.\end{aligned} \tag{2.7}$$

So

$$\begin{aligned}||x_{n+1} - q|| &\leq \left[1 + \frac{a_n}{1+a_n}(k_n - 1)\right]||x_n - q|| + a_n^2(1+k_n)||x_n - T^ny_n|| \\ &\quad + a_n||T^nx_{n+1} - T^ny_n|| + \gamma_n[1 + a_n(1+k_n)]||T^ny_n - u_n||.\end{aligned} \tag{2.8}$$

Furthermore, set $b_n := \beta_n + \delta_n$. Then from recursive formula (1.5), we have $y_n = (1 - b_n)x_n + b_nT^nx_n - \delta_n(T^nx_n - v_n)$. By condition (ii), we have $b_n \leq a_n$, for all $n \geq 0$. We make the following estimates:

$$\begin{aligned}||y_n - q|| &= ||(1-b_n)(x_n - q) + b_n(T^nx_n - q) - \delta_n(T^nx_n - v_n)|| \\ &\leq [1 + b_n(L-1)]||x_n - q|| + \delta_n||T^nx_n - v_n|| \\ &\leq L||x_n - q|| + \delta_nL||x_n - q|| + \delta_nM \\ &= L(1+\delta_n)||x_n - q|| + \delta_nM, \\ ||x_n - T^ny_n|| &\leq ||x_n - q|| + L||y_n - q|| \\ &\leq ||x_n - q|| + L^2(1+\delta_n)||x_n - q|| + L\delta_nM \\ &\leq [1 + L^2(1+\delta_n)]||x_n - q|| + L\delta_nM, \\ ||T^ny_n - u_n|| &\leq L||y_n - q|| + ||u_n - q|| \\ &\leq L^2(1+\delta_n)||x_n - q|| + (1 + L\delta_n)M, \\ ||T^nx_{n+1} - T^ny_n|| &\leq L||x_{n+1} - y_n|| \\ &= L||x_n - y_n + a_n(T^ny_n - x_n) - \gamma_n(T^ny_n - u_n)|| \\ &\leq L||x_n - y_n|| + La_n||T^ny_n - x_n|| + L\gamma_n||T^ny_n - u_n|| \\ &= L||b_n(x_n - T^nx_n) + \delta_n(T^nx_n - v_n)|| \\ &\quad + La_n||T^ny_n - x_n|| + L\gamma_n||T^ny_n - u_n||\end{aligned} \tag{2.9}$$

$$\begin{aligned}
&\le Lb_n||x_n - T^n x_n|| + L\delta_n||T^n x_n - v_n|| \\
&\quad + La_n||T^n y_n - x_n|| + L\gamma_n||T^n y_n - u_n|| \\
&\le La_n(1+L)||x_n - q|| + L^2\delta_n||x_n - q|| \\
&\quad + L\delta_n M + La_n[1 + L^2(1+\delta_n)]||x_n - q|| \\
&\quad + L^2 a_n\delta_n M + L^3\gamma_n(1+\delta_n)||x_n - q|| + L\gamma_n(1+L\delta_n)M.
\end{aligned} \tag{2.10}$$

Using (2.9) and (2.10) in (2.8), we obtain the following estimation:

$$\begin{aligned}
||x_{n+1} - q|| &\le \left[1 + \frac{a_n}{1+a_n}(k_n - 1)\right]||x_n - q|| + a_n^2(1+k_n)[1 + L^2(1+\delta_n)]||x_n - q|| \\
&\quad + a_n^2(1+k_n)L\delta_n M + La_n^2(1+L)||x_n - q|| + L^2 a_n\delta_n||x_n - q|| + La_n\delta_n M \\
&\quad + La_n^2[1 + L^2(1+\delta_n)]||x_n - q|| + L^2 a_n^2\delta_n M + L^3 a_n\gamma_n(1+\delta_n)||x_n - q|| \\
&\quad + La_n\gamma_n(1+L\delta_n)M + \gamma_n[1 + a_n(1+k_n)]L^2(1+\delta_n)||x_n - q|| \\
&\quad + \gamma_n[1 + a_n(1+k_n)](1+L\delta_n)M \\
&= \left\{1 + \frac{a_n}{1+a_n}(k_n - 1) + a_n^2(1+k_n)[1 + L^2(1+\delta_n)] + La_n^2(1+L) + L^2 a_n\delta_n \right. \\
&\quad + La_n^2[1 + L^2(1+\delta_n)] + L^3 a_n\gamma_n(1+\delta_n) \\
&\quad \left. + \gamma_n[1 + a_n(1+k_n)]L^2(1+\delta_n)\right\}||x_n - q|| \\
&\quad + \left\{a_n^2(1+k_n)L\delta_n M + La_n\delta_n M + L^2 a_n^2\delta_n M \right. \\
&\quad \left. + La_n\gamma_n(1+L\delta_n)M + \gamma_n[1 + a_n(1+k_n)](1+L\delta_n)M\right\}.
\end{aligned} \tag{2.11}$$

Set

$$\begin{aligned}
A_n &:= ||x_n - q||, \\
B_n &:= \frac{a_n}{1+a_n}(k_n - 1) + a_n^2(1+k_n)[1 + L^2(1+\delta_n)] + La_n^2(1+L) + L^2 a_n\delta_n \\
&\quad + La_n^2[1 + L^2(1+\delta_n)] + L^3 a_n\gamma_n(1+\delta_n) + \gamma_n[1 + a_n(1+k_n)]L^2(1+\delta_n), \\
C_n &:= a_n^2(1+k_n)L\delta_n M + La_n\delta_n M + L^2 a_n^2\delta_n M + La_n\gamma_n(1+L\delta_n)M \\
&\quad + \gamma_n[1 + a_n(1+k_n)](1+L\delta_n)M,
\end{aligned} \tag{2.12}$$

then inequality (2.11) is equal to

$$A_{n+1} \leq (1 + B_n)A_n + C_n. \tag{2.13}$$

By conditions (ii), (iii), and (iv), we know $\sum_{n=0}^{\infty} B_n < +\infty$, $\sum_{n=0}^{\infty} C_n < +\infty$. By Lemma 1.4, we know $\lim_{n\to\infty} \|x_n - q\|$ exists. □

Theorem 2.2. *Let E be a real uniformly smooth Banach space, let D be a nonempty closed convex subset of E, and let $T : D \to D$ be a uniformly L-Lipschitzian asymptotically pseudocontractive mapping with a sequence $\{k_n\} \subset [1, \infty)$, $\sum_{n=0}^{\infty}(k_n - 1) < +\infty$. Let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, and $\{\delta_n\}$ be four sequences in $[0, 1]$ satisfying the conditions (i)–(iv) in Lemma 2.1.*

Let $x_0 \in D$ be any given point and let $\{x_n\}$, $\{y_n\}$ be the modified Ishikawa iterative sequence with errors defined by (1.5).

(1) If $\{x_n\}$ converges strongly to a fixed point q of T in D, then there exists a nondecreasing function $\phi : [0, \infty) \to [0, \infty)$, $\phi(0) = 0$ such that

$$\langle T^n y_n - q, J(y_n - q)\rangle \leq k_n \|y_n - q\|^2 - \phi(\|y_n - q\|), \tag{*}$$

for all $n \geq 0$.

(2) Conversely, if there exists a strictly increasing function $\phi : [0, \infty) \to [0, \infty)$, $\phi(0) = 0$ satisfying condition (*), then $x_n \to q \in F(T)$.

Proof. Since E is uniformly smooth, the normalized duality mapping $J : E \to E^*$ is single-valued and uniformly continous on any bounded subset of E.

(1) Let $x_n \to q \in F(T)$. From conditions (ii)–(iv) in Lemma 2.1, we have $\beta_n \to 0$, $\delta_n \to 0$. Besides noticing $\|y_n - q\| \leq L(1 + \delta_n)\|x_n - q\| + \delta_n M$ in (2.9) of Lemma 2.1, we have

$$y_n \longrightarrow q \quad (n \longrightarrow \infty). \tag{2.14}$$

The rest of the proof is the same as Chang's [1, Theorem 2.1].

(2) By Lemma 2.1, we know $\lim_{n\to\infty} \|x_n - q\|$ exists. So $\{x_n\}$ is bounded. And by the proof of Lemma 2.1, we can also get $\{T^n y_n - y_n\}$, $\{x_n - T^n x_n\}$, $\{x_n - v_n\}$, and $\{u_n - y_n\}$; all are bounded. So the rest of the proof is the same as Chang's [1, Theorem 2.1]. □

Remark 2.3. (1) Theorem 2.2 removes the restriction on D which is bounded in Chang [1, Theorem 2.1].

(2) Respectively, we can get Chang [1, Theorems 2.2, 2.3, and 2.4], but without the restriction on D which is bounded.

(3) In Osilike and Akuchu [3], they discussed the common fixed points of a family of asymptotically pseudocontractive maps. Our paper is different from it.

Acknowledgments

This project is supported by National Natural Science Foundations of China (10561007) and the Natural Science Foundations of JiangXi Province (0411036).

References

[1] S. S. Chang, *Some results for asymptotically pseudo-contractive mappings and asymptotically nonexpansive mappings*, Proceedings of the American Mathematical Society **129** (2001), no. 3, 845–853.

[2] T. Kato, *Nonlinear semigroups and evolution equations*, Journal of the Mathematical Society of Japan **19** (1967), 508–520.

[3] M. O. Osilike and B. G. Akuchu, *Common fixed points of a finite family of asymptotically pseudocontractive maps*, Fixed Point Theory and Applications **2004** (2004), no. 2, 81–88.

[4] K.-K. Tan and H. K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, Journal of Mathematical Analysis and Applications **178** (1993), no. 2, 301–308.

Yuchao Tang: Department of Mathematics, NanChang University, NanChang, JiangXi 330047, China
E-mail address: hhaaoo1331@yahoo.com.cn

Liwei Liu: Department of Mathematics, NanChang University, NanChang, JiangXi 330047, China
E-mail address: liulws@yahoo.com.cn

Journal of Applied Mathematics and Decision Sciences

Special Issue on
Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods**: artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning
- **Application fields**: asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects**: decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site http://www.hindawi.com/journals/jamds/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk