NIELSEN NUMBER OF A COVERING MAP

JERZY JEZIERSKI

Received 23 November 2004; Revised 13 May 2005; Accepted 24 July 2005

We consider a finite regular covering $p_H : \widetilde{X}_H \to X$ over a compact polyhedron and a map $f : X \to X$ admitting a lift $\widetilde{f} : \widetilde{X}_H \to \widetilde{X}_H$. We show some formulae expressing the Nielsen number $N(f)$ as a linear combination of the Nielsen numbers of its lifts.

Copyright © 2006 Jerzy Jezierski. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let X be a finite polyhedron and let H be a normal subgroup of $\pi_1(X)$. We fix a covering $p_H : \widetilde{X}_H \to X$ corresponding to the subgroup H, that is, $p_\#(\pi_1(\widetilde{X}_H)) = H$.

We assume moreover that the subgroup H has finite rank, that is, the covering p_H is finite. Let $f : X \to X$ be a map satisfying $f(H) \subset H$. Then f admits a lift

$$\begin{array}{ccc} \widetilde{X}_H & \xrightarrow{\widetilde{f}} & \widetilde{X}_H \\ {\scriptstyle p_H}\downarrow & & \downarrow{\scriptstyle p_H} \\ X & \xrightarrow{f} & X \end{array} \tag{1.1}$$

Is it possible to find a formula expressing the Nielsen number $N(f)$ by the numbers $N(\widetilde{f})$ where $\widetilde{f}$ runs the set of all lifts? Such a formula seems very desirable since the difficulty of computing the Nielsen number often depends on the size of the fundamental group. Since $\pi_1\widetilde{X} \subset \pi_1 X$, the computation of $N(\widetilde{f})$ may be simpler. We will translate this problem to algebra. The main result of the paper is Theorem 4.2 expressing $N(f)$ as a linear combination of $\{N(\widetilde{f}_i)\}$, where the lifts are representing all the H-Reidemeister classes of f.

The discussed problem is analogous to the question about "the Nielsen number product formula" raised by Brown in 1967 [1]. A locally trivial fibre bundle $p : E \to B$ and a

Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2006, Article ID 37807, Pages 1–11
DOI 10.1155/FPTA/2006/37807

fibre map $f : E \to E$ were given and the question was how to express $N(f)$ by $N(\overline{f})$ and $N(f_b)$, where $\overline{f} : B \to B$ denoted the induced map of the base space and f_b was the restriction to the fibre over a fixed point $b \in \mathrm{Fix}(\overline{f})$. This problem was intensively investigated in 70ties and finally solved in 1980 by You [4]. At first sufficient conditions for the "product formula" were formulated: $N(f) = N(\overline{f})N(f_b)$ assuming that $N(f_b)$ is the same for all fixed points $b \in \mathrm{Fix}(\overline{f})$. Later it turned out that in general it is better to expect the formula

$$N(f) = N(f_{b_1}) + \cdots + N(f_{b_s}), \tag{1.2}$$

where $b_1, \ldots, b_s$ represent all the Nielsen classes of $\overline{f}$. One may find an analogy between the last formula and the formulae of the present paper. There are also other analogies: in both cases the obstructions to the above equalities lie in the subgroups $\{\alpha \in \pi_1 X;\ f_\# \alpha = \alpha\} \subset \pi_1 X$.

2. Preliminaries

We recall the basic definitions [2, 3]. Let $f : X \to X$ be a self-map of a compact polyhedron. Let $\mathrm{Fix}(f) = \{x \in X;\ f(x) = x\}$ denote the *fixed point set* of f. We define the *Nielsen relation* on $\mathrm{Fix}(f)$ putting $x \sim y$ if there is a path $\omega : [0,1] \to X$ such that $\omega(0) = x$, $\omega(1) = y$ and the paths ω, $f\omega$ are fixed end point homotopic. This relation splits the set $\mathrm{Fix}(f)$ into the finite number of classes $\mathrm{Fix}(f) = A_1 \cup \cdots \cup A_s$. A class $A \subset \mathrm{Fix}(f)$ is called *essential* if its fixed point index $\mathrm{ind}(f; A) \neq 0$. The number of essential classes is called the *Nielsen number* and is denoted by $N(f)$. This number has two important properties. It is a homotopy invariant and is the lower bound of the number of fixed points: $N(f) \leq \#\mathrm{Fix}(g)$ for every map g homotopic to f.

Similarly we define the *Nielsen relation modulo a normal subgroup* $H \subset \pi_1 X$. We assume that the map f preserves the subgroup H, that is, $f_\# H \subset H$. We say that then $x \sim_H y$ if $\omega = f\omega \bmod H$ for a path ω joining the fixed points x and y. This yields *H-Nielsen* classes and *H-Nielsen number* $N_H(f)$. For the details see [4].

Let us notice that each Nielsen class $\bmod H$ splits into the finite sum of ordinary Nielsen classes (i.e., classes modulo the trivial subgroup): $A = A_1 \cup \cdots \cup A_s$. On the other hand $N_H(f) \leq N(f)$.

We consider a regular finite covering $p : \widetilde{X}_H \to X$ as described above.

Let

$$\mathbb{O}_{XH} = \{\gamma : \widetilde{X}_H \longrightarrow \widetilde{X}_H;\ p_H \gamma = p_H\} \tag{2.1}$$

denote the group of natural transformations of this covering and let

$$\mathrm{lift}_H(f) = \left\{\widetilde{f} : \widetilde{X}_H \longrightarrow \widetilde{X}_H;\ p_H \widetilde{f} = f p_H\right\} \tag{2.2}$$

denote the set of all lifts.

We start by recalling classical results giving the correspondence between the coverings and the fundamental groups of a space.

Lemma 2.1. *There is a bijection $\mathbb{O}_{XH} = p_H^{-1}(x_0) = \pi_1(X)/H$ which can be described as follows:*

$$\gamma \sim \gamma(\widetilde{x}_0) \sim p_H(\widetilde{\gamma}). \tag{2.3}$$

We fix a point $\widetilde{x}_0 \in p_H^{-1}(x_0)$. For a natural transformation $\gamma \in \mathbb{O}_{XH}$, $\gamma(\widetilde{x}_0) \in p_H^{-1}(x_0)$ is a point and $\widetilde{\gamma}$ is a path in $\widetilde{X}_H$ joining the points $\widetilde{x}_0$ and $\gamma(\widetilde{x}_0)$. The bijection is not canonical. It depends on the choice of x_0 and $\widetilde{x}_0$.

Let us notice that for any two lifts $\widetilde{f}, \widetilde{f}' \in \mathrm{lift}_H(f)$ there exists a unique $\gamma \in \mathbb{O}_{XH}$ satisfying $\widetilde{f}' = \gamma\widetilde{f}$. More precisely, for a fixed lift $\widetilde{f}$, the correspondence

$$\mathbb{O}_{XH} \ni \alpha \longrightarrow \alpha\widetilde{f} \in \mathrm{lift}_H(f) \tag{2.4}$$

is a bijection. This correspondence is not canonical. It depends on the choice of $\widetilde{f}$.

The group $\mathbb{O}_{XH}$ is acting on $\mathrm{lift}_H(f)$ by the formula

$$\alpha \circ \widetilde{f} = \alpha \cdot \widetilde{f} \cdot \alpha^{-1} \tag{2.5}$$

and the orbits of this action are called *Reidemeister classes* mod H and their set is denoted $\mathcal{R}_H(f)$. Then one can easily check [3]

(1) $p_H(\mathrm{Fix}(\widetilde{f})) \subset \mathrm{Fix}(f)$ is either exactly one H-Nielsen class of the map f or is empty (for any $\widetilde{f} \in \mathrm{lift}_H(f)$)

(2) $\mathrm{Fix}(f) = \bigcup_{\widetilde{f}} p_H(\mathrm{Fix}(\widetilde{f}))$ where the summation runs the set $\mathrm{lift}_H(f)$

(3) if $p_H(\mathrm{Fix}(\widetilde{f})) \cap p_H(\mathrm{Fix}(\widetilde{f}')) \neq \emptyset$ then $\widetilde{f}$, $\widetilde{f}'$ represent the same Reidemeister class in $\mathcal{R}_H(f)$

(4) if $\widetilde{f}$, $\widetilde{f}'$ represent the same Reidemeister class then $p_H(\mathrm{Fix}(\widetilde{f})) = p_H(\mathrm{Fix}(\widetilde{f}'))$.

Thus $\mathrm{Fix}(f) = \bigcup_{\widetilde{f}} p_H(\mathrm{Fix}(\widetilde{f}))$ is the disjoint sum where the summation is over a subset containing exactly one lift $\widetilde{f}$ from each H-Reidemeister class. This gives the natural inclusion from the set of Nielsen classes modulo H into the set of H-Reidemeister classes

$$\mathcal{N}_H(f) \longrightarrow \mathcal{R}_H(f). \tag{2.6}$$

The H-Nielsen class A is sent into the H-Reidemeister class represented by a lift $\widetilde{f}$ satisfying $p_H(\mathrm{Fix}(\widetilde{f})) = A$. By (1) and (2) such lift exists, by (3) the definition is correct and (4) implies that this map is injective.

3. Lemmas

For a lift $\widetilde{f} \in \mathrm{lift}_H(f)$, a fixed point $x_0 \in \mathrm{Fix}(f)$ and an element $\beta \in \pi_1(X;x_0)$ we define the subgroups

$$\begin{aligned} \mathcal{Z}(\widetilde{f}) &= \left\{\gamma \in \mathbb{O}_{XH};\ \widetilde{f}\gamma = \gamma\widetilde{f}\right\} \\ C(f_\#,x_0;\beta) &= \{\alpha \in \pi_1(X;x_0);\ \alpha\beta = \beta f_\#(\alpha)\} \\ C_H(f_\#,x_0;\beta) &= \{[\alpha]_H \in \pi_1(X;x_0)/H(x_0);\ \alpha\beta = \beta f_\#(\alpha) \text{ modulo } H\}. \end{aligned} \tag{3.1}$$

If $\beta = 1$ we will write simply $C(f_\#,x_0)$ or $C_H(f_\#,x_0)$.

We notice that the canonical projection $j : \pi_1(X;x_0) \to \pi_1(X;x_0)/H(x_0)$ induces the homomorphism $j : C(f_\#,x_0;\beta) \to C_H(f_\#,x_0;\beta)$.

Lemma 3.1. *Let $\widetilde{f}$ be a lift of f and let $\widetilde{A}$ be a Nielsen class of $\widetilde{f}$. Then $p_H(\widetilde{A}) \subset \mathrm{Fix}(f)$ is a Nielsen class of f. On the other hand if $A \subset \mathrm{Fix}(f)$ is a Nielsen class of f then $p_H^{-1}(A) \cap \mathrm{Fix}(\widetilde{f})$ splits into the finite sum of Nielsen classes of $\widetilde{f}$.*

Proof. It is evident that $p_H(\widetilde{A})$ is contained in a Nielsen class $A \subset \mathrm{Fix}(f)$. Now we show that $A \subset p_H(\widetilde{A})$. Let us fix a point $\widetilde{x}_0 \in \widetilde{A}$ and let $x_0 = p_H(\widetilde{x}_0)$. Let $x_1 \in A$. We have to show that $x_1 \in p_H(\widetilde{A})$. Let $\omega : I \to X$ establish the Nielsen relation between the points $\omega(0) = x_0$ and $\omega(1) = x_1$ and let $h(t,s)$ denote the homotopy between $\omega = h(\cdot,0)$ and $f\omega = h(\cdot,1)$. Then the path ω lifts to a path $\widetilde{\omega} : I \to \widetilde{X}_H$, $\widetilde{\omega}(0) = \widetilde{x}_0$. Let us denote $\widetilde{\omega}(1) = \widetilde{x}_1$. It remains to show that $\widetilde{x}_1 \in \widetilde{A}$. The homotopy h lifts to $\widetilde{h} : I \times I \to \widetilde{X}_H$, $\widetilde{h}(0,s) = \widetilde{x}_0$. Then the paths $\widetilde{h}(\cdot,1)$ and $\widetilde{f}\widetilde{\omega}$ as the lifts of $f\omega$ starting from $\widetilde{x}_0$ are equal. Now $\widetilde{f}(\widetilde{x}_1) = \widetilde{f}(\widetilde{\omega}(1)) = \widetilde{h}(1,1) = \widetilde{h}(1,0) = \widetilde{\omega}(1) = \widetilde{x}_1$. Thus $\widetilde{x}_1 \in \mathrm{Fix}(\widetilde{f})$ and the homotopy $\widetilde{h}$ gives the Nielsen relation between $\widetilde{x}_0$ and $\widetilde{x}_1$ hence $\widetilde{x}_1 \in \widetilde{A}$.

Now the second part of the lemma is obvious. □

Lemma 3.2. *Let $\widetilde{A} \subset \mathrm{Fix}(\widetilde{f})$ be a Nielsen class of $\widetilde{f}$. Let us denote $A = p_H(\widetilde{A})$. Then*

1. *$p_H : \widetilde{A} \to A$ is a covering where the fibre is in bijection with the image $j_\#(C(f_\#,x)) \subset \pi_1(X;x)/H(x)$ for $x \in A$,*
2. *the cardinality of the fibre (i.e., $\#(p_H^{-1}(x) \cap \widetilde{A})$) does not depend on $x \in A$ and we will denote it by J_A,*
3. *if $\widetilde{A}'$ is another Nielsen class of $\widetilde{f}$ satisfying $p_H(\widetilde{A}') = p_H(\widetilde{A})$ then the cardinalities of $p_H^{-1}(x) \cap \widetilde{A}$ and $p_H^{-1}(x) \cap \widetilde{A}'$ are the same for each point $x \in A$.*

Proof. (1) Since p_H is a local homeomorphism, the projection $p_H : \widetilde{A} \to A$ is the covering.

(2) We will show a bijection $\phi : j(C(f_\#;x_0)) \to p_H^{-1}(x_0) \cap \widetilde{A}$ (for a fixed point $x_0 \in A$).

Let $\alpha \in C(f_\#)$. Let us fix a point $\widetilde{x}_0 \in p_H^{-1}(x_0)$. Let $\widetilde{\alpha} : I \to \widetilde{X}$ denote the lift of α starting from $\widetilde{\alpha}(0) = \widetilde{x}_0$. We define $\phi([\alpha]_H) = \widetilde{\alpha}(1)$. We show that

(2a) The definition is correct. Let $[\alpha]_H = [\alpha']_H$. Then $\alpha \equiv \alpha' \bmod H$ hence $\widetilde{\alpha}(1) = \widetilde{\alpha'}(1)$. Now we show that $\widetilde{\alpha}(1) \in \widetilde{A}$. Since $\alpha \in C(f_\#)$, there exists a homotopy h between the loops $h(\cdot,0) = \alpha$ and $h(\cdot,1) = f\alpha$. The homotopy lifts to $\widetilde{h} : I \times I \to \widetilde{X}_H$, $\widetilde{h}(0,s) = \widetilde{x}_0$. Then $\widetilde{x}_1 = \widetilde{h}(1,s)$ is also a fixed point of $\widetilde{f}$ and moreover $\widetilde{h}$ is the homotopy between the paths $\widetilde{\omega}$ and $\widetilde{f}\widetilde{\omega}$. Thus $\widetilde{x}_0, \widetilde{x}_1 \in \mathrm{Fix}(\widetilde{f})$ are Nielsen related hence $\widetilde{x}_1 \in \widetilde{A}$.

(2b) ϕ is onto. Let $\widetilde{x}_1 \in p_H^{-1}(x_0) \cap \widetilde{A}$. Now $\widetilde{x}_0, \widetilde{x}_1 \in \mathrm{Fix}(\widetilde{f})$ are Nielsen related. Let $\widetilde{\omega} : I \to \widetilde{X}_H$ establish this relation ($\widetilde{f}\widetilde{\omega} \sim \widetilde{\omega}$). Now

$$f(p_H\widetilde{\omega}) = p_H\widetilde{f}\widetilde{\omega} \sim p_H\widetilde{\omega} \tag{3.2}$$

hence $p_H\widetilde{\omega} \in C(f_\#; x_0)$. Moreover $\phi[p_H\widetilde{\omega}]_H = \widetilde{\omega}(1) = \widetilde{x}_1$.

(2c) ϕ is injective. Let $[\alpha]_H, [\alpha']_H \in j(C(f_\#))$ and let $\widetilde{\alpha}, \widetilde{\alpha}' : I \to \widetilde{X}_H$ be their lifts starting from $\widetilde{\alpha}(0) = \widetilde{\alpha}'(0) = \widetilde{x}_0$. Suppose that $\phi[\alpha]_H = \phi[\alpha']_H$. This means $\widetilde{\alpha}(1) = \widetilde{\alpha}'(1) \in \widetilde{X}_H$. Thus $p_H(\widetilde{\alpha} * \widetilde{\alpha}'^{-1}) = \alpha * \alpha'^{-1} \in H$ which implies $[\alpha]_H = [\alpha']_H$.

(3) Let $x_0 \in p_H(\widetilde{A}) = p_H(\widetilde{A}')$. Then by the above $\#(p^{-1}(x_0) \cap \widetilde{A}) = j_\#(C(f_\#)) = \#(p^{-1}(x_0) \cap \widetilde{A}')$. □

Lemma 3.3. *The restriction of the covering map $p_H : \mathrm{Fix}(\widetilde{f}) \to p_H(\mathrm{Fix}(\widetilde{f}))$ is a covering. The fibre over each point is in a bijection with the set*

$$\mathscr{Z}(\widetilde{f}) = \left\{\gamma \in \mathbb{O}_{XH};\ \widetilde{f}\gamma = \gamma\widetilde{f}\right\}. \tag{3.3}$$

Proof. Since the fibre of the covering p_H is discrete, the restriction $p_H : \mathrm{Fix}(\widetilde{f}) \to p_H(\mathrm{Fix}(\widetilde{f}))$ is a locally trivial bundle. Let us fix points $x_0 \in p_H(\mathrm{Fix}(\widetilde{f}))$, $\widetilde{x}_0 \in p_H^{-1}(x_0) \cap \mathrm{Fix}(\widetilde{f})$. We recall that

$$\alpha : p_H^{-1}(x_0) \longrightarrow \mathbb{O}_{XH}, \tag{3.4}$$

where $\alpha_{\widetilde{x}} \in \mathbb{O}_{XH}$ is characterized by $\alpha_{\widetilde{x}}(\widetilde{x}_0) = \widetilde{x}$, is a bijection. We will show that $\alpha(p_H^{-1}(x_0) \cap \mathrm{Fix}(\widetilde{f})) = \mathscr{Z}(\widetilde{f})$.

Let $\widetilde{f}(\widetilde{x}) = \widetilde{x}$ for an $\widetilde{x} \in p_H^{-1}(x_0)$. Then

$$\widetilde{f}\alpha_{\widetilde{x}}(\widetilde{x}_0) = \widetilde{f}(\widetilde{x}) = \widetilde{x} = \alpha_{\widetilde{x}}(\widetilde{x}_0) = \alpha_{\widetilde{x}}\widetilde{f}(\widetilde{x}_0) \tag{3.5}$$

which implies $\widetilde{f}\alpha_{\widetilde{x}} = \alpha_{\widetilde{x}}\widetilde{f}$ hence $\alpha_{\widetilde{x}} \in \mathscr{Z}(\widetilde{f})$.

Now we assume that $\widetilde{f}\alpha_{\widetilde{x}} = \alpha_{\widetilde{x}}\widetilde{f}$. Then in particular $\widetilde{f}\alpha_{\widetilde{x}}(\widetilde{x}_0) = \alpha_{\widetilde{x}}\widetilde{f}(\widetilde{x}_0)$ which gives $\widetilde{f}(\widetilde{x}) = \alpha_{\widetilde{x}}(\widetilde{x}_0)$, $\widetilde{f}(\widetilde{x}) = \widetilde{x}$ hence $\widetilde{x} \in \mathrm{Fix}(\widetilde{f})$. □

We will denote by I_{A_H} the cardinality of the subgroup $\#\mathscr{Z}(\widetilde{f})$ for the H-Nielsen class $A_H = p_H(\mathrm{Fix}(\widetilde{f}))$. We will also write $I_{A_i} = I_{A_H}$ for any Nielsen class A_i of f contained in A.

Lemma 3.4. *Let $A_0 \subset \mathrm{Fix}(f)$ be a Nielsen class and let $\widetilde{A}_0 \subset \mathrm{Fix}(\widetilde{f})$ be a Nielsen class contained in $p_H^{-1}(A_0)$. Then, by Lemma 3.1 $A_0 = p_H(\widetilde{A}_0)$ and moreover*

$$\begin{aligned} \mathrm{ind}\,(\widetilde{f}; p_H^{-1}(A_0)) &= I_{A_0} \cdot \mathrm{ind}\,(f; A_0) \\ \mathrm{ind}\,(\widetilde{f}; \widetilde{A}_0) &= J_{A_0} \cdot \mathrm{ind}\,(f; A_0). \end{aligned} \tag{3.6}$$

Proof. Since the index is the homotopy invariant we may assume that $\mathrm{Fix}(f)$ is finite. Now for any fixed points $x_0 \in \mathrm{Fix}(f)$, $\widetilde{x}_0 \in \mathrm{Fix}(\widetilde{f})$ satisfying $p_H(\widetilde{x}_0) = x_0$ we have $\mathrm{ind}(\widetilde{f}_0;\widetilde{x}_0) = \mathrm{ind}(f_0;x_0)$ since the projection p_H is a local homeomorphism. Thus

$$\begin{aligned}\mathrm{ind}\left(\widetilde{f};p_H^{-1}(A_0)\right) &= \sum_{x\in A_0}\mathrm{ind}\left(\widetilde{f};p_H^{-1}(x)\right) = \sum_{x\in A_0} I_{A_0}\cdot\mathrm{ind}(f;x)\\ &= I_{A_0}\sum_{x\in A_0}\mathrm{ind}(f;x) = I_{A_0}\cdot\mathrm{ind}\left(f;A_0\right).\end{aligned} \tag{3.7}$$

Similarly we prove the second equality:

$$\begin{aligned}\mathrm{ind}\left(\widetilde{f};\widetilde{A}_0\right) &= \sum_{x\in A_0}\mathrm{ind}\left(\widetilde{f};p_H^{-1}(x)\cap\widetilde{A}_0\right) = \sum_{x\in A_0}\sum_{\widetilde{x}\in p_H^{-1}(x)\cap\widetilde{A}_0}\mathrm{ind}\left(\widetilde{f};\widetilde{x}\right)\\ &= \sum_{x\in A_0} J_{A_0}\cdot\mathrm{ind}(f;x) = J_{A_0}\cdot\left(\sum_{x\in A_0}\mathrm{ind}(f;x)\right) = J_{A_0}\cdot\mathrm{ind}\left(f;A_0\right).\end{aligned} \tag{3.8}$$

□

To get a formula expressing $N(f)$ by the numbers $N(\widetilde{f})$ we will need the assumption that the numbers $J_A = J_{A'}$ for any two H-Nielsen related classes $A, A' \subset \mathrm{Fix}(f)$. The next lemma gives a sufficient condition for such equality.

Lemma 3.5. *Let $x_0 \in p(\mathrm{Fix}(\widetilde{f}))$. If the subgroups $H(x_0), C(f,x_0) \subset \pi_1(X,x_0)$ commute, that is, $h\cdot\alpha = \alpha\cdot h$, for any $h \in H(x_0)$, $\alpha \in C(f,x_0)$, then $J_A = J_{A'}$ for all Nielsen classes $A, A' \subset p(\mathrm{Fix}(\widetilde{f}))$.*

Proof. Let $x_1 \in p(\mathrm{Fix}(\widetilde{f}))$ be another point. The points $x_0, x_1 \in p(\mathrm{Fix}(\widetilde{f}))$ are H-Nielsen related, that is, there is a path $\omega : [0,1] \to X$ satisfying $\omega(0) = x_0$, $\omega(1) = x_1$ such that $\omega * f(\omega^{-1}) \in H(x_0)$. We will show that the conjugation

$$\pi_1(X,x_0) \ni \alpha \longrightarrow \omega^{-1} * \alpha * \omega \in \pi_1(X,x_1) \tag{3.9}$$

sends $C(f,x_0)$ onto $C(f,x_1)$. Let $\alpha \in C(f,x_0)$. We will show that $\omega^{-1} * \alpha * \omega \in C(f,x_1)$. In fact $f(\omega^{-1} * \alpha * \omega) = \omega^{-1} * \alpha * \omega \Leftrightarrow (\omega * f\omega^{-1}) * \alpha = \alpha * (\omega * f\omega^{-1})$ but the last equality holds since $\omega * f\omega^{-1} \in H(x_0)$ and $\alpha \in C(f,x_0)$. □

Remark 3.6. The assumption of the above lemma is satisfied if at least one of the groups $H(x_0)$, $C(f,x_0)$ belongs to the center of $\pi_1(X;x_0)$.

Remark 3.7. Let us notice that if the subgroups $H(x_0), C(f,x_0) \subset \pi_1(X,x_0)$ commute then so do the corresponding subgroups at any other point $x_1 \in p_H(\mathrm{Fix}(\widetilde{f}))$.

Proof. Let us fix a path $\omega : [0,1] \to X$. We will show that the conjugation

$$\pi_1(X,x_0) \ni \alpha \longrightarrow \omega^{-1} * \alpha * \omega \in \pi_1(X,x_1) \tag{3.10}$$

sends $C(f,x_0)$ onto $C(f,x_1)$. Let $\alpha \in C(f,x_0)$. We will show that $\omega^{-1} * \alpha * \omega \in C(f,x_1)$. But the last means $f(\omega^{-1} * \alpha * \omega) = \omega^{-1} * \alpha * \omega$ hence $f(\omega^{-1}) * f\alpha * f\omega = \omega^{-1} * \alpha * \omega \Leftrightarrow f(\omega^{-1}) * \alpha * f\omega = \omega^{-1} * \alpha * \omega \Leftrightarrow (\omega * f\omega^{-1}) * \alpha = \alpha * (\omega * f\omega^{-1})$ and the last

holds since $(\omega * f\omega^{-1}) \in H(x_0)$ and $\alpha \in C(f,x_0)$. Now it remains to notice that the elements of $H(x_1)$, $C(f;x_1)$ are of the form $\omega^{-1} * \gamma * \omega$ and $\omega^{-1} * \alpha * \omega$ respectively for some $\gamma \in H(x_0)$ and $\alpha \in C(f,x_0)$. □

Now we will express the numbers I_A, J_A in terms of the homotopy group homomorphism $f_\# : \pi_1(X,x_0) \to \pi_1(X,x_0)$ for a fixed point $x_0 \in \mathrm{Fix}(f)$. Let $\widetilde{f} : \widetilde{X}_H \to \widetilde{X}_H$ be a lift satisfying $\widetilde{x}_0 \in p_H^{-1}(x_0) \cap \mathrm{Fix}(\widetilde{f})$. We also fix the isomorphism

$$\pi_1(X;x_0)/H(x_0) \ni \alpha \longrightarrow \gamma_\alpha \in \mathbb{O}_{XH}, \tag{3.11}$$

where $\gamma_\alpha(\widetilde{x}_0) = \widetilde{\alpha}(1)$ and $\widetilde{\alpha}$ denotes the lift of α starting from $\widetilde{\alpha}(0) = \widetilde{x}_0$.

We will describe the subgroup corresponding to $C(\widetilde{f})$ by this isomorphism and then we will do the same for the other lifts $\widetilde{f}' \in \mathrm{lift}_H(f)$.

Lemma 3.8.

$$\widetilde{f}\gamma_\alpha = \gamma_{f\alpha}\widetilde{f}. \tag{3.12}$$

Proof.

$$\widetilde{f}\gamma_\alpha(\widetilde{x}_0) = \widetilde{f}\widetilde{\alpha}(1) = \gamma_{f\alpha}(\widetilde{x}_0) = \gamma_{f\alpha}\widetilde{f}(\widetilde{x}_0), \tag{3.13}$$

where the middle equality holds since $\widetilde{f}\widetilde{\alpha}$ is a lift of the path $f\alpha$ from the point $\widetilde{x}_0$. □

Corollary 3.9. *There is a bijection between*

$$\begin{aligned} \mathfrak{L}(\widetilde{f}) &= \left\{\gamma \in \mathbb{O}_{XH};\ \widetilde{f}\gamma = \gamma\widetilde{f}\right\}, \\ C_H(f) &= \{\alpha \in \pi_1(X;x_0)/H(x_0);\ f_{H\#}(\alpha) = \alpha\}. \end{aligned} \tag{3.14}$$

Thus

$$I_A/J_A = \#\mathfrak{L}(\widetilde{f})/\#j(C(f)) = \#(C_H(f)/j(C(f))). \tag{3.15}$$

Let us emphasize that $C(f)$, $C_H(f)$ are the subgroups of $\pi_1(X;x_0)$ or $\pi_1(X;x_0)/H(x_0)$ respectively where the base point is the chosen fixed point. Now will take another fixed point $x_1 \in \mathrm{Fix}(f)$ and we will denote $C'(f) = \{\alpha' \in \pi_1(X;x_1);\ f_\#\alpha = \alpha\}$ and similarly we define $C'_H(f)$. We will express the cardinality of these subgroups in terms of the group $\pi_1(X;x_0)$.

Lemma 3.10. *Let $\eta : [0,1] \to X$ be a path from x_0 to x_1. This path gives rise to the isomorphism $P_\eta : \pi_1(X;x_1) \to \pi_1(X;x_0)$ by the formula $P_\eta(\alpha) = \eta\alpha\eta^{-1}$. Let $\delta = \eta \cdot (f\eta)^{-1}$. Then*

$$\begin{aligned} P_\eta(C'(f)) &= \{\alpha \in \pi_1(X;x_0);\ \alpha\delta = \delta f_\#(\alpha)\} \\ P_\eta(C'_H(f)) &= \{[\alpha] \in \pi_1(X;x_0)/H(x_0);\ \alpha\delta = \delta f_\#(\alpha) \textit{ modulo } H\}. \end{aligned} \tag{3.16}$$

Proof. We notice that δ is a loop based at x_0 representing the Reidemeister class of the point x_1 in $\mathcal{R}(f) = \pi_1(X;x_0)/\mathcal{R}$.

We will denote the right-hand side of the above equalities by $C(f;\delta)$ and $C_H(f;\delta)$ respectively. Let $\alpha' \in \pi_1(X;x_1)$. We denote $\alpha = P_\eta(\alpha') = \eta\alpha'\eta^{-1}$. We will show that $\alpha \in C(f;\delta) \Leftrightarrow \alpha' \in C'(f)$.

In fact $\alpha \in C(f;\delta) \Leftrightarrow \alpha\delta = \delta \cdot f\alpha \Leftrightarrow (\eta\alpha'\eta^{-1})(\eta \cdot f\eta^{-1}) = (\eta \cdot f\eta^{-1})(f\eta \cdot f\alpha' \cdot (f\eta)^{-1}) \Leftrightarrow \eta\alpha' \cdot (f\eta)^{-1} = \eta \cdot f\alpha' \cdot (f\eta)^{-1} \Leftrightarrow \alpha' = f\alpha'$.

Similarly we prove the second equality. □

Thus we get the following formulae for the numbers I_A, J_A.

Corollary 3.11. *Let $\delta \in \pi_1(X;x_0)$ represent the Reidemeister class $A \in \mathcal{R}(f)$. Then $I_A = \#C_H(f;j(\delta))$, $J_A = \#j(C(f;\delta))$.*

4. Main theorem

Lemma 4.1. *Let $A \subset p_H(\mathrm{Fix}(\widetilde{f}))$ be a Nielsen class of f. Then $p_H^{-1}A$ contains exactly I_A/J_A fixed point classes of $\widetilde{f}$.*

Proof. Since the projection of each Nielsen class $\widetilde{A} \subset p_H^{-1}(A) \cap \mathrm{Fix}(\widetilde{f})$ is onto A (Lemma 3.1), it is enough to check how many Nielsen classes of $\widetilde{f}$ cut $p_H^{-1}(a)$ for a fixed point $a \in A$. But by Lemma 3.3 $p_H^{-1}(a) \cap \mathrm{Fix}(\widetilde{f})$ contains I_A points and by Lemma 3.2 each class in this set has exactly J_A common points with $p_H^{-1}(a)$. Thus exactly I_A/J_A Nielsen classes of $\widetilde{f}$ are cutting $p_H^{-1}(a) \cap \mathrm{Fix}(\widetilde{f})$. □

Let $f : X \to X$ be a self-map of a compact polyhedron admitting a lift $\widetilde{f} : \widetilde{X}_H \to \widetilde{X}_H$. We will need the following auxiliary assumption:

> *for any Nielsen classes $A, A' \in \mathrm{Fix}(f)$ representing the same class modulo the subgroup H the numbers $J_A = J_{A'}$.*

We fix lifts $\widetilde{f}_1, \ldots, \widetilde{f}_s$ representing all H-Nielsen classes of f, that is,

$$\mathrm{Fix}(f) = p_H\Big(\mathrm{Fix}\,(\widetilde{f}_1)\Big) \cup \cdots \cup p_H\big(\mathrm{Fix}\,(\widetilde{f}_s)\big) \tag{4.1}$$

is the mutually disjoint sum. Let I_i, J_i denote the numbers corresponding to a (Nielsen class of f) $A \subset p_H(\mathrm{Fix}(\widetilde{f}_i))$. By the remark after Lemma 3.3 and by the above assumption these numbers do not depend on the choice of the class $A \subset p_H(\mathrm{Fix}(\widetilde{f}_i))$. We also notice that Lemmas 3.3, 3.2 imply

$$\begin{aligned} I_i &= \#\mathcal{L}(\widetilde{f}_i) = \#\big\{\gamma \in \mathbb{O}_{XH};\ \gamma\widetilde{f}_i = \widetilde{f}_i\gamma\big\} \\ J_i &= \#j(C(f_\#;x)) = \#j(\{\gamma \in \pi_1(X,x_i);\ f_\#\gamma = \gamma\}) \end{aligned} \tag{4.2}$$

for an $x_i \in A_i$.

Theorem 4.2. *Let X be a compact polyhedron, $P_H : \widetilde{X}_H \to \widetilde{X}$ a finite regular covering and let $f : X \to X$ be a self-map admitting a lift $\widetilde{f} : \widetilde{X}_H \to \widetilde{X}_H$. We assume that for each two Nielsen classes $A, A' \subset \mathrm{Fix}(f)$, which represent the same Nielsen class modulo the subgroup H, the numbers $J_A = J_{A'}$. Then*

$$N(f) = \sum_{i=1}^{s} (J_i/I_i) \cdot N(\widetilde{f}_i), \tag{4.3}$$

where I_i, J_i denote the numbers defined above and the lifts $\widetilde{f}_i$ represent all H-Reidemeister classes of f, corresponding to nonempty H-Nielsen classes.

Proof. Let us denote $A_i = p_H(\mathrm{Fix}(\widetilde{f}_i))$. Then A_i is the disjoint sum of Nielsen classes of f. Let us fix one of them $A \subset A_i$. By Lemma 3.1 $p_H^{-1}A \cap \mathrm{Fix}(\widetilde{f}_i)$ splits into I_A/J_A Nielsen classes in $\mathrm{Fix}(\widetilde{f}_i)$. By Lemma 3.4 A is essential iff one (hence all) Nielsen classes in $p_H^{-1}A \subset \mathrm{Fix}\,\widetilde{f}_i$ is essential. Summing over all essential classes of $\widetilde{f}$ in $A_i = p_A(\mathrm{Fix}(\widetilde{f}_i))$ we get

$$\begin{aligned} &\text{the number of essential Nielsen classes of } f \text{ in } A_i \\ &\quad = \sum_{A} (J_A/I_A) \cdot (\text{number of essential Nielsen classes of } \widetilde{f}_i \text{ in } p_H^{-1}A), \end{aligned} \tag{4.4}$$

where the summation runs the set of all essential Nielsen classes contained in A_i.

But $J_A = J_i$, $I_A = I_i$ for all $A \subset A_i$ hence

$$(\text{the number of essential Nielsen classes of } f \text{ in } A_i) = J_i/I_i \cdot N(\widetilde{f}_i). \tag{4.5}$$

Summing over all lifts $\{\widetilde{f}_i\}$ representing non-empty H-Nielsen classes of f we get

$$N(f) = \sum_{i} (J_i/I_i) \cdot N(\widetilde{f}_i) \tag{4.6}$$

since $N(f)$ equals the number of essential Nielsen classes in $\mathrm{Fix}(f) = \bigcup_{i=1}^{s} p_H \mathrm{Fix}(\widetilde{f}_i)$. □

Corollary 4.3. *If moreover, under the assumptions of Theorem 4.2, $C = J_i/I_i$ does not depend on i then*

$$N(f) = C \cdot \sum_{i=1}^{s} N(\widetilde{f}_i). \tag{4.7}$$

5. Examples

In all examples given below the auxiliary assumption $J_A = J_{A'}$ holds, since the assumptions of Lemma 3.5 are satisfied (in 1, 2, 3 and 5 the fundamental groups are commutative and in 4 the subgroup $C(f, x_0)$ is trivial).

(1) If $\pi_1 X$ is finite and $p : \widetilde{X} \to X$ is the universal covering (i.e., $H = 0$) then $\widetilde{X}$ is simply connected hence for any lift $\widetilde{f} : \widetilde{X} \to \widetilde{X}$

$$N(\widetilde{f}) = \begin{cases} 1 & \text{for } L(\widetilde{f}) \neq 0 \\ 0 & \text{for } L(\widetilde{f}) = 0. \end{cases} \tag{5.1}$$

But $L(\widetilde{f}) \neq 0$ if and only if the Nielsen class $p(\mathrm{Fix}(\widetilde{f})) \subset \mathrm{Fix}(f)$ is essential (Lemma 3.4). Thus

$$N(f) = \text{number of essential classes} = N(\widetilde{f}_1) + \cdots + N(\widetilde{f}_s), \tag{5.2}$$

where the lifts $\widetilde{f}_1, \ldots, \widetilde{f}_s$ represent all Reidemeister classes of f.

(2) Consider the commutative diagram

$$\begin{array}{ccc} S^1 & \xrightarrow{p_l} & S^1 \\ {\scriptstyle p_k}\downarrow & & \downarrow{\scriptstyle p_k} \\ S^1 & \xrightarrow{p_l} & S^1 \end{array} \tag{5.3}$$

Where $p_k(z) = z^k$, $p_l(z) = z^l$, $k,l \geq 2$. The map p_k is regarded as k-fold regular covering map. Then each natural transformation map of this covering is of the form $\alpha(z) = \exp(2\pi p/k) \cdot z$ for $p = 0, \ldots, k-1$ hence is homotopic to the identity map. Now all the lifts of the map p_l are maps of degree l hence their Nielsen numbers equal $l-1$. On the other hand the Reidemeister relation of the map $p_l : S^1 \to S^1$ modulo the subgroup $H = \mathrm{im} p_{k\#}$ is given by

$$\begin{aligned} \alpha \sim \beta &\iff \beta = \alpha + p(l-1) \in k \cdot \mathbb{Z} \quad \text{for a } p \in \mathbb{Z} \\ &\iff \beta = \alpha + p(l-1) + qk \quad \text{for some } p, q \in \mathbb{Z} \\ &\iff \alpha = \beta \text{ modulo g.c.d. } (l-1, k). \end{aligned} \tag{5.4}$$

Thus $\#\mathcal{R}_H(p_l) = \text{g.c.d.}(l-1,k)$. Now the sum

$$\sum_{p_l'} N(p_l') = \big(\text{g.c.d.}(l-1,k)\big) \cdot (l-1), \tag{5.5}$$

(where the summation runs the set having exactly one common element with each H-Reidemeister class) equals $N(p_l) = l-1$ iff the numbers k, $l-1$ are relatively prime.

Notice that in our notation $I = \text{g.c.d.}(l-1,k)$ while $J = 1$.

(3) Let us consider the action of the cyclic group $\mathbb{Z}_8$ on $S^3 = \{(z,z') \in \mathbb{C} \times \mathbb{C};\ |z|^2 + |z'|^2 = 1\}$ given by the cyclic homeomorphism

$$S^3 \ni (z,z') \longrightarrow \big(\exp(2\pi i/8) \cdot z, \exp(2\pi i/8) \cdot z'\big) \in S^3. \tag{5.6}$$

The quotient space is the lens space which we will denote L_8. We will also consider the quotient space of S^3 by the action of the subgroup $2\mathbb{Z}_4 \subset \mathbb{Z}_8$. Now the quotient group is

also a lens space which we will denote by L_4. Let us notice that there is a natural 2-fold covering $p_H : L_4 \to L_8$

$$L_4 = S^3/\mathbb{Z}_4 \ni [z,z'] \longrightarrow [z,z'] \in S^3/\mathbb{Z}_8 = L_8. \tag{5.7}$$

The group of natural transformations $\mathbb{O}_L$ of this covering contains two elements: the identity and the map $A[z,z'] = [\exp(2\pi i/8)\cdot z, \exp(2\pi i/8)\cdot z']$. Now we define the map $f : L_8 \to L_8$ putting $f[z,z'] = [z^7/|z|^6, z'^7/|z|'^6]$. This map admits the lift $\widetilde{f} : L_4 \to L_4$ given by the same formula and the lift $A\widetilde{f}$. We notice that each of the maps f, $\widetilde{f}$, $A\widetilde{f}$ is a map of a closed oriented manifold of degree 49. Since $H_1(L;\mathbb{Q}) = H_2(L;\mathbb{Q}) = 0$ for all lens spaces, the Lefschetz number of each of these three maps equals; $L(f) = 1 - 49 = -48 \neq 0$. On the other hand since the lens spaces are Jiang [3], all involved Reidemeister classes are essential hence the Nielsen number equals the Reidemeister number in each case.

Now

$$\mathcal{R}(f) = \operatorname{coker}(\mathrm{id} - 7\cdot\mathrm{id}) = \operatorname{coker}(-6\cdot\mathrm{id}) = \operatorname{coker}(2\cdot\mathrm{id}) = \mathbb{Z}_2. \tag{5.8}$$

Similarly $\mathcal{R}(\widetilde{f}) = \mathbb{Z}_2$ and $\mathcal{R}(A\cdot\widetilde{f}) = \mathcal{R}(\widetilde{f}) = \mathbb{Z}_2$ since A is homotopic to the identity. Thus

$$R(f) = 2 \neq 2 + 2 = R(\widetilde{f}) + R(A\cdot\widetilde{f}). \tag{5.9}$$

Since all the classes are essential, the same inequality holds for the Nielsen numbers.

(4) If the group $\{\alpha \in \pi_1(X;x)/H(x);\ f_\#\alpha = \alpha\}$ is trivial for each $x \in \mathrm{Fix}(f)$ lying in an essential Nielsen class of f then all the numbers $I_i = J_i = 1$ and the sum formula holds.

(5) If $\pi_1 X/H$ is abelian then the rank of the groups

$$C(f_{H\#}) = \{\alpha \in \pi_1(X,x)/H(x);\ f_\#\alpha = \alpha\} = \ker\left(\mathrm{id} - f_\#\right) : \pi_1(X,x)/H(x) \longrightarrow \pi_1(X,x)/H(x) \tag{5.10}$$

does not depend on $x \in \mathrm{Fix}(f)$ hence I is constant. If moreover $\pi_1 X$ is abelian then also the group $C(f_\#) = \ker(\mathrm{id} - f_\#)$ does not depend on $x \in \mathrm{Fix}(f)$. Then we get

$$N(f) = J/I \cdot \left(N(\widetilde{f}_1) + \cdots + N(\widetilde{f}_s)\right). \tag{5.11}$$

References

[1] R. F. Brown, *The Nielsen number of a fibre map*, Annals of Mathematics. Second Series **85** (1967), 483–493.

[2] ———, *The Lefschetz Fixed Point Theorem*, Scott, Foresman, Illinois, 1971.

[3] B. J. Jiang, *Lectures on Nielsen Fixed Point Theory*, Contemporary Mathematics, vol. 14, American Mathematical Society, Rhode Island, 1983.

[4] C. Y. You, *Fixed point classes of a fiber map*, Pacific Journal of Mathematics **100** (1982), no. 1, 217–241.

Jerzy Jezierski: Department of Mathematics, University of Agriculture, Nowoursynowska 159, 02 766 Warszawa, Poland

E-mail address: jezierski_acn@waw.pl

Journal of Applied Mathematics and Decision Sciences

Special Issue on
Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods**: artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning
- **Application fields**: asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects**: decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site http://www.hindawi.com/journals/jamds/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk