Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2007, Article ID 19745, 7 pages
doi:10.1155/2007/19745

Research Article

Diametrically Contractive Multivalued Mappings

S. Dhompongsa and H. Yingtaweesittikul

Received 1 March 2007; Accepted 2 May 2007

Recommended by Tomas Dominguez Benavides

Diametrically contractive mappings on a complete metric space are introduced by V. I. Istratescu. We extend and generalize this idea to multivalued mappings. An easy example shows that our fixed point theorem is more applicable than a former one obtained by H. K. Xu. A convergence theorem of Picard iteratives is also provided for multivalued mappings on hyperconvex spaces, thereby extending a Proinov's result.

Copyright © 2007 S. Dhompongsa and H. Yingtaweesittikul. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let (X,d) be a complete metric space. A mapping $T : X \to X$ is a *contraction* if for some $\alpha \in (0,1)$,

$$d(Tx,Ty) \le \alpha d(x,y) \quad \forall x,y \in X. \tag{1.1}$$

The mapping T is said to be *contractive* if

$$d(Tx,Ty) < d(x,y) \quad \forall x,y \in X,\ x \neq y. \tag{1.2}$$

By the well-known Banach's contraction principle, every contraction has a unique fixed point x_0 and the Picard iteration $\{T^n x\}$ converges to x_0 for every $x \in X$. Examples in [1, 2] show that a contractive mapping may fail to have a fixed point. However, a question of the existence of a fixed point is of interest. In fact, it has been left open the following question.

Question 1.1 [3]. Let M be a weakly compact subset of a Banach space and let $T : M \to M$ be contractive. Does T have a fixed point?

Istratescu [4] introduced a proper subclass of the class of the contractive mappings, whose elements are called the diametrically contractive mappings. Xu [2] proved, in the framework of Banach spaces, the following theorem.

Theorem 1.2 [2, Theorem 2.3]. *Let M be a weakly compact subset of a Banach space X and let $T : M \to M$ be a diametrically contractive mapping. Then T has a fixed point.*

The following problems raised in [2] had been answered in the negative way in [5].

Problem 1.3. Can we substitute "weakly compact" subset with "closed convex bounded" one in Theorem 1.2?

Problem 1.4. If T is diametrically contractive and x^* is the fixed point of T, do we have $T^n x \to x^*$ for all (or at least for some) $x \in M$?

In this paper, we weaken the condition in the definition of diametrically contractive mappings and obtain a corresponding fixed point theorem for nonself multivalued mappings. Moreover, we also apply a Proinov's fixed point theorem to a selection of a multivalued mapping with externally hyperconvex values and obtain its unique fixed point on a hyperconvex metric space.

2. Diametrically contractive mappings

In [4], Istratescu introduced a new class of mappings strictly lying between contractions and contractive mappings.

Definition 2.1. A mapping T on a complete metric space (X, d) is said to be *diametrically contractive* if $\delta(TA) < \delta(A)$ for all closed subsets A with $0 < \delta(A) < \infty$.

(Here $\delta(A) := \sup\{d(x, y) : x, y \in A\}$ is the diameter of $A \subset X$.)

In the following, we consider a multivalued version of mappings in Theorem 1.2. We also can weaken the condition required in Definition 2.1.

Let $\mathcal{F}(X)$ be the collection of nonempty closed subsets of X and let $\operatorname{Fix} T$ denote the set of fixed points of T. Recall that $TA = \bigcup_{a \in A} Ta$.

Theorem 2.2. *Let M be a weakly compact subset of a Banach space X and let $T : M \to \mathcal{F}(X)$, $Tx \cap M \neq \emptyset$ for all $x \in M$ and $\delta(TA \cap A) < \delta(A)$ for all closed sets A with $\delta(A) > 0$. Then T has a unique fixed point.*

Proof. The uniqueness of the fixed point is obvious. To prove the existence we consider the family $\mathcal{U} := \{A \subset M : A \text{ is a nonempty weakly compact subset of } M,\ TA \cap M \subset A\}$. Clearly, $\mathcal{U} \neq \emptyset$. Partially order $\mathcal{U}$ by saying that $A_1 \preceq A_2$ if $A_1 \supset A_2$ for $A_1, A_2 \in \mathcal{U}$. Every chain $\mathcal{C}$ in $\mathcal{U}$ has a finite intersection property, thus it has a nonempty intersection, that is, $B := \bigcap_{A \in \mathcal{C}} A \neq \emptyset$. Since $TB \cap M \subset TA \cap M \subset A$ for all $A \in \mathcal{C}$, $TB \cap M \subset B$, that is, $B \in \mathcal{U}$, and it is an upper bound of $\mathcal{C}$. Thus $\mathcal{U}$ has a maximal element, say A. Fix $x \in A$. As $A \in \mathcal{U}$ we see that $Tx \cap M \subset TA \cap M \subset A$. That is to say $Tx \cap A \neq \emptyset$ for all $x \in A$.

Put $A_0 = \overline{TA \cap A}^w$. Therefore $A_0 = \overline{TA \cap A}^w \subset \overline{TA \cap M}^w \subset \overline{A}^w = A$ and so $A_0 \subset A$. Moreover, we have $TA_0 \cap M \subset TA \cap M \subset A$. Therefore $TA_0 \cap M \subset TA \cap A \subset \overline{TA \cap A}^w = A_0$. Thus $A_0 \in \mathcal{U}$ and by maximality of A, we have $A = A_0 = \overline{TA \cap A}^w$. By lower semicontinuity of the norm of X we see that $\delta(A) = \delta(\overline{TA \cap A}^w) = \delta(TA \cap A)$. Since T is diametrically contractive we must have $\delta(A) = 0$ and A consists of exactly one point, say ξ. By the condition $\emptyset \neq TA \cap M \subset A$ we see that $\xi \in T\xi$, and we have a fixed point. □

The proof given above is a modification of the proof of Theorem 1.2. The following example shows that Theorem 2.2 is strictly stronger than Theorem 1.2.

Example 2.3. $M = [0,5]$, $T : M \to \mathbb{R}$ defined by $Tx = x + 1$ if $x \leq 3$, and $Tx = 4$ if $x > 3$. Now, if A is a closed subset of M, then there will be a, b in M such that $A \subset [a,b]$ and $\delta(A) = b - a$. If $[a,b] \subset [0,3]$, then $TA \subset [a+1, b+1]$ and $TA \cap A \subset [a+1, b]$. Thus $\delta(TA \cap A) \leq b - a - 1 < \delta(A)$. If $a \leq 3 \leq b$, then $TA \subset [a+1, 4]$ and therefore $\delta(TA \cap A) \leq 3 - a < b - a = \delta(A)$. The case when $[a,b] \subset [3,5]$, T clearly satisfies $\delta(TA \cap A) = 0 < \delta(A)$. Thus T has a fixed point by Theorem 2.2. Note that 4 is the unique fixed point of T. We observe that T does not satisfy the condition in Theorem 1.2 because $\delta(T[0,1]) = 1 = \delta([0,1])$.

Example 2.4. Let $Tx = [0, x - \log(x+1)]$ for $x \in [0,100]$. If A is a bounded closed subset of $[0,100]$, then for some $a, b > 0$ we have $A \subset [a,b]$, and $\delta(A) = b - a$. Clearly $TA \subset \bigcup_{x \in A} [0, x - \log(x+1)] \subset [0, b - \log(b+1)]$, and so $TA \cap A \subset [a, b - \log(b+1)]$. Therefore $\delta(TA \cap A) < \delta(A)$. 0 is the unique fixed point of T.

Next we will replace the diameter $\delta(A)$ of a set A by $\alpha(A)$, where α is the Kuratowski measure of noncompactness:

$$\alpha(A) = \inf\{\epsilon > 0 : A \text{ can be covered by finitely many sets with diameters } \leq \epsilon\}. \quad (2.1)$$

Definition 2.5. Let M be a nonempty subset of a metric space (X,d). A mapping $T : M \to 2^X$ is said to be *a k-set contraction* if, for each $A \subset M$ with A bounded, TA is bounded and $\alpha(TA) \leq k\alpha(A)$. If $\alpha(TA) < \alpha(A)$ for all such A, then T is said to be *α-condensing*.

Suppose that M is a bounded subset of a metric space (X,d). Then:

(i) $\mathrm{co}(M) = \bigcap\{B \subset X : B \text{ is a closed ball in } X \text{ such that } M \subset B\}$, and

(ii) M is said to be *subadmissible* [6], if for each $A \in \langle M \rangle$, $\mathrm{co}(A) \subset M$, where $\langle M \rangle$ denotes the class of all nonempty finite subsets of M.

For a nonempty subset M of X and a topological space Y, if two set-valued mappings $T, F : M \to 2^Y$ satisfy the condition $T(\mathrm{co}(A) \cap M) \subset F(A)$, for any $A \in \langle M \rangle$, then F is called a generalized KKM mapping with respect to T.

Let $T : M \to 2^Y$ be a set-valued mapping such that the family $\{\overline{Fx} : x \in M\}$ has the finite intersection property (where $\overline{Fx}$ denotes the closure of Fx) for each generalized KKM mapping $F : M \to 2^Y$ with respect to T, then we say that T has the KKM property. Denote

$$\mathrm{KKM}(M,Y) = \{T : M \longrightarrow 2^Y : T \text{ has the KKM property}\}. \quad (2.2)$$

Theorem 2.6 [7, Theorem 1]. *Let (X,d) be a complete metric space and let M be a nonempty bounded nearly subadmissible subset of X. If $T \in \mathrm{KKM}(M,M)$ is a k-set contraction, $0 < k < 1$, and closed with $\overline{TM} \subset M$, then T has a fixed point in M.*

The next result shows that we can replace k-set contractions in Theorem 2.6 by α-condensing mappings.

Theorem 2.7. *Let (X,d) be a complete metric space and let M be a nonempty bounded nearly subadmissible subset of X. If $T \in \mathrm{KKM}(M,M)$ is α-condensing, and closed with $\overline{TM} \subset M$, then T has a fixed point in M.*

In the course of the proof, we will apply the technique in the proof of the following lemma.

Lemma 2.8 [8, Lemma 2.2]. *Let F be a selfmapping of an arbitrary set Y and let $f : Y \to \mathbb{R}_+$ be a nonnegative valued function defined on Y. Suppose that the following conditions hold:*

(i) *there exists a function $\varphi \in \Phi_1$ (i.e., $\varphi : \mathbb{R}_+ \to \mathbb{R}_+$ satisfying: for any $\epsilon > 0$, there exists $\delta > \epsilon$ such that $\epsilon < t < \delta$ implies $\varphi(t) \le \epsilon$) such that $f(Fy) \le \varphi(f(y))$ for all $y \in Y$;*

(ii) *$f(y) > 0$ implies $f(Fy) < f(y)$ and $f(y) = 0$ implies $f(Fy) = 0$.*

Then $\lim f(F^n y) = 0$ for each $y \in Y$.

Proof of Theorem 2.7. We follow the proof of Theorem 2.6. Let $y \in M$ be any point, and let $M_0 = M$. Define $M_1 = \mathrm{co}(T(M_0) \cup \{y\}) \cap M$, and $M_{n+1} = \mathrm{co}(T(M_n) \cup \{y\}) \cap M$, for each n. Then

$$\alpha(M_{n+1}) \le \alpha(T(M_n)) < \alpha(M_n) \le \cdots < \alpha(M), \tag{2.3}$$

for each n (see [7]).

If we can prove that

$$\lim \alpha(M_n) = 0, \tag{2.4}$$

then the rest of the proof will follow the same lines as of Theorem 2.6. To achieve (2.4), we will apply the proof of Lemma 2.8. For each $t \in R_+$, let $A_t = \{M_n : \alpha(M_n) \le t\}$ and $B_t = \{\alpha(M_{n+1}) : M_n \in A_t\}$. From (2.3), it is seen that $B_t \ne \emptyset$ and bounded if $A_t \ne \emptyset$. Define $\varphi(t) = \sup B_t$ if $A_t \ne \emptyset$. Otherwise, put $\varphi(t) = 0$. We claim that $\varphi \in \Phi_1$. Consider the set A_ϵ for $\epsilon > 0$. If $\alpha(M_n) \le \epsilon$ for some n, let n_0 be the smallest such n. If $n_0 = 0$, then by (2.3) it is seen that $\varphi(t) \le \epsilon$ for all $t > \epsilon$. Otherwise, let $\delta = \alpha(M_{n_0-1})$. Thus $\delta > \epsilon$, and if $\epsilon < t < \delta$, then by (2.3) we have $\varphi(t) \le \alpha(M_{n_0+1}) < \alpha(M_{n_0}) \le \epsilon$. Therefore $\varphi \in \Phi_1$. We now prove (2.4).

Clearly, we have

$$\alpha(M_{n+1}) \le \varphi(\alpha(M_n)), \quad \alpha(M_{n+1}) < \alpha(M_n) \quad \text{by (2.3) for each } n. \tag{2.5}$$

It follows from (2.3) that $\{\alpha(M_n)\}$ is strictly decreasing, hence it converges to some $\epsilon \ge 0$. Suppose $\epsilon > 0$. Since $\varphi \in \Phi_1$, we have for some $\delta > \epsilon$, $\varphi(t) \le \epsilon$ for all $t \in (\epsilon, \delta)$. Choose n_0 so that $\epsilon < \alpha(M_{n_0}) < \delta$. Thus $\varphi(\alpha(M_{n_0})) \le \epsilon$. But then (2.5) implies $\alpha(M_n) \le \epsilon$ for all $n > n_0$ which contradicts to (2.3). Hence (2.4) follows. □

3. Picard iteratives for multivalued mappings on hyperconvex metric spaces

A metric space (X,d) is *hyperconvex* if for any family of points $\{x_\alpha\}$ in X and any family of positive numbers $\{r_\alpha\}$ satisfying $d(x_\alpha,x_\beta)\le r_\alpha+r_\beta$, we have $\bigcap_\alpha B(x_\alpha,r_\alpha)\neq\emptyset$ where $B(x,r)$ is the closed ball with center at x and radius r. A subset E of X is said to be *externally hyperconvex* if for any of those families $\{x_\alpha\}$, $\{r_\alpha\}$ with $d(x_\alpha,x_\beta)\le r_\alpha+r_\beta$ and $\operatorname{dist}(x_\alpha,E)\le r_\alpha$, we have $\bigcap_\alpha B(x_\alpha,r_\alpha)\cap E\neq\emptyset$. The class of all externally hyperconvex subsets of X will be denoted by $\mathscr{E}(X)$. Let H be the Hausdorff metric.

Let t be a single-valued selfmapping on a metric space (X,d). A fixed point of t is said to be *contractive* (cf. [9]) if all Picard iteratives of t converge to this fixed point. A selfmapping t on a metric space (X,d) is said to be *asymptotically regular* (cf. [10]) if $\lim d(t^n(x),t^{n+1}(x))=0$ for each x in X. Extend the concept naturally to multivalued mappings with the Hausdorff metric taken into action.

Theorem 3.1 [8, Theorem 4.1]. *Let t be a continuous and asymptotically regular selfmapping on a complete metric space satisfying the following conditions:*

(i) *there exists $\varphi\in\Phi_1$ such that $d(t(x),t(y))\le\varphi(D(x,y))$ for all $x,y\in X$;*

(ii) *$d(t(x),t(y))<D(x,y)$ for all $x,y\in X$ with $x\neq y$.*

Then t has a contractive fixed point. Here $D(x,y)=d(x,y)+r[d(x,t(x))+d(y,t(y))]$, $r\ge 0$.

Replacing D by d, we present a multivalued version of Theorem 3.1 on a special setting, namely, on the class of hyperconvex metric spaces.

Theorem 3.2. *Let (X,d) be a bounded hyperconvex metric space, and let $T:X\to\mathscr{E}(X)$ be asymptotically regular satisfying the following conditions:*

(i) *there exists $\varphi\in\Phi_1$ such that $\varphi(x)\le x$, $\varphi(x+y)\le\varphi(x)+\varphi(y)$, $\varphi(x)=0$ if and only if $x=0$, and $H(Tx,Ty)\le\varphi(d(x,y))$ for all x,y in X;*

(ii) *$H(Tx,Ty)<d(x,y)$ for all $x,y\in X$ with $x\neq y$.*

Then, if $\delta(T^nx)\to 0$ for each $x\in X$, T has a contractive fixed point. That is, there exists a unique point ξ in X such that, for each $x\in X$, $T^nx\to\{\xi\}=\operatorname{Fix}T$.

Proof. The uniqueness of the fixed point is evident. We are going to find a selection $t:X\to X$ so that $t(x)\in Tx$ for all $x\in X$ and t satisfies the conditions in Theorem 3.1. Thus, there is a point ξ in $\operatorname{Fix}t$ satisfying $t^n(x)\to\xi$ for all $x\in X$. To find a selection t, we apply Zorn's lemma to the family $\mathscr{F}=\{(A,t):\emptyset\neq A\subset X,\ t:A\to A$ asymptotically regular, $t(a)\in Ta$ for all $a\in A$, and t satisfies (i) and (ii) in Theorem 3.1$\}$. Partially order $\mathscr{F}$ by $(A_1,t_1)\preceq(A_2,t_2)$ if $A_1\subset A_2$ and $t_2|_{A_1}=t_1$.

Suppose $A=\emptyset$ or $(A,t)\in\mathscr{F}$ and $x_0\in X\setminus A$. We will define a countable set $\{x_0,x_1,x_2,\dots\}$, possibly finite, and an extension function t^* of t over $A\cup\{x_0,x_1,x_2,\dots\}$ so that $(A\cup\{x_0,x_1,x_2,\dots\},t^*)\in\mathscr{F}$. Let $x_0,x_1,\dots,x_n\in X\setminus A$ have been defined for some $n\ge 0$ so that, for $1\le k\le n$, $x_k\in Tx_{k-1}$, and when $n\ge 2$, $d(x_{i+1},x_{j+1})\le\varphi(d(x_i,x_j))$ and $d(x_{i+1},x_{j+1})<d(x_i,x_j)$ for $i<j$ in $\{1,\dots,n-1\}$. Moreover, $d(t(x),x_{i+1})\le\varphi(d(x,x_i))$ and $d(t(x),x_{i+1})<d(x,x_i)$ for $i\in\{1,\dots,n-1\}$ and all $x\in A$.

Put $r_k = \varphi(d(x_{k-1},x_n))$, and $r_{t(x)} = \varphi(d(x,x_n))$ for each $1 \le k \le n$, and for all $x \in A$. Thus, for $1 \le k \le n$, $x \in A$, and for $i < j$ in $\{1,\dots,n-1\}$,

(i) $\operatorname{dist}(x_k, Tx_n) \le H(Tx_{k-1}, Tx_n) \le \varphi(d(x_{k-1},x_n)) = r_k$,

(ii) $\operatorname{dist}(t(x), Tx_n) \le H(Tx, Tx_n) \le r_{t(x)}$,

(iii) $d(t(x),x_k) \le \varphi(d(x,x_{k-1})) \le \varphi(d(x,x_n)) + \varphi(d(x_{k-1},x_n)) = r_{t(x)} + r_k$, and

(iv) $d(x_i,x_j) \le \varphi(d(x_{i-1},x_{j-1})) \le \varphi(d(x_{i-1},x_n)) + \varphi(d(x_{j-1},x_n)) = r_i + r_j$.

Finally, for $x, y \in A$, $d(t(x),t(y)) \le \varphi(d(x,y)) \le \varphi(d(x,x_n)) + \varphi(d(y,x_n)) = r_{t(x)} + r_{t(y)}$. Therefore there exists a point $x_{n+1} \in \bigcap_{x\in A} B(t(x), r_{t(x)}) \bigcap_{k=1}^{n} B(x_k, r_k) \cap Tx_n$. The point x_{n+1} has the following property: for $k \le n$, $d(x_k,x_{n+1}) \le r_k = \varphi(d(x_{k-1},x_n))$, and for each $x \in A$, $d(t(x),x_{n+1}) \le r_{t(x)} = \varphi(d(x,x_n))$. Clearly, $d(x_k,x_{n+1}) < d(x_{k-1},x_n)$ and $d(t(x),x_{n+1}) < d(x,x_n)$.

If $x_{n+1} \in A$, the process terminates. Otherwise, we obtain a subset $\{x_0,x_1,\dots,x_n,\dots\}$ of $X \setminus A$ satisfying the conditions (i) and (ii) in Theorem 3.1 where we extend t to t^* by defining $t^*(x_n) = x_{n+1}$ for $n \ge 0$. Thus $(A \cup \{x_0,x_1,\dots\}, t^*) \in \mathcal{F}$.

In summary, the above argument shows that, if $A = \emptyset$, then $(\{x_0,x_1,\dots\}, t^*) \in \mathcal{F}$, that is, $\mathcal{F} \ne \emptyset$. On the other hand if (A,t) is a maximal element in $\mathcal{F}$ (by Zorn's lemma), we must have $A = X$, that is, (X,t) belongs to $\mathcal{F}$ for some t. Apply Theorem 3.1, to conclude that there exists a fixed point ξ of t such that $t^n(x) \to \xi$ for each $x \in X$. Consequently, $T^n x \to \{\xi\}$ for each $x \in X$ and $\operatorname{Fix} T = \{\xi\}$. □

Acknowledgments

The authors would like to thank the referees for valuable comments. They also would like to thank the Thailand Research Fund (Grant BRG4780013) and the Development and Promotion of Science and Technology Talent Project (DPST) for their support.

References

[1] W. A. Kirk, "A fixed point theorem for mappings which do not increase distances," *The American Mathematical Monthly*, vol. 72, no. 9, pp. 1004–1006, 1965.

[2] H.-K. Xu, "Diametrically contractive mappings," *Bulletin of the Australian Mathematical Society*, vol. 70, no. 3, pp. 463–468, 2004.

[3] K. Deimling, *Multivalued Differential Equations*, vol. 1 of *de Gruyter Series in Nonlinear Analysis and Applications*, Walter de Gruyter, Berlin, Germany, 1992.

[4] V. I. Istratescu, "Some fixed theorems for convex contraction mappings and mappings with convex diminishing diameters. IV nonexpansive diameter mappings in uniformly convex spaces," preliminary report. Abstract of the American Mathematiccal Society, 82T-46-316, 1982.

[5] M. Baronti, E. Casini, and P. L. Papini, "Diametrically contractive maps and fixed points," *Fixed Point Theory and Applications*, vol. 2006, Article ID 79075, 8 pages, 2006.

[6] A. Amini, M. Fakhar, and J. Zafarani, "KKM mappings in metric spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 60, no. 6, pp. 1045–1052, 2005.

[7] C.-M. Chen, "KKM property and fixed point theorems in metric spaces," *Journal of Mathematical Analysis and Applications*, vol. 323, no. 2, pp. 1231–1237, 2006.

[8] P. D. Proinov, "Fixed point theorems in metric spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 3, pp. 546–557, 2006.

[9] S. Leader, "Fixed points for operators on metric spaces with conditional uniform equivalence of orbits," *Journal of Mathematical Analysis and Applications*, vol. 61, no. 2, pp. 466–474, 1977.

[10] F. E. Browder and W. V. Petryshyn, "The solution by iteration of nonlinear functional equations in Banach spaces," *Bulletin of the American Mathematical Society*, vol. 72, pp. 571–575, 1966.

S. Dhompongsa: Department of Mathematics, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand
Email address: sompongd@chiangmai.ac.th

H. Yingtaweesittikul: Department of Mathematics, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand
Email address: g4825119@cm.edu

Journal of Applied Mathematics and Decision Sciences

Special Issue on
Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods**: artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning
- **Application fields**: asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects**: decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site http://www.hindawi.com/journals/jamds/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk