Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2007, Article ID 39465, 6 pages
doi:10.1155/2007/39465

Research Article

A Note on Asymptotic Contractions

Marina Arav, Francisco Eduardo Castillo Santos,
Simeon Reich, and Alexander J. Zaslavski

Received 27 August 2006; Revised 16 October 2006; Accepted 16 October 2006

Recommended by Brailey Sims

We provide sufficient conditions for the iterates of an asymptotic contraction on a complete metric space X to converge to its unique fixed point, uniformly on each bounded subset of X.

Copyright © 2007 Marina Arav et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let (X,d) be a complete metric space. The following theorem is the main result of Chen [1]. It improves upon Kirk's original theorem [2]. In this connection, see also [3, 4].

Theorem 1.1. *Let $T : X \to X$ be such that*

$$d(T^n x, T^n y) \le \phi_n(d(x,y)) \tag{1.1}$$

for all $x, y \in X$ and all natural numbers n, where $\phi_n : [0,\infty) \to [0,\infty)$ and $\lim_{n\to\infty} \phi_n = \phi$, uniformly on any bounded interval $[0,b]$. Suppose that ϕ is upper semicontinuous and that $\phi(t) < t$ for all $t > 0$. Furthermore, suppose that there exists a positive integer n_* such that ϕ_{n_*} is upper semicontinuous and $\phi_{n_*}(0) = 0$. If there exists $x_0 \in X$ which has a bounded orbit $O(x_0) = \{x_0, Tx_0, T^2x_0, \ldots\}$, then T has a unique fixed point $x_* \in X$ and $\lim_{n\to\infty} T^n x = x_*$ for all $x \in X$.

Note that Theorem 1.1 does not provide us with uniform convergence of the iterates of T on bounded subsets of X, although this does hold for many classes of mappings of contractive type (e.g., [5, 6]). This property is important because it yields stability of the

convergence of iterates even in the presence of computational errors [7]. In the present paper we show that this conclusion can be derived in the setting of Theorem 1.1. To this end, we first prove a somewhat more general result (Theorem 1.2) which, when combined with Theorem 1.1, yields our strengthening of Chen's result (Theorem 1.3).

THEOREM 1.2. *Let $x_* \in X$ be a fixed point of $T : X \to X$. Assume that*

$$d(T^n x, x_*) \le \phi_n(d(x, x_*)) \quad \forall x \in X \text{ and all natural numbers } n, \tag{1.2}$$

where $\phi_n : [0,\infty) \to [0,\infty)$ and $\lim_{n\to\infty} \phi_n = \phi$, uniformly on any bounded interval $[0,b]$. Suppose that ϕ is upper semicontinuous and that $\phi(t) < t$ for all $t > 0$. Then $T^n x \to x_*$ as $n \to \infty$, uniformly on each bounded subset of X.

THEOREM 1.3. *Let $T : X \to X$ be such that*

$$d(T^n x, T^n y) \le \phi_n(d(x, y)) \tag{1.3}$$

for all $x, y \in X$ and all natural numbers n, where $\phi_n : [0,\infty) \to [0,\infty)$ and $\lim_{n\to\infty} \phi_n = \phi$, uniformly on any bounded interval $[0,b]$. Suppose that ϕ is upper semicontinuous and that $\phi(t) < t$ for all $t > 0$. Furthermore, suppose that there exists a positive integer n_* such that ϕ_{n_*} is upper semicontinuous and $\phi_{n_*}(0) = 0$. If there exists $x_0 \in X$ which has a bounded orbit $O(x_0) = \{x_0, Tx_0, T^2x_0, \ldots\}$, then T has a unique fixed point $x_* \in X$ and $\lim_{n\to\infty} T^n x = x_*$, uniformly on each bounded subset of X.

2. Proof of Theorem 1.2

We may assume without loss of generality that $\phi(0) = 0$ and $\phi_n(0) = 0$ for all integers $n \ge 1$.

For each $x \in X$ and each $r > 0$, set

$$B(x, r) = \{y \in X : d(x, y) \le r\}. \tag{2.1}$$

We first prove three lemmas.

LEMMA 2.1. *Let $K > 0$. Then there exists a natural number $\overline{q}$ such that for all integers $s \ge \overline{q}$,*

$$T^s(B(x_*, K)) \subset B(x_*, K + 1). \tag{2.2}$$

Proof. There exists a natural number $\overline{q}$ such that for all integers $s \ge \overline{q}$,

$$|\phi_s(t) - \phi(t)| < 1 \quad \forall t \in [0, K]. \tag{2.3}$$

Let $s \geq \overline{q}$ be an integer. Then for all $x \in B(x_*, K)$,

$$d(T^s x, x_*) \leq \phi_s(d(x, x_*)) < \phi(d(x, x_*)) + 1 < d(x, x_*) + 1 < K + 1. \tag{2.4}$$

Lemma 2.1 is proved. □

Lemma 2.2. *Let $0 < \epsilon_1 < \epsilon_0$. Then there exists a natural number q such that for each integer $j \geq q$,*

$$T^j(B(x_*, \epsilon_1)) \subset B(x_*, \epsilon_0). \tag{2.5}$$

Proof. There exists an integer $q \geq 1$ such that for each integer $j \geq q$,

$$|\phi_j(t) - \phi(t)| < (\epsilon_0 - \epsilon_1)/2 \quad \forall t \in [0, \epsilon_0]. \tag{2.6}$$

Assume that

$$j \in \{q, q+1, \ldots\}, \qquad x \in B(x_*, \epsilon_1). \tag{2.7}$$

By (1.2) and (2.6),

$$\begin{aligned} d(T^j x, x_*) &\leq \phi_j(d(x, x_*)) < \phi(d(x, x_*)) + \frac{(\epsilon_0 - \epsilon_1)}{2} \\ &\leq \epsilon_1 + \frac{(\epsilon_0 - \epsilon_1)}{2} = \frac{(\epsilon_0 + \epsilon_1)}{2}. \end{aligned} \tag{2.8}$$

Lemma 2.2 is proved. □

Lemma 2.3. *Let $K, \epsilon > 0$. Then there exists a natural number q such that for each $x \in B(x_*, K)$,*

$$\min\{d(T^j x, x_*) : j = 1, \ldots, q\} \leq \epsilon. \tag{2.9}$$

Proof. By Lemma 2.1, there is a natural number $\overline{q}$ such that

$$T^n(B(x_*, K)) \subset B(x_*, K+1) \text{ for all natural numbers } n \geq \overline{q}. \tag{2.10}$$

We may assume without loss of generality that $\epsilon < K/8$. Since the function $t - \phi(t)$, $t \in (0, \infty)$, is lower semicontinuous and positive, there is

$$\delta \in \left(0, \frac{\epsilon}{8}\right) \tag{2.11}$$

such that

$$t - \phi(t) \geq 2\delta \quad \forall t \in \left[\frac{\epsilon}{2}, K+1\right]. \tag{2.12}$$

There is a natural number $s \geq \overline{q}$ such that

$$\left|\phi(t) - \phi_s(t)\right| \leq \delta \quad \forall t \in [0, K+1]. \tag{2.13}$$

By (2.12) and (2.13), we have, for all $t \in [\epsilon/2, K+1]$,

$$\phi_s(t) \leq \phi(t) + \delta \leq t - 2\delta + \delta = t - \delta. \tag{2.14}$$

In view of (2.13) and (2.11), we have, for all $t \in [0, \epsilon/2]$,

$$\phi_s(t) \leq \phi(t) + \delta \leq t + \delta \leq \frac{\epsilon}{2} + \delta < \frac{3}{4}\epsilon. \tag{2.15}$$

Choose a natural number p such that

$$p > 4 + \delta^{-1}(K+1). \tag{2.16}$$

Let

$$x \in B(x_*, K). \tag{2.17}$$

We will show that

$$\min\{d(T^j x, x_*) : j = 1, 2, \ldots, ps\} \leq \epsilon. \tag{2.18}$$

Let us assume the contrary. Then

$$d(T^j x, x_*) > \epsilon \quad \forall j = s, \ldots, ps. \tag{2.19}$$

By (2.17) and (2.10),

$$T^j x \in B(x_*, K+1), \quad j = s, \ldots, ps. \tag{2.20}$$

Let a natural number i satisfy $i \leq p-1$. By (2.19) and (2.20),

$$d(T^{is} x, x_*) > \epsilon, \qquad d(T^{is} x, x_*) \leq K+1. \tag{2.21}$$

It follows from (1.2), (2.21), and (2.14) that

$$d(T^s(T^{is} x), x_*) \leq \phi_s(d(T^{is} x, x_*)) \leq d(T^{is} x, x_*) - \delta. \tag{2.22}$$

Thus for each natural number $i \le p-1$,

$$d(T^{(i+1)s}x, x_*) \le d(T^{is}x, x_*) - \delta. \tag{2.23}$$

This inequality implies that

$$d(T^{ps}x, x_*) \le d(T^{(p-1)s}x, x_*) - \delta \le \cdots \le d(T^{s}x, x_*) - (p-1)\delta. \tag{2.24}$$

When combined with (2.20) and (2.16), this implies, in turn, that

$$d(T^{ps}x, x_*) \le K + 1 - (p-1)\delta < 0. \tag{2.25}$$

The contradiction we have reached proves (2.18) and completes the proof of Lemma 2.3. □

Completion of the proof of Theorem 1.2. Let $K, \epsilon > 0$. Choose $\epsilon_1 \in (0, \epsilon)$. By Lemma 2.2, there exists a natural number q_1 such that

$$T^j(B(x_*, \epsilon_1)) \subset B(x_*, \epsilon) \text{ for all integers } j \ge q_1. \tag{2.26}$$

By Lemma 2.3, there exists a natural number q_2 such that

$$\min\{d(T^j x, x_*) : j = 1, \ldots, q_2\} \le \epsilon_1 \quad \forall x \in B(x_*, K). \tag{2.27}$$

Assume that

$$x \in B(x_*, K). \tag{2.28}$$

By (2.27), there is a natural number $j_1 \le q_2$ such that

$$d(T^{j_1}x, x_*) \le \epsilon_1. \tag{2.29}$$

In view of (2.29) and (2.26),

$$T^j(T^{j_1}x) \in B(x_*, \epsilon) \text{ for all integers } j \ge q_1. \tag{2.30}$$

Inclusion (2.30) and the inequality $j_1 \le q_2$ now imply that

$$T^i x \in B(x_*, \epsilon) \text{ for all integers } i \ge q_1 + q_2. \tag{2.31}$$

Theorem 1.2 is proved.

Acknowledgments

Part of the first author's research was carried out when she was visiting the Technion—Israel Institute of Technology. The third author was partially supported by the Fund for the Promotion of Research at the Technion and by the Technion VPR Fund—B. and G. Greenberg Research Fund (Ottawa).

References

[1] Y.-Z. Chen, "Asymptotic fixed points for nonlinear contractions," *Fixed Point Theory and Applications*, vol. 2005, no. 2, pp. 213–217, 2005.
[2] W. A. Kirk, "Fixed points of asymptotic contractions," *Journal of Mathematical Analysis and Applications*, vol. 277, no. 2, pp. 645–650, 2003.
[3] I. D. Arandelović, "On a fixed point theorem of Kirk," *Journal of Mathematical Analysis and Applications*, vol. 301, no. 2, pp. 384–385, 2005.
[4] J. Jachymski and I. Jóźwik, "On Kirk's asymptotic contractions," *Journal of Mathematical Analysis and Applications*, vol. 300, no. 1, pp. 147–159, 2004.
[5] F. E. Browder, "On the convergence of successive approximations for nonlinear functional equations," *Indagationes Mathematicae*, vol. 30, pp. 27–35, 1968.
[6] E. Rakotch, "A note on contractive mappings," *Proceedings of the American Mathematical Society*, vol. 13, no. 3, pp. 459–465, 1962.
[7] D. Butnariu, S. Reich, and A. J. Zaslavski, "Asymptotic behavior of inexact orbits for a class of operators in complete metric spaces," to appear in Journal of Applied Analysis.

Marina Arav: Department of Mathematics and Statistics, Georgia State University,
Atlanta, GA 30303, USA
Email address: matmxa@langate.gsu.edu

Francisco Eduardo Castillo Santos: School of Mathematical and Physical Sciences,
The University of Newcastle, Newcastle, NSW 2308, Australia
Email address: francisco.castillosantos@studentmail.newcastle.edu.au

Simeon Reich: Department of Mathematics, The Technion–Israel Institute of Technology,
Haifa 32000, Israel
Email address: sreich@techunix.technion.ac.il

Alexander J. Zaslavski: Department of Mathematics, The Technion–Israel Institute of Technology,
Haifa 32000, Israel
Email address: ajzasl@tx.technion.ac.il

Mathematical Problems in Engineering

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at http://www.hindawi.com/journals/mpe/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/ according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru