Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2007, Article ID 61434, 5 pages
doi:10.1155/2007/61434

Review Article

Remarks of Equivalence among Picard, Mann, and Ishikawa Iterations in Normed Spaces

Xue Zhiqun

Received 1 April 2007; Revised 16 April 2007; Accepted 21 June 2007

Recommended by J. R. L. Webb

We show that the convergence of Picard iteration is equivalent to the convergence of Mann iteration schemes for various Zamfirescu operators. Our result extends of Soltuz (2005).

Copyright © 2007 Xue Zhiqun. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let E be a real normed space, D a nonempty convex subset of E, and T a self-map of D, let $p_0, u_0, x_0 \in D$. The Picard iteration is defined by

$$p_{n+1} = Tp_n, \quad n \geq 0. \tag{1.1}$$

The Mann iteration is defined by

$$u_{n+1} = (1 - a_n)u_n + a_n Tu_n, \quad n \geq 0. \tag{1.2}$$

The Ishikawa iteration is defined by

$$\begin{aligned} y_n &= (1 - b_n)x_n + b_n Tx_n, \quad n \geq 0, \\ x_{n+1} &= (1 - a_n)x_n + a_n Ty_n, \quad n \geq 0, \end{aligned} \tag{1.3}$$

where $\{a_n\}, \{b_n\}$ are sequences of positive numbers in $[0, 1]$. Obviously, for $a_n = 1$, the Mann iteration (1.2) reduces to the Picard iteration, and for $b_n = 0$, the Ishikawa iteration (1.3) reduces to the Mann iteration (1.2).

Definition 1.1 [1, Definition 1]. Let $T : D \to D$ be a map for which there exist real numbers a, b, c satisfying $0 < a < 1$, $0 < b < 1/2$, $0 < c < 1/2$. Then T is called a Zamfirescu operator

if, for each pair x, y in D, T satisfies at least one of the following conditions given in (1)–(3):

(1) $\|Tx - Ty\| \leq a\|x - y\|$;

(2) $\|Tx - Ty\| \leq b(\|x - Tx\| + \|y - Ty\|)$;

(3) $\|Tx - Ty\| \leq c(\|x - Ty\| + \|y - Tx\|)$.

It is easy to show that every Zamfirescu operator T satisfies the inequality

$$\|Tx - Ty\| \leq \delta\|x - y\| + 2\delta\|x - Tx\| \tag{1.4}$$

for all $x, y \in D$, where $\delta = \max\{a, b/(1-b), c/(1-c)\}$ with $0 < \delta < 1$ (See Şoltuz [1]). Recently, Şoltuz [1] had studied that the equivalence of convergence for Picard, Mann, and Ishikawa iterations, and proved the following results.

Theorem 1.2 [1, Theorem 1]. *Let X be a normed space, D a nonempty, convex, closed subset of X, and $T : D \to D$ an operator satisfying condition Z (Zamfirescu operator). If $u_0 = x_0 \in D$, let $\{u_n\}_{n=0}^{\infty}$ be defined by (1.2) for $u_0 \in D$, and let $\{x_n\}_{n=0}^{\infty}$ be defined by (1.3) for $x_0 \in D$ with $\{a_n\}$ in $[0,1]$ satisfying $\sum_{n=0}^{\infty} a_n = \infty$. Then the following are equivalent:*

(i) *the Mann iteration (1.2) converges to the fixed point of T;*

(ii) *the Ishikawa iteration (1.3) converges to the fixed point of T.*

Theorem 1.3 [1, Theorem 2]. *Let X be a normed space, D a nonempty, convex, closed subset of X, and $T : D \to D$ an operator satisfying condition Z (Zamfirescu operator). If $u_0 = p_0 \in D$, let $\{p_n\}_{n=0}^{\infty}$ be defined by (1.1) for $p_0 \in D$, and let $\{u_n\}_{n=0}^{\infty}$ be defined by (1.2) for $u_0 \in D$ with $\{a_n\}$ in $[0,1]$ satisfying $\sum_{n=0}^{\infty} a_n = \infty$ and $a_n \to 0$ as $n \to \infty$. Then*

(i) *if the Mann iteration (1.2) converges to x^* and $\lim_{n\to\infty}(\|u_{n+1} - u_n\|/a_n) = 0$, then the Picard iteration (1.1) converges to x^*,*

(ii) *if the Picard iteration (1.1) converges to x^* and $\lim_{n\to\infty}(\|p_{n+1} - p_n\|/a_n) = 0$, then the Mann iteration (1.2) converges to x^*.*

However, in the above-mentioned theorem, it is unnecessary that, for two conditions, $\lim_{n\to\infty}(\|u_{n+1} - u_n\|/a_n) = 0$ and $\lim_{n\to\infty}(\|p_{n+1} - p_n\|/a_n) = 0$. The aim of this paper is to show that the convergence of Picard iteration schemes is equivalent to the convergence of the Mann iteration for Zamfirescu operators in normed spaces. The result improves ones announced by Şoltuz [1, Theorem 2]. We will use a special case of the following lemma.

Lemma 1.4 [2]. *Let $\{a_n\}$ and $\{\sigma_n\}$ be nonnegative real sequences satisfying the following inequality:*

$$a_{n+1} \leq (1 - \lambda_n)a_n + \sigma_n, \tag{1.5}$$

where $\lambda_n \in (0,1)$, for all $n \geq n_0$, $\sum_{n=1}^{\infty} \lambda_n = \infty$, and $\sigma_n/\lambda_n \to 0$ as $n \to \infty$. Then $\lim_{n\to\infty} a_n = 0$.

This lemma is apparently due to Vasilen, it is given as [2, Lemma 2.3.6, page 96]. It was rediscovered with a different proof by Weng [3].

2. Main results

Theorem 2.1. *Let E be a normed space, D a nonempty closed convex subset of E, and $T : D \to D$ a Zamfirescu operator. Suppose that T has a fixed point $q \in D$. Let $\{p_n\}_{n=0}^{\infty}$ be defined by (1.1) for $p_0 \in D$, and let $\{u_n\}_{n=0}^{\infty}$ be defined by (1.2) for $u_0 \in D$ with $\{a_n\}$ in $[0,1]$ and satisfying $\sum_{n=0}^{\infty} a_n = \infty$. Then the following are equivalent:*

(i) *the Picard iteration (1.1) converges to the fixed point of T;*

(ii) *the Mann iteration (1.2) converges to the fixed point of T.*

Proof. Let q be a fixed point of T. We will prove (ii)⇒(i). Suppose that $\|u_n - q\| \to 0$ as $n \to \infty$. Applying (1.1) and (1.2), we have

$$\begin{aligned} ||u_{n+1} - p_{n+1}|| &\le (1 - a_n)||u_n - Tp_n|| + a_n||Tu_n - Tp_n|| \\ &\le (1 - a_n)||u_n - Tu_n|| + ||Tu_n - Tp_n||. \end{aligned} \tag{2.1}$$

Using (1.4) with $x = u_n$, $y = p_n$, we have

$$||Tu_n - Tp_n|| \le \delta||u_n - p_n|| + 2\delta||u_n - Tu_n||. \tag{2.2}$$

Therefore, from (2.1), we get

$$\begin{aligned} ||u_{n+1} - p_{n+1}|| &\le \delta\|u_n - p_n\| + (1 - a_n + 2\delta)||u_n - Tu_n|| \\ &\le \delta||u_n - p_n|| + (1 - a_n + 2\delta)(||u_n - q|| + ||Tu_n - Tq||) \\ &\le \delta||u_n - p_n|| + (1 - a_n + 2\delta)(1 + \delta)||u_n - q||, \end{aligned} \tag{2.3}$$

denoted by $A_n = \|u_n - p_n\|$, $\delta = 1 - \lambda$, and $B_n = (1 - a_n + 2\delta)(1 + \delta)\|u_n - q\|$. By Lemma 1.4, we obtain $A_n = \|u_n - p_n\| \to 0$ as $n \to \infty$. Hence by $\|p_n - q\| \le \|u_n - p_n\| + \|u_n - q\|$, we get $\|p_n - q\| \to 0$ as $n \to \infty$.

Next, we will prove (i)⇒(ii), that is, if the Picard iteration converges, then the Mann iteration does too. Now by using Picard iteration (1.1) and Mann iteration (1.2), we have

$$\begin{aligned} ||u_{n+1} - p_{n+1}|| &\le (1 - a_n)||u_n - Tp_n|| + a_n||Tu_n - Tp_n|| \\ &\le (1 - a_n)||u_n - p_n|| + (1 - a_n)||p_n - Tp_n|| + a_n||Tu_n - Tp_n|| \\ &\le (1 - a_n)||u_n - p_n|| + (1 - a_n)(||p_n - q|| + ||Tp_n - Tq||) + a_n||Tu_n - Tp_n||. \end{aligned} \tag{2.4}$$

On using (1.4) with $x = p_n$, $y = u_n$, we get

$$\begin{aligned} ||Tu_n - Tp_n|| &\le a_n\delta||u_n - p_n|| + 2a_n\delta||p_n - Tp_n|| \\ &\le a_n\delta||u_n - p_n|| + 2a_n\delta(||p_n - q|| + ||Tp_n - Tq||). \end{aligned} \tag{2.5}$$

Again, using (1.4) with $x = q$, $y = p_n$, we get

$$||Tp_n - Tq|| \leq \delta||p_n - q||. \tag{2.6}$$

Hence by (2.4)–(2.6), we obtain

$$\begin{aligned}
||u_{n+1} - p_{n+1}|| &\leq (1-(1-\delta)a_n)||u_n - p_n|| + (1 - a_n + 2a_n\delta)(1+\delta)||p_n - q|| \\
&\leq (1-(1-\delta)a_n)||u_n - q|| + (1 - a_n + 2a_n\delta)(2+\delta)||p_n - q|| \\
&\leq (1-\lambda a_n)(1-\lambda a_{n-1})||u_{n-1} - q|| + (1 - a_n + 2a_n\delta)(2+\delta)||p_n - q|| \\
&\leq (1-\lambda a_n)(1-\lambda a_{n-1})\cdots(1-\lambda a_0)||u_0 - q|| \\
&\quad + (1 - a_n + 2a_n\delta)(2+\delta)||p_n - q|| \\
&\leq \exp\left(-\lambda\sum_{i=0}^{n} a_i\right)||u_0 - q|| + (1 - a_n + 2a_n\delta)(2+\delta)||p_n - q||,
\end{aligned} \tag{2.7}$$

where $1 - \delta = \lambda$. Since $\sum_{n=0}^{\infty} a_n = \infty$ and $\|p_n - q\| \to 0$ as $n \to \infty$, hence $\|u_n - p_n\| \to 0$ as $n \to \infty$. And thus, $\|u_n - q\| \leq \|u_n - p_n\| + \|p_n - q\| \to 0$ as $n \to \infty$. This completes the proof. □

Remark 2.2. Theorem 2.1 improves [1, Theorem 2] in the following sense.

(1) Both hypotheses $\lim_{n\to\infty}(\|u_{n+1} - u_n\|/a_n) = 0$ and $\lim_{n\to\infty}(\|p_{n+1} - p_n\|/a_n) = 0$ have been removed, and the conclusion remains valid.

(2) The assumption that $u_0 = p_0$ in [1] is superfluous.

Theorem 2.3. *Let E be a normed space, D a nonempty closed convex subset of E, and $T : D \to D$ a Zamfirescu operator. Suppose that T has a fixed point $q \in D$. Let $\{p_n\}_{n=0}^{\infty}$ be defined by (1.1) for $p_0 \in D$, and let $\{x_n\}_{n=0}^{\infty}$ be defined by (1.3) for $x_0 \in D$ with $\{a_n\}$ and $\{b_n\}$ in $[0,1]$ and satisfying $\sum_{n=0}^{\infty} a_n = \infty$. Then the following are equivalent:*

(i) *the Picard iteration (1.1) converges to the fixed point of T;*

(ii) *the Ishikawa iteration (1.3) converges to the fixed point of T.*

Remark 2.4. As previously suggested, Theorem 2.3 reproduces exactly [1, Theorem 1]. Therefore we have the following conclusion: Picard iteration converges to the fixed point of $T \Leftrightarrow$ Mann iteration converges to the fixed point of $T \Leftrightarrow$ Ishikawa iteration converges to the fixed point of T.

Acknowledgments

The author would like to thank the referee and the editor for their careful reading of the manuscript and their many valuable comments and suggestions. The project is supported by the National Science Foundation of China and Shijiazhuang Railway Institute Sciences Foundation.

References

[1] Ş. M. Şoltuz, "The equivalence of Picard, Mann and Ishikawa iterations dealing with quasi-contractive operators," *Mathematical Communications*, vol. 10, no. 1, pp. 81–88, 2005.

[2] F. P. Vasilev, *Numerical Methods for Solving Extremal Problems*, Nauka, Moscow, Russia, 2nd edition, 1988.
[3] X. Weng, "Fixed point iteration for local strictly pseudo-contractive mapping," *Proceedings of the American Mathematical Society*, vol. 113, no. 3, pp. 727–731, 1991.

Xue Zhiqun: Department of Mathematics and Physics, Shijiazhuang Railway Institute, Shijiazhuang 050043, China
Email address: xuezhiqun@126.com

Mathematical Problems in Engineering

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at http://www.hindawi.com/journals/mpe/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/ according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru