Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2007, Article ID 78706, 6 pages
doi:10.1155/2007/78706

Research Article

A Fixed Point Theorem Based on Miranda

Uwe Schäfer

Received 5 June 2007; Revised 17 August 2007; Accepted 1 October 2007

Recommended by Robert F. Brown

A new fixed point theorem is proved by using the theorem of Miranda.

Copyright © 2007 Uwe Schäfer. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In 1940, Miranda published the following theorem ([1]).

Theorem 1.1. *Let* $\Omega = \{x \in \mathbb{R}^n : |x_i| \le L,\ i = 1,\dots,n\}$ *and let* $f : \Omega \to \mathbb{R}^n$ *be continuous satisfying*

$$\begin{aligned} f_i(x_1, x_2, \dots, x_{i-1}, -L, x_{i+1}, \dots, x_n) &\ge 0, \\ f_i(x_1, x_2, \dots, x_{i-1}, +L, x_{i+1}, \dots, x_n) &\le 0, \end{aligned} \quad \forall\, i\{1,\dots,n\}. \tag{1.1}$$

Then, $f(x) = 0$ *has a solution in* Ω.

For $n = 1$, Theorem 1.1 reduces to the well-known intermediate-value theorem. Miranda proved his theorem using the Brouwer fixed point theorem. Using the Brouwer degree of a mapping, Vrahatis gave another short proof of Theorem 1.1 (see [2]). Following this proof it is easy to see that Theorem 1.1 is also true, if L is dependent of i; that is, Ω can also be a rectangle and need not to be a cube. Even some L_i can be zero. Very often, the theorem of Miranda is stated as in the following corollary (see also [3, 4]), which is not the theorem of Miranda in its original form, but a consequence of it.

Corollary 1.2. *Let* $\hat{x} \in \mathbb{R}^n$, $L = (l_i) \in \mathbb{R}^n$, $l_i \ge 0$, *for* $i = 1,\dots,n$, *let* Ω *be the rectangle* $\Omega := \{x \in \mathbb{R}^n : |x_i - \hat{x}_i| \le l_i, i = 1,\dots,n\}$ *and let* $f : \Omega \to \mathbb{R}^n$ *be a continuous function on* Ω.

Also let

$$F_i^+ := \{x \in \Omega : x_i = \widehat{x}_i + l_i\}, \quad F_i^- := \{x \in \Omega : x_i = \widehat{x}_i - l_i\}, \quad i = 1,\dots,n, \tag{1.2}$$

be the pairs of parallel opposite faces of the rectangle Ω. *If for all* $i = 1,\dots,n$

$$f_i(x)\cdot f_i(y) \le 0, \quad \forall\, x \in F_i^+,\ \forall\, y \in F_i^-, \tag{1.3}$$

then there exists some $x^* \in \Omega$ *satisfying* $f(x^*) = 0$.

In principle, Corollary 1.2 says that Theorem 1.1 is also true if the $\le$ -sign and the $\ge$ -sign are exchanged with each other in (1.1). Corollary 1.2 also says that Theorem 1.1 is not restricted to a rectangle with 0 as its center.

Many generalizations have been given (see, e.g., [2, 4–6] for the finite-dimensional case and see [7, 8] for the infinite-dimensional case). In the presented paper we give a generalization of Corollary 1.2 in the infinite-dimensional Hilbert space l^2. Finally, we prove a fixed point version of Theorem 1.1 in l^2.

2. The infinite-dimensional case

Let l^2 be the infinite-dimensional Hilbert space of all square summable sequences of real numbers equipped with the natural order

$$x \le y :\Longleftrightarrow x_i \le y_i, \quad \forall_i \in \mathbb{N}, \tag{2.1}$$

and equipped with the norm $\|x\| := \sqrt{\sum_{i=1}^{\infty} x_i^2}$.

Theorem 2.1. *Let* $\widehat{x} = \{\widehat{x}_i\}_{i=1}^{\infty} \in l^2$, $L = \{l_i\}_{i=1}^{\infty} \in l^2$, $l_i \ge 0$, *for all* $i \in \mathbb{N}$, $\Omega := \{x \in l^2 : |x_i - \widehat{x}_i| \le l_i, f \text{ for all } i \in \mathbb{N}\}$ *and let* $f : \Omega \to l^2$ *be a continuous function on* Ω. *Also let*

$$F_i^+ := \{x \in \Omega : x_i = \widehat{x}_i + l_i\}, \quad F_i^- := \{x \in \Omega : x_i = \widehat{x}_i - l_i\}, \quad \forall\, i \in \mathbb{N}. \tag{2.2}$$

If for all $i \in \mathbb{N}$ *it holds that*

$$f_i(x)\cdot f_i(y) \le 0, \quad \forall\, x \in F_i^+,\ \forall\, y \in F_i^-, \tag{2.3}$$

then there exists some $x^* \in \Omega$ *satisfying* $f(x^*) = 0$.

Proof. For fixed $n \in \mathbb{N}$, we consider the function $\widetilde{h}^{(n)} : \Omega \to l^2$ defined by

$$\widetilde{h}^{(n)}(x) := \begin{pmatrix} f_1(x_1, x_2, \dots, x_{n-1}, x_n, x_{n+1}, \dots) \\ \vdots \\ f_n(x_1, x_2, \dots, x_{n-1}, x_n, x_{n+1}, \dots) \\ 0 \\ \vdots \end{pmatrix}. \tag{2.4}$$

Since Ω is compact and since f is continuous, the set $f(\Omega)$ is compact. Therefore, for given $\varepsilon > 0$ there is a finite set of elements $v^{(1)},\dots,v^{(p)} \in f(\Omega)$ such that if $f(x) \in f(\Omega)$,

then there is a $v \in \{v^{(1)},\dots,v^{(p)}\}$ such that

$$\|f(x)-v\| \le \varepsilon \tag{2.5}$$

and there exists $n_1 = n_1(\varepsilon) \in \mathbb{N}$ such that for all $n > n_1$ it holds that

$$\sqrt{\sum_{j=n+1}^{\infty} (v_j)^2} \le \varepsilon, \quad \forall v \in \{v^{(1)},\dots,v^{(p)}\}. \tag{2.6}$$

So, if $n > n_1$ is valid, then for all $f(x) \in f(\Omega)$ we have some $v \in \{v^{(1)},\dots,v^{(p)}\}$ such that

$$\left\|f(x)-\widetilde{h}^{(n)}(x)\right\| = \left\| \begin{pmatrix} 0 \\ \vdots \\ 0 \\ f_{n+1}(x) \\ f_{n+2}(x) \\ \vdots \end{pmatrix} \right\| \le \|f(x)-v\| + \left\| \begin{pmatrix} 0 \\ \vdots \\ 0 \\ v_{n+1} \\ v_{n+2} \\ \vdots \end{pmatrix} \right\| \le 2\varepsilon \tag{2.7}$$

for all $x \in \Omega$. Now, for fixed $n \in \mathbb{N}$ we define

$$\Omega_n := \begin{pmatrix} [\hat{x}_1 - l_1, \hat{x}_1 + l_1] \\ \vdots \\ [\hat{x}_n - l_n, \hat{x}_n + l_n] \end{pmatrix} \tag{2.8}$$

and $h^{(n)} : \Omega_n \to \mathbb{R}^n$ by

$$h^{(n)}(x) := \begin{pmatrix} f_1(x_1, x_2, \dots, x_{n-1}, x_n, \hat{x}_{n+1}, \hat{x}_{n+2}, \dots) \\ \vdots \\ f_n(x_1, x_2, \dots, x_{n-1}, x_n, \hat{x}_{n+1}, \hat{x}_{n+2}, \dots) \end{pmatrix}. \tag{2.9}$$

Due to (2.3) and Corollary 1.2 there exists $x^{(n)} \in \Omega_n$ with

$$h^{(n)}(x^{(n)}) = 0. \tag{2.10}$$

Setting

$$\widetilde{x}^{(n)} := \begin{pmatrix} x^{(n)} \\ \hat{x}_{n+1} \\ \hat{x}_{n+2} \\ \vdots \end{pmatrix}, \tag{2.11}$$

it holds that

$$\widetilde{x}^{(n)} \in \Omega, \quad \widetilde{h}^{(n)}(\widetilde{x}^{(n)}) = 0. \tag{2.12}$$

Now, let $n > n_1$. Then,

$$||f(\widetilde{x}^{(n)})|| = ||f(\widetilde{x}^{(n)}) - \widetilde{h}^{(n)}(\widetilde{x}^{(n)})|| \leq 2\varepsilon. \tag{2.13}$$

Hence, $\lim_{n\to\infty} f(\widetilde{x}^{(n)}) = 0$. Since Ω is compact, the sequence $\widetilde{x}^{(n)}$ has an accumulation point in Ω, say x^*. Without loss of generality, we assume that $\lim_{n\to\infty}\widetilde{x}^{(n)} = x^*$ holds. On the one hand, it follows that $\lim_{n\to\infty} f(\widetilde{x}^{(n)}) = f(x^*)$, since f is continuous. On the other hand, it follows that $f(x^*) = 0$, since the limit is unique. □

Next, we prove the fixed point version of Theorem 1.1 in l^2.

Theorem 2.2. *Let $L = \{l_i\}_{i=1}^{\infty} \in l^2$, $l_i \geq 0$, for all $i \in \mathbb{N}$. Let $\Omega = \{x \in l^2 : |x_i| \leq l_i, \forall i \in \mathbb{N}\}$ and suppose that the mapping $g : \Omega \to l^2$ is continuous satisfying*

$$\begin{aligned} &g_i(x_1, x_2, \ldots, x_{i-1}, -l_i, x_{i+1}, \ldots) \geq 0, \\ &g_i(x_1, x_{2,} \ldots, x_{i-1}, +l_i, x_{i+1}, \ldots) \leq 0, \end{aligned} \quad \forall\, i \in \mathbb{N}. \tag{2.14}$$

Then, $g(x) = x$ has a solution in Ω.

Proof. We consider the continuous function

$$f(x) := g(x) - x, \quad x \in \Omega. \tag{2.15}$$

Since for all $i \in \mathbb{N}$

$$\begin{aligned} f_i(x_1, \ldots, x_{i-1}, -l_i, x_{i+1}, \ldots) &= g_i(x_1, \ldots, x_{i-1}, -l_i, x_{i+1}, \ldots) + l_i \geq 0, \\ f_i(x_1, \ldots, x_{i-1}, +l_i, x_{i+1}, \ldots) &= g_i(x_1, \ldots, x_{i-1}, +l_i, x_{i+1}, \ldots) - l_i \leq 0, \end{aligned} \tag{2.16}$$

due to Theorem 2.1 there exists $x \in \Omega$ satisfying $f(x) = 0$; that is, $g(x) = x$. □

Example 2.3. Let $b \in l^2$ and $A = (a_{ik})$ satisfying $\sum_{i,k=1}^{\infty} |a_{ik}|^2 < \infty$. Then, the mapping

$$g(x) := \left(b_1 - \sum_{k=1}^{\infty} a_{1k}x_k, b_2 - \sum_{k=1}^{\infty} a_{2k}x_k, \ldots\right) \tag{2.17}$$

is (even) a compact mapping from l^2 to l^2. Now, if A is some kind of diagonally dominant in the sense that there exists some $L = \{l_i\}_{i=1}^{\infty} \in l^2$such that for all $i \in \mathbb{N}$

$$a_{ii} \cdot l_i \geq |b_i| + \sum_{k=1, k\neq i}^{\infty} |a_{ik}| \cdot l_k, \tag{2.18}$$

then by Theorem 2.1 there exists some $\xi \in \Omega = \{x \in l^2 : |x_i| \leq l_i, \forall i \in \mathbb{N}\}$ with $A\xi = b$. By Theorem 2.2 it follows that there exists $\eta \in \Omega$ satisfying $\eta = b - A\eta$.

Remark 2.4. Note that in Theorem 2.2 it is not necessary that g is a self-mapping as it is assumed in many other fixed point theorems.

Remark 2.5. Theorem 2.2 is also valid in $\mathbb{R}^n$ of course. Note, however, that the conditions (2.14) cannot be changed analogously as the conditions (1.1) have been changed to (1.3). We demonstrate this in Figure 2.1 for $n = 1$.




FIGURE 2.1. In both pictures the thick line is the graph of a function $y = g(x)$, $x \in \Omega$. In the left picture, $\Omega = [-L, L]$ and $g(-L) < 0$, $g(L) > 0$. According to Corollary 1.2 $g(x)$ has a zero in Ω. However, $g(x)$ has no fixed point in Ω, which is no contradiction to Theorem (2.2), since $g(-L) \geq 0$, $g(L) \leq 0$ is not valid, here. In the right picture, $\Omega = [\hat{x} - L, \hat{x} + L]$ and $g(\hat{x} - L) > 0$, $g(\hat{x} + L) < 0$. According to Corollary 1.2, $g(x)$ has a zero in Ω. However, $g(x)$ has no fixed point in Ω.

Acknowledgments

The author would like to thank the anonymous referee(s) for many suggestions and comments that helped to improve the paper. Furthermore, he would like to thank Professor Mitsuhiro Nakao for his invitation to the Kyushu University in Fukuoka, where this work was started.

References

[1] C. Miranda, "Un'osservazione su un teorema di Brouwer," *Bollettino dell'Unione Matematica Italiana*, vol. 3, pp. 5–7, 1940.
[2] M. N. Vrahatis, "A short proof and a generalization of Miranda's existence theorem," *Proceedings of the American Mathematical Society*, vol. 107, no. 3, pp. 701–703, 1989.
[3] J. B. Kioustelidis, "Algorithmic error estimation for approximate solutions of nonlinear systems of equations," *Computing*, vol. 19, no. 4, pp. 313–320, 1978.
[4] J. Mayer, "A generalized theorem of Miranda and the theorem of Newton-Kantorovich," *Numerical Functional Analysis and Optimization*, vol. 23, no. 3-4, pp. 333–357, 2002.
[5] G. Alefeld, A. Frommer, G. Heindl, and J. Mayer, "On the existence theorems of Kantorovich, Miranda and Borsuk," *Electronic Transactions on Numerical Analysis*, vol. 17, pp. 102–111, 2004.
[6] N. H. Pavel, "Theorems of Brouwer and Miranda in terms of Bouligand-Nagumo fields," *Analele Stiintifice ale Universitatii Al. I. Cuza din Iasi. Serie Noua. Matematica*, vol. 37, no. 2, pp. 161–164, 1991.
[7] C. Avramescu, "A generalization of Miranda's theorem," *Seminar on Fixed Point Theory Cluj-Napoca*, vol. 3, pp. 121–127, 2002.
[8] C. Avramescu, "Some remarks about Miranda's theorem," *Analele Universitatii din Craiova. Seria Matematica Informatica*, vol. 27, pp. 6–13, 2000.

Uwe Schäfer: Institut für Angewandte und Numerische Mathematik, Fakultät für Mathematik, Universität Karlsruhe (TH), D-76128 Karlsruhe, Germany
Email address: Uwe.Schaefer@math.uni-karlsruhe.de

Journal of Applied Mathematics and Decision Sciences

Special Issue on
Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods**: artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning
- **Application fields**: asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects**: decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site http://www.hindawi.com/journals/jamds/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk