Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2007, Article ID 87471, 5 pages
doi:10.1155/2007/87471

Research Article

On Fuzzy ε-Contractive Mappings in Fuzzy Metric Spaces

Dorel Miheţ

Received 24 December 2006; Accepted 1 March 2007

Recommended by Donal O'Regan

We answer into affirmative an open question raised by A. Razani in 2005. An essential role in our proofs is played by the separation axiom in the definition of a fuzzy metric space in the sense of George and Veeramani.

Copyright © 2007 Dorel Miheţ. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Preliminaries

In this section, we recall some definitions and results that will be used in the sequel.

Definition 1.1 (see [1]). A triple $(X, M, *)$, where X is an arbitrary set, $*$ is a continuous t-norm, and M is a fuzzy set on $X^2 \times (0, \infty)$, is said to be a *fuzzy metric space* (in the sense of George and Veeramani) if the following conditions are satisfied for all $x, y \in X$ and $s, t > 0$:
(GV-1) $M(x, y, t) > 0$;
(GV-2) $M(x, y, t) = 1$ if and only if $x = y$;
(GV-3) $M(x, y, t) = M(y, x, t)$;
(GV-4) $M(x, y, \cdot)$ is continuous;
(GV-5) $M(x, z, t + s) \geq M(x, y, t) * M(y, z, s)$.
Note (see [2]) that the "separation" condition (GV-2) means that

$$\begin{aligned} &M(x,x,t) = 1 \quad \forall x \in X,\ \forall t > 0, \\ &x \neq y \Longrightarrow M(x,y,t) < 1 \quad \forall t > 0. \end{aligned} \tag{1.1}$$

Definition 1.2 (see [1]). Let $(X, M, *)$ be a fuzzy metric space. A sequence $(x_n)_{n \in \mathbb{N}}$ in X is said to be *convergent* if there is $x \in X$ such that $\lim_{n \to \infty} M(x_n, x, t) = 1$ for each $t > 0$

(the notation $\lim_{n\to\infty} x_n = x$ or $x_n \to x$ will be used). A mapping $f : X \to X$ is said to be *continuous* if $f(x_n) \to f(x)$ whenever (x_n) is a sequence in X convergent to x.

Definition 1.3 (see [3]). Let $(X, M, *)$ be a fuzzy metric space and $0 < \varepsilon < 1$. A mapping $f : X \to X$ is called *fuzzy ε-contractive* if $M(f(x), f(y), t) > M(x, y, t)$ whenever $1 - \varepsilon < M(x, y, t) < 1$.

The next continuity lemma can be found in [4] (also see [5, Theorem 12.2.3]).

Lemma 1.4. *Let $(X, M, *)$ be a fuzzy metric space. If $\lim_{n\to\infty} x_n = x$ and $\lim_{n\to\infty} y_n = y$, then $\lim_{n\to\infty} M(x_n, y_n, t) = M(x, y, t)$ for all $t > 0$.*

2. Main results

The following theorem has been proved by Razani in [3].

Theorem 2.1 (see [3, Theorem 3.3]). *Let $(X, M, *)$ be a fuzzy metric space, where the continuous t-norm is defined as $a * b = \min\{a, b\}$. Suppose f is a fuzzy ε-contractive self-mapping of X such that there exists a point $x \in X$ whose sequence of iterates $(f^n(x))$ contains a convergent subsequence $(f^{n_i}(x))$. Then $\xi = \lim_{i\to\infty} f^{n_i}(x)$ is a periodic point, that is, there is a positive integer k such that $f^k(\xi) = \xi$.*

In [3, Question 3.7], it has been asked whether Theorem 2.1 would remain true if $*$ is replaced by an arbitrary t-norm.

With Theorem 2.3, we answer into affirmative this question. In the proofs of our theorems, we need the following.

Lemma 2.2. *Every fuzzy ε-contractive mapping in a fuzzy metric space is continuous.*

Proof. The continuity of the fuzzy ε-contractive mapping f is an immediate consequence of the implication

$$M(x, y, t) > 1 - \varepsilon \Longrightarrow M(f(x), f(y), t) \geq M(x, y, t) \tag{2.1}$$

which can be proved as follows: if $M(x, y, t) < 1$, then $M(x, y, t) > 1 - \varepsilon$ implies $M(f(x), f(y), t) > M(x, y, t)$, while if $M(x, y, t) = 1$ then, due to (GV-2), we have $x = y$, hence $M(f(x), f(y), t) = M(x, y, t)$. □

Theorem 2.3. *Let $(X, M, *)$ be a fuzzy metric space. Then for every fuzzy ε-contractive mapping f on X with the property that there exists a point $x \in X$ whose sequence of iterates $(f^n(x))_{n\in\mathbb{N}}$ contains a convergent subsequence, the point $\xi = \lim_{i\to\infty} f^{n_i}(x)$ is a periodic point.*

Proof. Since $*$ is continuous, there is $\delta \in (0, \varepsilon)$ such that $(1 - \delta) * (1 - \delta) > 1 - \varepsilon$. Also, there is a positive integer N_1 such that $i \geq N_1$ implies $M(f^{n_i}(x), \xi, t/2) > 1 - \delta$, for all $t > 0$. Fix a $k \geq N_1$ and denote $n_{k+1} - n_k$ by s. As f is fuzzy ε-contractive and $M(f^{n_k}(x), \xi, t/2) > 1 - \varepsilon$, we have

$$M\left(f^{n_k+1}(x), f(\xi), \frac{t}{2}\right) \geq M\left(f^{n_k}(x), \xi, \frac{t}{2}\right) > 1 - \delta > 1 - \varepsilon \tag{2.2}$$

and, after $n_{k+1}-n_k$ iterations, $M(f^{n_{k+1}}(x),f^s(\xi),t/2)>1-\delta$. Therefore,

$$\begin{aligned} M(\xi,f^s(\xi),t) &\geq M\left(f^{n_{k+1}}(x),\xi,\frac{t}{2}\right)*M\left(f^{n_{k+1}}(x),f^s(\xi),\frac{t}{2}\right) \\ &\geq (1-\delta)*(1-\delta)>1-\varepsilon \quad \forall t>0. \end{aligned} \tag{2.3}$$

Since f is continuous, $\lim_{i\to\infty}f^{n_i}(x)=\xi$ implies $\lim_{i\to\infty}f^{n_i+s}(x)=f^s(\xi)$, therefore, by Lemma 1.4,

$$\lim_{i\to\infty}M\left(f^{n_i}(x),f^{n_i+s}(x),t\right)=M(\xi,f^s(\xi),t) \quad \forall t>0. \tag{2.4}$$

As the sequence of real numbers $(z_n)_{n\geq n_k}$, $z_n:=M(f^n(x),f^{n+s}(x),t)_{n\geq n_k}$ is convergent for every $t>0$ (being nondecreasing and bounded), one has

$$\lim_{n\to\infty}M\left(f^n(x),f^{n+s}(x),t\right)=M(\xi,f^s(\xi),t) \quad \forall t>0. \tag{2.5}$$

On the other hand, from $f^{n_i}(f(x))=f(f^{n_i}(x))\to_{i\to\infty}f(\xi)$ and $f^{n_i}(f^{s+1}(x))=f^{s+1}(f^{n_i}(x))\to f^{s+1}(\xi)$, it follows that

$$\lim_{i\to\infty}M\left(f^{n_i+1}(x),f^{n_i+1+s}(x),t\right)=M\left(f(\xi),f^{s+1}(\xi),t\right) \quad \forall t>0, \tag{2.6}$$

that is,

$$M(\xi,f^s(\xi),t)=M\left(f(\xi),f^{s+1}(\xi),t\right) \quad \forall t>0. \tag{2.7}$$

We claim that $f^s(\xi)=\xi$. Indeed, if $f^s(\xi)\neq\xi$ then, due to (GV-2), $M(\xi,f^s(\xi),t)<1$, for all $t>0$ and since $M(\xi,f^s(\xi),t)>1-\varepsilon$ for all $t>0$, we have $1-\varepsilon<M(\xi,f^s(\xi),t)<1$ for all $t>0$. This implies $M(\xi,f^s(\xi),t)<M(f(\xi),f^{s+1}(\xi),t)$ for all $t>0$, which is a contradiction. Therefore, $f^s(\xi)=\xi$, concluding the proof. □

Example 2.4. Consider fuzzy metric space $(N^*,M,*)$, where $N^*=\{1,2,\dots\}$, $a*b=\min\{a,b\}$, and

$$M(x,y,t)=\begin{cases}\dfrac{1}{2}, & x\neq y,\\ 1, & x=y,\end{cases} \tag{2.8}$$

for all $t>0$. The mapping $f:N^*\to N^*$,

$$f(x)=\begin{cases}1 & \text{if } x \text{ is even},\\ 2 & \text{if } x \text{ is odd},\end{cases} \tag{2.9}$$

is fuzzy 1/2-contractive, in the absence of the condition $1-\varepsilon<M(x,y,t)<1$. The sequence of the successive approximations of 1 is $2,1,2,1,2,1,\dots$, and its subsequence $1,1,\dots$ converges to 1, which is a periodic point for f.

In the following, we show that the assertion [3, Corollary 3.5] claiming that in the conditions of Theorem 2.1 we cannot have $M(\xi,f(\xi),t)>1-\varepsilon$, is not correct. As a matter

of fact, we will show in Theorem 2.7 that $M(\xi, f(\xi), t) > 1 - \varepsilon$ is a sufficient condition for the existence of a fixed point for a fuzzy ε-contractive mapping.

Example 2.5. Consider the standard fuzzy metric space $(X, M, *)$, where $X = (-\infty, \infty)$, $M(x, y, t) = t/(t + |x - y|)$, $a * b = \min\{a, b\}$, and the mapping $f : X \to X$, $f(x) = x/2$. Since

$$M(f(x), f(y), t) = \frac{2t}{2t + |x - y|} > \frac{t}{t + |x - y|} = M(x, y, t) \tag{2.10}$$

for all $x, y \in X$, $x \neq y$, and $t > 0$, f is fuzzy ε-contractive for every $\varepsilon \in (0, 1)$ and it is immediate that the sequence of iterates of any point converges to 0. As 0 is a fixed point of f, we have $M(0, f(0), t) = 1 > 1 - \varepsilon$ for every $\varepsilon \in (0, 1)$.

The error in the proof of the corollary derives from the fact that the (strict) inequality $M(f^2(\xi), f(\xi), t) > M(f(\xi), \xi, t)$ (see [3]) takes place only if $M(f(\xi), \xi, t) \neq 1$, that is, (due to (GV-2)) only if $f(\xi) \neq \xi$. The next proposition is a correct version of [3, Corollary 3.5].

Proposition 2.6. *Let $(X, M, *)$ be a fuzzy metric space and let $f : X \to X$ be a fuzzy ε-contractive mapping. Suppose that there is $\zeta \in X$ such that $M(\xi, f(\xi), t) > 1 - \varepsilon$ for some $t > 0$ and $f^k(\xi) = \xi$ for some integer $k \geq 1$. Then $f(\xi) = \xi$.*

Proof. From $M(\xi, f(\xi), t) > 1 - \varepsilon$, it follows that

$$M(f^l(\xi), f^{l+1}(\xi), t) \geq M(f(\xi), f^2(\xi), t) \tag{2.11}$$

for all $l \geq 1$. Thus,

$$M(\xi, f(\xi), t) = M(f^k(\xi), f^{k+1}(\xi), t) \geq M(f(\xi), f^2(\xi), t). \tag{2.12}$$

If we had $f(\xi) \neq \xi$, then due to (GV-2), $M(\xi, f(\xi), t) \neq 1$. As $M(\xi, f(\xi), t) > 1 - \varepsilon$, from the definition of a ε-fuzzy contractive mapping, the strict inequality

$$M(f(\xi), f^2(\xi), t) > M(\xi, f(\xi), t) \tag{2.13}$$

would follow, and thus we would obtain

$$M(\xi, f(\xi), t) \geq M(f(\xi), f^2(\xi), t) > M(\xi, f(\xi), t). \tag{2.14}$$

This contradiction completes the proof. □

A sufficient condition for the existence of a fixed point for a fuzzy ε-contraction is given in the next theorem.

Theorem 2.7. *Let $(X, M, *)$ be a fuzzy metric space and let $f : X \to X$ be a fuzzy ε-contractive mapping. Suppose that for some $x \in X$, the sequence $(f^n(x))_{n \in \mathbb{N}}$ contains a convergent subsequence and let $\zeta \in X$ be its limit. If there exists $t_0 > 0$ such that $M(x, f(x), t_0) > 1 - \varepsilon$ and $M(\zeta, f(\zeta), t_0) > 1 - \varepsilon$, then ζ is a fixed point of f.*

Proof. Let $x_n = f^n(x)$ and let $(x_{n_k})_{k \in \mathbb{N}}$ be a convergent subsequence of (x_n). As the sequence $(f(x_{n_k}))_{k \in \mathbb{N}}$ converges to $f(\zeta)$ and the sequence $(f(f(x_{n_k})))_{k \in \mathbb{N}}$ converges to

$f(f(\zeta))$ (see Lemma 2.2), we have (see Lemma 1.4)

$$\begin{aligned} M(x_{n_k}, f(x_{n_k}), t) &\longrightarrow M(\zeta, f(\zeta), t) \quad \forall t > 0, \\ M(f(x_{n_k}), f^2(x_{n_k}), t) &\longrightarrow M(f(\zeta), f^2(\zeta), t) \quad \forall t > 0. \end{aligned} \tag{2.15}$$

Since $M(x, f(x), t_0) > 1 - \varepsilon$, the sequence $(z_n)_{n\in\mathbb{N}}$, $z_n := M(x_n, f(x_n), t_0)$ is a nondecreasing sequence of numbers in $[0,1]$, therefore it is convergent. As its subsequence $(M(x_{n_k}, f(x_{n_k}), t_0))$ converges to $M(\zeta, f(\zeta), t_0)$, it follows that z_n converges to $M(\zeta, f(\zeta), t_0)$. Also,

$$\lim_{n\to\infty} z_{n+1} = \lim_{n\to\infty} M(f(x_n), f^2(x_n), t_0) = M(f(\zeta), f^2(\zeta), t_0), \tag{2.16}$$

therefore the equality $M(\zeta, f(\zeta), t_0) = M(f(\zeta), f^2(\zeta), t_0)$ holds.

Suppose $\zeta \neq f(\zeta)$. Then, due to (GV-2), $M(\zeta, f(\zeta), t_0)$ is not 1, hence $1 - \varepsilon < M(\zeta, f(\zeta), t_0) < 1$. This implies that $M(f(\zeta), f^2(\zeta), t_0) > M(\zeta, f(\zeta), t_0)$, contradicting the above equality. Therefore, ζ is a fixed point of f. □

Example 2.8. Let $X = (0, \infty)$, $M(x, y, t) = \min\{x, y\}/\max\{x, y\}$ for all $t > 0$ and $a * b = ab$. Then (see [6]), $(X, M, *)$ is a fuzzy metric space. Since $\sqrt{t} > t$ for all $t \in (0,1)$, the mapping $f : X \to X$, $f(x) = \sqrt{x}$, is fuzzy ε-contractive for every $\varepsilon \in (0,1)$ and the sequence $(f^n(1))_{n\in\mathbb{N}}$ is convergent to 1, the fixed point of f. Note that the condition $M(1, f(1), t) > 1 - \varepsilon$ is not satisfied by the mapping in Example 2.4.

Remark 2.9. Theorem 2.3 is Theorem 2.4 in our archived manuscript 35106, submitted in the 4th of August 2005 to FPTA. Recently, Ćirić et al. [7] solved a similar question of Razani for mappings in intuitionistic fuzzy metric spaces.

References

[1] A. George and P. Veeramani, "On some results in fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 64, no. 3, pp. 395–399, 1994.

[2] V. Gregori and S. Romaguera, "Characterizing completable fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 144, no. 3, pp. 411–420, 2004.

[3] A. Razani, "A contraction theorem in fuzzy metric spaces," *Fixed Point Theory and Applications*, vol. 2005, no. 3, pp. 257–265, 2005.

[4] M. Grabiec, "Fixed points in fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 27, no. 3, pp. 385–389, 1988.

[5] B. Schweizer and A. Sklar, *Probabilistic Metric Spaces*, North-Holland Series in Probability and Applied Mathematics, North-Holland, New York, NY, USA, 1983.

[6] V. Radu, "Some suitable metrics on fuzzy metric spaces," *Fixed Point Theory*, vol. 5, no. 2, pp. 323–347, 2004.

[7] L. Ćirić, S. Ješić, and J. S. Ume, "The existence theorems for fixed and periodic points of nonexpansive mappings in intuitionistic fuzzy metric spaces," to appear in *Chaos, Solitons & Fractals*.

Dorel Miheţ: Faculty of Mathematics and Computer Science, West University of Timişoara, Bv. V. Parvan 4, 300223 Timişoara, Romania

Email addresses: doru_mihet@yahoo.com; mihet@math.uvt.ro

Journal of Applied Mathematics and Decision Sciences

Special Issue on
Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods**: artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning
- **Application fields**: asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects**: decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site http://www.hindawi.com/journals/jamds/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk