Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2007, Article ID 96941, 6 pages
doi:10.1155/2007/96941

Research Article

Existence Principle for Advanced Integral Equations on Semiline

Adela Chiş

Received 18 May 2007; Accepted 16 July 2007

Recommended by Andrzej Szulkin

The continuation principle for generalized contractions in gauge spaces is used to discuss nonlinear integral equations with advanced argument.

Copyright © 2007 Adela Chiş. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

This paper deals with an integral equation with advanced argument. The advanced argument makes necessary the use of two pseudometrics in the contraction condition. For this reason we will apply the continuation principle established in Chiş and Precup [1] involving contractions in Gheorghiu's sense, with respect to a family of pseudometrics rather than the existence principle from Frigon [2, 3].

In what follows we recall some notions and results from papers Chiş and Precup [1] and Chiş [4].

First recall the notion of a contraction on a gauge space introduced by Gheorghiu [5].

Definition 1.1 (Gheorghiu [5]). Let $(X,\mathcal{P})$ be a gauge space with the family of pseudometrics $\mathcal{P} = \{p_\alpha\}_{\alpha\in A}$, where A is a set of indices. A map $F : D \subset X \to X$ is a *contraction* if there exists a function $\varphi : A \to A$ and $a \in \mathbb{R}_+^A, a = \{a_\alpha\}_{\alpha\in A}$ such that

$$\begin{aligned} &p_\alpha(F(x),F(y)) \le a_\alpha p_{\varphi(\alpha)}(x,y), \quad \forall \alpha \in A,\ x,y \in D, \\ &\sum_{n=1}^{\infty} a_\alpha a_{\varphi(\alpha)} a_{\varphi^2(\alpha)} \cdots a_{\varphi^{n-1}(\alpha)} p_{\varphi^n(\alpha)}(x,y) < \infty, \end{aligned} \tag{1.1}$$

for every $\alpha \in A$ and $x, y \in D$. Here, φ^n is the nth iteration of φ.

Theorem 1.2 (Chiş [4]). *Let X be a set endowed with two separating gauge structures: $\mathcal{P} = \{p_\alpha\}_{\alpha\in A}$ and $\mathcal{Q} = \{q_\beta\}_{\beta\in B}$, let D_0 and D be two subsets of X with $D_0 \subset D$, and let $F : D \to X$ be a map. Assume that $F(D_0) \subset D_0$ and D is $\mathcal{P}$-closed. In addition, assume that the following conditions are satisfied:*

(i) *there is a function $\psi : A \to B$ and $c \in (0,\infty)^A$, $c = \{c_\alpha\}_{\alpha\in A}$ such that*

$$p_\alpha(x,y) \le c_\alpha q_{\psi(\alpha)}(x,y), \quad \forall \alpha \in A,\ x,y \in X; \tag{1.2}$$

(ii) *$(X,\mathcal{P})$ is a sequentially complete gauge space;*

(iii) *if $x_0 \in D$, $x_n = F(x_{n-1})$, for $n = 1,2,\ldots$, and $\mathcal{P} - \lim_{n\to\infty} x_n = x$ for some $x \in D$, then $F(x) = x$;*

(iv) *F is a $\mathcal{Q}$-contraction on D_0.*

Then F has at least one fixed point which can be obtained by successive approximations starting from any element of D_0.

For a map $H : D \times [0,1] \to X$, where $D \subset X$, we will use the following notations:

$$\begin{aligned} \Sigma &= \{(x,\lambda) \in D \times [0,1] : H(x,\lambda) = x\}, \\ S &= \{x \in D : H(x,\lambda) = x, \text{ for some } \lambda \in [0,1]\}, \\ \Lambda &= \{\lambda \in [0,1] : H(x,\lambda) = x, \text{ for some } x \in D\}. \end{aligned} \tag{1.3}$$

Theorem 1.3 (Chiş and Precup [1]). *Let X be a set endowed with the separating gauge structures $\mathcal{P} = \{p_\alpha\}_{\alpha\in A}$ and $\mathcal{Q}^\lambda = \{q^\lambda_\beta\}_{\beta\in B}$, for $\lambda \in [0,1]$. Let $D \subset X$ be $\mathcal{P}$-sequentially closed, $H : D \times [0,1] \to X$ a map, and assume that the following conditions are satisfied:*

(i) *for each $\lambda \in [0,1]$, there exists a function $\varphi_\lambda : B \to B$ and $a^\lambda \in [0,1)^B$, $a^\lambda = \{a^\lambda_\beta\}_{\beta\in B}$ such that*

$$\begin{aligned} &q^\lambda_\beta(H(x,\lambda),H(y,\lambda)) \le a^\lambda_\beta q^\lambda_{\varphi_\lambda(\beta)}(x,y), \\ &\sum_{n=1}^{\infty} a^\lambda_\beta a^\lambda_{\varphi_\lambda(\beta)} a^\lambda_{\varphi^2_\lambda(\beta)} \cdots a^\lambda_{\varphi^{n-1}_\lambda(\beta)} < \infty, \end{aligned} \tag{1.4}$$

for every $\beta \in B$ and $x,y \in D$;

(ii) *there exists $\rho > 0$ such that for each $(x,\lambda) \in \Sigma$, there is a $\beta \in B$ with*

$$\inf\{q^\lambda_\beta(x,y) : y \in X \setminus D\} > \rho; \tag{1.5}$$

(iii) *for each $\lambda \in [0,1]$, there is a function $\psi : A \to B$ and $c \in (0,\infty)^A$, $c = \{c_\alpha\}_{\alpha\in A}$ such that*

$$p_\alpha(x,y) \le c_\alpha q^\lambda_{\psi(\alpha)}(x,y), \quad \forall \alpha \in A,\ x,y \in X; \tag{1.6}$$

(iv) *$(X,\mathcal{P})$ is a sequentially complete gauge space;*

(v) *if $\lambda \in [0,1]$, $x_0 \in D$, $x_n = H(x_{n-1},\lambda)$, for $n = 1,2,\ldots$, and $\mathcal{P} - \lim_{n\to\infty} x_n = x$, then $H(x,\lambda) = x$;*

(vi) *for every* $\varepsilon > 0$, *there exists* $\delta = \delta(\varepsilon) > 0$ *with*

$$q^{\lambda}_{\varphi^n_\lambda(\beta)}(x, H(x,\lambda)) \leq (1 - a^{\lambda}_{\varphi^n_\lambda(\beta)})\varepsilon, \tag{1.7}$$

for $(x,\mu) \in \Sigma$, $|\lambda - \mu| \leq \delta$, *all* $\beta \in B$, *and* $n \in \mathbb{N}$.

In addition, assume that $H_0 := H(\cdot, 0)$ *has a fixed point. Then, for each* $\lambda \in [0,1]$, *the map* $H_\lambda := H(\cdot,\lambda)$ *has at least a fixed point.*

2. The main result

We consider the integral equation inspired from biomathematics (see O'Regan and Precup [6]):

$$x(t) = \int_{t-1}^{t} f(s, x(s+2)), \quad t \in [0,\infty). \tag{2.1}$$

Let $I = [-1,\infty)$ and for a function $u \in L^1(a,b)$ we denote by $|u|_{L^1(a,b)}$ the norm in $L^1(a,b)$. We have the following existence principle for (2.1).

Theorem 2.1. *Let* $(E, \|\cdot\|)$ *be a Banach space, and let* $f : I \times E \to E$ *be a continuous function. Assume that the following conditions hold:*

(a) *there exists* $k : I \to (0,\infty)$, $k \in L^1_{\text{loc}}(I)$ *with* $|k|_{L^1_{\text{loc}}(I)} < 1$ *such that*

$$\|f(t,x) - f(t,y)\| \leq k(t)|x - y| \tag{2.2}$$

for all $x, y \in E$, *and* $t \in I$;

(b) *for each* $n \in \mathbb{N}$ *there exists* $r_n > 0$ *such that, any continuous solution* x *of the equation*

$$x(t) = \lambda\left(\int_{t-1}^{t} f(s, x(s+1))ds\right), \quad t \in [0,\infty) \tag{2.3}$$

with $\lambda \in [0,1]$, *satisfies* $\|x(t)\| \leq r_n$ *for any* $t \in [n, 2n+1]$;

(c) *there exists* $\alpha \in L^1_{\text{loc}}(I)$ *şi* $\beta : [0,\infty) \to (0,\infty)$ *nondecreasing such that*

$$\|f(t,x)\| \leq \alpha(t)\beta(\|x\|) \tag{2.4}$$

for all $t \in I$ *and* $x \in E$;

(d) *there exists* $C > 0$ *such that* $\beta(r_{k+1})/(1 - L_k) \leq C$ *for any* $k \in \mathbb{N}$, *where* $L_n = \int_{n-1}^{2n+1} k(s)ds$.

Then there exists at least one solution $x \in C(\mathbb{R}_+, E)$ *of the integral equation (2.1).*

Proof. For the proof we use Theorem 1.3. Let $X = C(R_+, E)$. For each $n \in \mathbb{N}$ we define the map $|\cdot|_n : X \to \mathbb{R}_+$ by $|x|_n = \max_{t\in[n,2n+1]} \|x(t)\|$. This map is a seminorm on X, and let $d_n : X \times X \to \mathbb{R}_+$ be given by

$$d_n(x,y) = |x - y|_n = \max_{t\in[n,2n+1]} \|x(t) - y(t)\|. \tag{2.5}$$

It is easy to show that d_n is a pseudometric on X and the family $\{d_n\}_{n\in\mathbb{N}}$ defines on X a gauge structure, separated and complete by sequences.

Here $\mathcal{P} = \mathcal{Q}^{\lambda} = \{d_n\}_{n\in\mathbb{N}}$ for each $\lambda \in [0,1]$. Let D be the closure in X of the set

$$\{x \in X : \text{ there exists } n \in \mathbb{N} \text{ such that } d_n(x,0) \le r_n + \delta\}, \tag{2.6}$$

where $\delta > 0$ is a fixed number. We define $H : D \times [0,1] \to X$ by $H(x,\lambda) = \lambda A(x)$, where

$$A(x)(t) = \int_{t-1}^{t} f(s, x(s+2))\,ds. \tag{2.7}$$

First, we verify condition (i) from Theorem 1.3.

Let $t \in [n, 2n+1]$, where $n \ge 0$. We have

$$\begin{aligned} ||H(x,\lambda)(t) - H(y,\lambda)(t)|| &\le \lambda \int_{t-1}^{t} ||f(s,x(s+2)) - f(s,y(s+2))||\,ds \\ &\le \int_{n-1}^{2n+1} k(s)||x(s+2) - y(s+2)||\,ds \\ &\le \max_{s\in[n-1,2n+1]} ||x(s+2) - y(s+2)|| \int_{n-1}^{2n+1} k(s)\,ds \\ &\le \max_{\tau\in[n+1,2n+3]} ||x(\tau) - y(\tau)|| \int_{n-1}^{2n+1} k(s)\,ds \\ &= L_n d_{n+1}(x,y). \end{aligned} \tag{2.8}$$

If we take the maximum with respect to t, we obtain

$$d_n(H(x,\lambda), H(y,\lambda)) \le L_n d_{n+1}(x,y) \tag{2.9}$$

for all $x, y \in D$ and all $n \in \mathbb{N}$. Hence, condition (i) in Theorem 1.3 holds with $\varphi_\lambda = \varphi$ where $\varphi : \mathbb{N} \to \mathbb{N}$ is defined by $\varphi(n) = n+1$. In addition, the series $\sum_{n=1}^{\infty} L_n L_{n+1} \cdots L_{2n}$ is finite since from assumption (a) we know that $|k|_{L^1_{\mathrm{loc}}(I)} < 1$ so $L_n \le |k|_{L^1_{\mathrm{loc}}(I)} < 1$.

Condition (ii) in our case becomes: there exists $\rho > 0$ such that for any solution $(x,\lambda) \in D \times [0,1]$, to $x = H(x,\lambda)$, there exists $n \in \mathbb{N}$ with

$$\inf\{d_n(x,y) : y \in X \setminus D\} > \rho. \tag{2.10}$$

If $y \in X \setminus D$, we have that $d_n(y,0) > r_n + \delta$ for each $n \in \mathbb{N}$. So there exists at least one $t \in [n, 2n+1]$ with

$$||x(t) - y(t)|| \ge ||y(t)|| - ||x(t)|| > r_n + \delta - r_n = \delta. \tag{2.11}$$

Hence $d_n(x,y) > \delta$ and (2.10) holds for any $\rho \in (0,\delta)$.

Condition (iii) in Theorem 1.3 is trivial since $\mathcal{P} = \mathcal{Q}^{\lambda}$ for any $\lambda \in [0,1]$.

Condition (iv) in Theorem 1.3 becomes: $(X, \{d_n\}_{n\in\mathbb{N}})$ is a gauge space sequatially complete because E is a Banach space.

Condition (v): Let $\lambda \in [0,1]$, $x_0 \in D$, $x_n = H(x_{n-1},\lambda)$ for $n = 1,2,\ldots$, and assume $\mathcal{P} - \lim_{n\to\infty} x_n = x$. We will prove that $H(x,\lambda) = x$.

Let $m \in \mathbb{N}$ and $t \in [m, 2m+1]$. We have

$$\begin{aligned}
||H(x,\lambda)(t) - x(t)|| &= ||H(x,\lambda)(t) - x_n(t) + x_n(t) - x(t)|| \\
&\leq ||H(x,\lambda)(t) - x_n(t)|| + ||x_n(t) - x(t)|| \\
&= ||H(x,\lambda)(t) - H(x_{n-1},\lambda)(t)|| + ||x_n(t) - x(t)|| \\
&\leq \int_{t-1}^{t} k(s)||x(s+2) - x_{n-1}(s+2)||ds + \max_{t\in[m,2m+1]} ||x_n(t) - x(t)|| \\
&\leq L_m \max_{s\in[m-1,2m+1]} ||x(s+2) - x_{n-1}(s+2)|| + d_m(x_n, x) \\
&= L_m \max_{\tau\in[m+1,2m+3]} ||x(\tau) - x_{n-1}(\tau)|| + d_m(x_n, x) \\
&= L_m d_{m+1}(x, x_{n-1}) + d_m(x_n, x).
\end{aligned} \tag{2.12}$$

Consequently, passing to maximum after $t \in [m, 2m+1]$ we have

$$d_m(H(x,\lambda), x) \leq L_m d_{m+1}(x, x_{n-1}) + d_m(x_n, x) \tag{2.13}$$

for all $m \in \mathbb{N}$. Letting $n \to \infty$, we deduce that $d_m(H(x,\lambda), x) = 0$ for each $m \in \mathbb{N}$ and since the family $\{d_m\}_{m\in\mathbb{N}}$ is separated, we have $H(x,\lambda) = x$.

Condition (vi) becomes: for each $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that

$$d_{\varphi^n(m)}(x, H(x,\lambda)) \leq (1 - L_{\varphi^n(m)})\varepsilon \tag{2.14}$$

for each $(x,\mu) \in D \times [0,1]$, $H(x,\mu) = x$, $|\lambda - \mu| \leq \delta$, and $n, m \in \mathbb{N}$.

We have $\varphi^n(m) = n + m$. Let $t \in [n+m, 2(n+m)+1]$, and using conditions (c) and (d) we obtain

$$\begin{aligned}
||x(t) - H(x,\lambda)(t)|| &= ||H(x,\mu)(t) - H(x,\lambda)(t)|| \\
&= |\mu - \lambda| \left|\left| \int_{t-1}^{t} f(s, x(s+2))ds \right|\right| \\
&\leq |\mu - \lambda| \int_{t-1}^{t} \alpha(s)\beta(||x(s+2)||)ds \\
&\leq |\mu - \lambda|\beta(r_{m+n+1}) \int_{n+m-1}^{2(n+m)+1} \alpha(s)ds \\
&\leq |\mu - \lambda||\alpha|_{L^1_{\mathrm{loc}}(I)} C(1 - L_{m+n}).
\end{aligned} \tag{2.15}$$

So condition (vi) is true with $\delta(\varepsilon) = \varepsilon / C|\alpha|_{L^1_{\mathrm{loc}}(I)}$.

In addition, $H(\cdot, 0) = 0$. So $H(\cdot, 0)$ has a fixed point.

Therefore, all the assumptions of Theorem 1.3 are satisfied. Now the conclusion follows from Theorem 1.3. □

Other existence results for integral and differential equations established by the continuation method (see O'Regan and Precup [6]) are given in Chiş [4, 7].

References

[1] A. Chiş and R. Precup, "Continuation theory for general contractions in gauge spaces," *Fixed Point Theory and Applications*, vol. 2004, no. 3, pp. 173–185, 2004.

[2] M. Frigon, "Fixed point results for generalized contractions in gauge spaces and applications," *Proceedings of the American Mathematical Society*, vol. 128, no. 10, pp. 2957–2965, 2000.

[3] M. Frigon, "Fixed point results for multivalued contractions on gauge spaces," in *Set Valued Mappings with Applications in Nonlinear Analysis*, R. P. Agarwal and D. O'Regan, Eds., vol. 4 of *Series in Mathematical Analysis and Applications*, pp. 175–181, Taylor & Francis, London, UK, 2002.

[4] A. Chiş, "Initial value problem on semi-line for differential equations with advanced argument," *Fixed Point Theory*, vol. 7, no. 1, pp. 37–42, 2006.

[5] N. Gheorghiu, "Contraction theorem in uniform spaces," *Studii şi Cercetări Matematice*, vol. 19, pp. 119–122, 1967 (Romanian).

[6] D. O'Regan and R. Precup, *Theorems of Leray-Schauder Type and Applications*, vol. 3 of *Series in Mathematical Analysis and Applications*, Gordon and Breach Science, Amsterdam, The Netherlands, 2001.

[7] A. Chiş, "Continuation methods for integral equations in locally convex spaces," *Studia Universitatis Babeş-Bolyai Mathematica*, vol. 50, no. 3, pp. 65–79, 2005.

Adela Chiş: Department of Mathematics, Technical University of Cluj-Napoca,
400020 Cluj-Napoca, Romania
Email address: adela.chis@math.utcluj.ro

Journal of Applied Mathematics and Decision Sciences

Special Issue on
Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods**: artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning
- **Application fields**: asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects**: decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site http://www.hindawi.com/journals/jamds/. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk