

Research Article

Convergence of Iterative Sequences for Fixed Point and Variational Inclusion Problems

Li Yu and Ma Liang

School of Management, University of Shanghai for Science and Technology, Shanghai 200093, China

Correspondence should be addressed to Li Yu, brucemath@139.com

Received 14 November 2010; Accepted 8 February 2011

Academic Editor: Yeol J. Cho

Copyright © 2011 L. Yu and M. Liang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

An iterative process is considered for finding a common element in the fixed point set of a strict pseudocontraction and in the zero set of a nonlinear mapping which is the sum of a maximal monotone operator and an inverse strongly monotone mapping. Strong convergence theorems of common elements are established in real Hilbert spaces.

1. Introduction and Preliminaries

Throughout this paper, we always assume that H is a real Hilbert space with the inner product $\langle \cdot, \cdot \rangle$ and the norm $\| \cdot \|$.

Let C be a nonempty closed convex subset of H and $S : C \rightarrow C$ a nonlinear mapping. In this paper, we use $F(S)$ to denote the fixed point set of S . Recall that the mapping S is said to be nonexpansive if

$$\|Sx - Sy\| \leq \|x - y\|, \quad \forall x, y \in C. \quad (1.1)$$

S is said to be κ -strictly pseudocontractive if there exists a constant $\kappa \in [0, 1)$ such that

$$\|Sx - Sy\|^2 \leq \|x - y\|^2 + \kappa \|(x - Sx) - (y - Sy)\|^2, \quad \forall x, y \in C. \quad (1.2)$$

The class of strictly pseudocontractive mappings was introduced by Browder and Petryshyn [1] in 1967. It is easy to see that every nonexpansive mapping is a 0-strictly pseudocontractive mapping.

Let $A : C \rightarrow H$ be a mapping. Recall that A is said to be monotone if

$$\langle Ax - Ay, x - y \rangle \geq 0, \quad \forall x, y \in C. \quad (1.3)$$

A is said to be inverse strongly monotone if there exists a constant $\alpha > 0$ such that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|Ax - Ay\|^2, \quad \forall x, y \in C. \quad (1.4)$$

For such a case, A is also said to be α -inverse strongly monotone.

Let $M : H \rightarrow 2^H$ be a set-valued mapping. The set $D(M)$ defined by $D(M) = \{x \in H : Mx \neq \emptyset\}$ is said to be the domain of M . The set $R(M)$ defined by $R(M) = \bigcup_{x \in H} Mx$ is said to be the range of M . The set $G(M)$ defined by $G(M) = \{(x, y) \in H \times H : x \in D(M), y \in R(M)\}$ is said to be the graph of M .

Recall that M is said to be monotone if

$$\langle x - y, f - g \rangle > 0, \quad \forall (x, f), (y, g) \in G(M). \quad (1.5)$$

M is said to be maximal monotone if it is not properly contained in any other monotone operator. Equivalently, M is maximal monotone if $R(I + rM) = H$ for all $r > 0$. For a maximal monotone operator M on H and $r > 0$, we may define the single-valued resolvent $J_r = (I + rM)^{-1} : H \rightarrow D(M)$. It is known that J_r is firmly nonexpansive and $M^{-1}(0) = F(J_r)$.

Recall that the classical variational inequality problem is to find $x \in C$ such that

$$\langle Ax, y - x \rangle \geq 0, \quad \forall y \in C. \quad (1.6)$$

Denote by $VI(C, A)$ of the solution set of (1.6). It is known that $x \in C$ is a solution to (1.6) if and only if x is a fixed point of the mapping $P_C(I - \lambda A)$, where $\lambda > 0$ is a constant and I is the identity mapping.

Recently, many authors considered the convergence of iterative sequences for the variational inequality (1.6) and fixed point problems of nonlinear mappings see, for example, [1–32].

In 2005, Iiduka and Takahashi [7] proved the following theorem.

Theorem IT. *Let C be a closed convex subset of a real Hilbert space H . Let A be an α -inverse-strongly monotone mapping of C into H , and let S be a nonexpansive mapping of C into itself such that $F(S) \cap VI(C, A) \neq \emptyset$. Suppose that $x_1 = x \in C$ and $\{x_n\}$ is given by*

$$x_{n+1} = \alpha_n x + (1 - \alpha_n) SP_C(x_n - \lambda_n Ax_n), \quad (1.7)$$

for every $n = 1, 2, \dots$, where $\{\alpha_n\}$ is a sequence in $[0, 1)$ and $\{\lambda_n\}$ is a sequence in $[a, b]$. If $\{\alpha_n\}$ and $\{\lambda_n\}$ are chosen so that $\{\lambda_n\} \in [a, b]$ for some a, b with $0 < a < b < 2\alpha$,

$$\lim_{n \rightarrow \infty} \alpha_n = 0, \quad \sum_{n=1}^{\infty} \alpha_n = \infty, \quad \sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty, \quad \sum_{n=1}^{\infty} |\lambda_{n+1} - \lambda_n| < \infty, \quad (1.8)$$

then $\{x_n\}$ converges strongly to $P_{F(S) \cap VI(C, A)} x$.

In 2007, Y. Yao and J.-C. Yao [31] further obtained the following theorem.

Theorem YY. *Let C be a closed convex subset of a real Hilbert space H . Let A be an α -inverse-strongly monotone mapping of C into H , and let S be a nonexpansive mapping of C into itself such that $F(S) \cap \Omega \neq \emptyset$, where Ω denotes the set of solutions of a variational inequality for the α -inverse-strongly monotone mapping. Suppose that $x_1 = u \in C$ and $\{x_n\}, \{y_n\}$ are given by*

$$\begin{aligned} x_1 &= u \in C, \\ y_n &= P_C(x_n - \lambda_n A x_n), \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n S P_C(I - \lambda_n A) y_n, \quad n \geq 1, \end{aligned} \tag{1.9}$$

where $\{\alpha_n\}, \{\beta_n\}$, and $\{\gamma_n\}$ are three sequences in $[0, 1]$ and $\{\lambda_n\}$ is a sequence in $[0, 2a]$. If $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$, and $\{\lambda_n\}$ are chosen so that $\lambda_n \in [a, b]$ for some a, b with $0 < a < b < 2a$ and

- (a) $\alpha_n + \beta_n + \gamma_n = 1$, for all $n \geq 1$,
- (b) $\lim_{n \rightarrow \infty} \alpha_n = 0, \sum_{n=1}^{\infty} \alpha_n = \infty$,
- (c) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$,
- (d) $\lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0$,

then $\{x_n\}$ converges strongly to $P_{F(S) \cap \Omega} u$.

In this work, motivated by the above results, we consider the problem of finding a common element in the fixed point set of a strict pseudocontraction and in the zero set of a nonlinear mapping which is the sum of a maximal monotone operator and a inverse strongly monotone mapping. Strong convergence theorems of common elements are established in real Hilbert spaces. The results presented in this paper improve and extend the corresponding results announced by Iiduka and Takahashi [7] and Y. Yao and J.-C. Yao [31].

In order to prove our main results, we also need the following lemmas.

Lemma 1.1 (see [22]). *Let C be a nonempty closed convex subset of a Hilbert space H , $A : C \rightarrow H$ a mapping, and $M : H \rightarrow 2^H$ a maximal monotone mapping. Then,*

$$F(J_r(I - rA)) = (A + M)^{-1}(0), \quad \forall r > 0. \tag{1.10}$$

Lemma 1.2 (see [1]). *Let C be a nonempty closed convex subset of a real Hilbert space H and $S : C \rightarrow C$ a κ -strict pseudocontraction with a fixed point. Define $S_a : C \rightarrow C$ by $S_a x = ax + (1 - a)Sx$ for each $x \in C$. If $a \in [\kappa, 1)$, then S_a is nonexpansive with $F(S_a) = F(S)$.*

Lemma 1.3 (see [25]). *Let C be a nonempty closed convex subset of a Hilbert space H and $S : C \rightarrow C$ a κ -strict pseudocontraction. Then,*

- (a) S is $((1 + \kappa)/(1 - \kappa))$ -Lipschitz,
- (b) $I - S$ is demi-closed, this is, if $\{x_n\}$ is a sequence in C with $x_n \rightharpoonup x$ and $x_n - Sx_n \rightarrow 0$, then $x \in F(S)$.

Lemma 1.4 (see [28]). Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Hilbert space H , and let $\{\beta_n\}$ be a sequence in $(0, 1)$ with

$$0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1. \quad (1.11)$$

Suppose that $x_{n+1} = (1 - \beta_n)y_n + \beta_n x_n$ for all integers $n \geq 1$ and

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (1.12)$$

Then, $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$.

Lemma 1.5 (see [29]). Assume that $\{\alpha_n\}$ is a sequence of nonnegative real numbers such that

$$\alpha_{n+1} \leq (1 - \gamma_n)\alpha_n + \delta_n, \quad (1.13)$$

where $\{\gamma_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a sequence such that

- (a) $\sum_{n=1}^{\infty} \gamma_n = \infty$,
- (b) $\limsup_{n \rightarrow \infty} \delta_n / \gamma_n \leq 0$ or $\sum_{n=1}^{\infty} |\delta_n| < \infty$.

Then, $\lim_{n \rightarrow \infty} \alpha_n = 0$.

Lemma 1.6 (see [24]). Let H be a Hilbert space and M a maximal monotone operator on H . Then, the following holds:

$$\|J_r x - J_s x\|^2 \leq \frac{r-x}{r} \langle J_r x - J_s x, J_r x - x \rangle, \quad \forall s, t > 0, \quad x \in H, \quad (1.14)$$

where $J_r = (I + rM)^{-1}$ and $J_s = (I + sM)^{-1}$.

2. Main Results

Theorem 2.1. Let H be a real Hilbert space H and C a nonempty close and convex subset of H . Let $M : H \rightarrow 2^H$ and $W : H \rightarrow 2^H$ two maximal monotone operators such that $D(M) \subset C$ and $D(W) \subset C$, respectively. Let $S : C \rightarrow C$ be a κ -strict pseudocontraction, $A : C \rightarrow H$ an α -inverse strongly monotone mapping, and $B : C \rightarrow H$ a β -inverse strongly monotone mapping. Assume that $\mathcal{F} := F(S) \cap (A + M)^{-1}(0) \cap (B + W)^{-1}(0) \neq \emptyset$. Let $\{x_n\}$ be a sequence generated in the following manner:

$$\begin{aligned} x_1 &\in C, \\ y_n &= J_{s_n}(x_n - s_n B x_n), \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n (\delta_n J_{r_n}(y_n - r_n A y_n) + (1 - \delta_n) S J_{r_n}(y_n - r_n A y_n)), \quad \forall n \geq 1, \end{aligned} \quad (2.1)$$

where $u \in C$ is a fixed element, $J_{r_n} = (I + r_n M)^{-1}$ and $J_{s_n} = (I + s_n W)^{-1}$, $\{r_n\}$ is a sequence in $(0, 2\alpha)$, $\{s_n\}$ is a sequence in $(0, 2\beta)$ and $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, and $\{\delta_n\}$ are sequences in $[0, 1]$.

Assume that the following restrictions are satisfied:

- (a) $0 < a \leq r_n \leq b < 2\alpha$, $\lim_{n \rightarrow \infty} (r_n - r_{n+1}) = 0$,
- (b) $0 < c \leq s_n \leq d < 2\beta$, $\lim_{n \rightarrow \infty} (s_n - s_{n+1}) = 0$,
- (c) $0 \leq \kappa \leq \delta_n < e < 1$, $\lim_{n \rightarrow \infty} (\delta_n - \delta_{n+1}) = 0$,
- (d) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (e) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \liminf_{n \rightarrow \infty} \beta_n < 1$.

Then, the sequence $\{x_n\}$ converges strongly to $q = P_{\mathcal{F}}u$.

Proof. The proof is split into five steps.

Step 1. Show that $\{x_n\}$ is bounded.

Note that $(I - r_n A)$ and $(I - s_n B)$ are nonexpansive for each fixed $n \geq 1$. Indeed, we see from the restriction (a) that

$$\begin{aligned} \|(I - r_n A)x - (I - r_n A)y\|^2 &= \|x - y\|^2 - 2r_n \langle x - y, Ax - Ay \rangle + r_n^2 \|Ax - Ay\|^2 \\ &\leq \|x - y\|^2 - r_n(2\alpha - r_n) \|Ax - Ay\|^2 \\ &\leq \|x - y\|^2, \quad \forall x, y \in C. \end{aligned} \quad (2.2)$$

This shows that $(I - r_n A)$ is nonexpansive for each fixed $n \geq 1$, so is $(I - s_n B)$. Put

$$S_n x = \delta_n x + (1 - \delta_n) Sx, \quad \forall x \in C. \quad (2.3)$$

In view of the restriction (c), we obtain from Lemma 1.2 that S_n is a nonexpansive mapping with $F(S_n) = F(S)$ for each fixed $n \geq 1$. Fixing $p \in \mathcal{F}$ and since J_{r_n} and $I - r_n A$ are nonexpansive, we see that

$$\begin{aligned} \|x_{n+1} - p\| &\leq \alpha_n \|u - p\| + \beta_n \|x_n - p\| + \gamma_n \|S_n J_{r_n} (y_n - r_n A y_n) - p\| \\ &\leq \alpha_n \|u - p\| + \beta_n \|x_n - p\| + \gamma_n \|J_{r_n} (y_n - r_n A y_n) - p\| \\ &\leq \alpha_n \|u - p\| + \beta_n \|x_n - p\| + \gamma_n \|y_n - p\| \\ &\leq \alpha_n \|u - p\| + (1 - \alpha_n) \|x_n - p\|. \end{aligned} \quad (2.4)$$

By mathematical inductions, we see that $\{x_n\}$ is bounded and so is $\{y_n\}$. This completes Step 1.

Step 2. Show that $\|x_{n+1} - x_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Notice from Lemma 1.6 that

$$\begin{aligned}
\|y_{n+1} - y_n\| &\leq \|(x_{n+1} - s_{n+1}Bx_{n+1}) - (x_n - s_nBx_n)\| \\
&\quad + \|J_{s_{n+1}}(x_n - s_nBx_n) - J_{s_n}(x_n - s_nBx_n)\| \\
&\leq \|x_{n+1} - x_n\| + |s_{n+1} - s_n|\|Bx_n\| \\
&\quad + \frac{|s_{n+1} - s_n|}{s_{n+1}} \|J_{s_{n+1}}(x_n - s_nBx_n) - (x_n - s_nBx_n)\| \\
&\leq \|x_{n+1} - x_n\| + 2M_1|s_{n+1} - s_n|,
\end{aligned} \tag{2.5}$$

where M_1 is an appropriate constant such that

$$M_1 = \max \left\{ \sup_{n \geq 1} \{\|Bx_n\|\}, \sup_{n \geq 1} \left\{ \frac{\|J_{s_{n+1}}(x_n - s_nBx_n) - (x_n - s_nBx_n)\|}{s_{n+1}} \right\} \right\}. \tag{2.6}$$

Put

$$z_n = J_{r_n}(y_n - r_nAy_n), \quad \forall n \geq 1. \tag{2.7}$$

In a similar way, we can obtain from Lemma 1.6 that

$$\begin{aligned}
\|z_{n+1} - z_n\| &\leq \|(y_{n+1} - r_{n+1}Ay_{n+1}) - (y_n - r_nAy_n)\| \\
&\quad + \|J_{r_{n+1}}(y_n - r_nAy_n) - J_{r_n}(y_n - r_nAy_n)\| \\
&\leq \|y_{n+1} - y_n\| + |r_{n+1} - r_n|\|Ay_n\| \\
&\quad + \frac{|r_{n+1} - r_n|}{r_{n+1}} \|J_{r_{n+1}}(y_n - r_nAy_n) - (y_n - r_nAy_n)\| \\
&\leq \|y_{n+1} - y_n\| + 2M_2|r_{n+1} - r_n|,
\end{aligned} \tag{2.8}$$

where M_2 is an appropriate constant such that

$$M_2 = \max \left\{ \sup_{n \geq 1} \{\|Ay_n\|\}, \sup_{n \geq 1} \left\{ \frac{\|J_{r_{n+1}}(y_n - r_nAy_n) - (y_n - r_nAy_n)\|}{r_{n+1}} \right\} \right\}. \tag{2.9}$$

Substituting (2.5) into (2.8) yields that

$$\|z_{n+1} - z_n\| \leq \|x_{n+1} - x_n\| + M_3(|s_{n+1} - s_n| + |r_{n+1} - r_n|), \tag{2.10}$$

where M_3 is an appropriate constant such that

$$M_3 = \max\{2M_1, 2M_2\}. \quad (2.11)$$

It follows from (2.10) that

$$\begin{aligned} \|S_{n+1}z_{n+1} - S_n z_n\| &\leq \|z_{n+1} - z_n\| + \|z_n - S_n z_n\| |\delta_n - \delta_{n+1}| \\ &\leq \|x_{n+1} - x_n\| + M_4(|s_{n+1} - s_n| + |r_{n+1} - r_n| + |\delta_n - \delta_{n+1}|), \end{aligned} \quad (2.12)$$

where M_4 is an appropriate constant such that

$$M_4 = \max\left\{\sup_{n \geq 1}\{\|z_n - S_n z_n\|\}, M_3\right\}. \quad (2.13)$$

Put

$$l_n = \frac{x_{n+1} - \beta_n x_n}{1 - \beta_n}, \quad \forall n \geq 1. \quad (2.14)$$

Note that

$$\begin{aligned} l_{n+1} - l_n &= \frac{\alpha_{n+1}u + \gamma_{n+1}S_{n+1}z_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n u + \gamma_n S_n z_n}{1 - \beta_n} \\ &= \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n}\right)u + \frac{\gamma_{n+1}}{1 - \beta_{n+1}}(S_{n+1}z_{n+1} - S_n z_n) \\ &\quad + \left(\frac{\gamma_{n+1}}{1 - \beta_{n+1}} - \frac{\gamma_n}{1 - \beta_n}\right)S_n z_n \\ &= \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n}\right)(u - S_n z_n) + \frac{\gamma_{n+1}}{1 - \beta_{n+1}}(S_{n+1}z_{n+1} - S_n z_n). \end{aligned} \quad (2.15)$$

It follows from (2.12) that

$$\begin{aligned} \|l_{n+1} - l_n\| &\leq \left|\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n}\right| \|u - S_n z_n\| + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \|S_{n+1}z_{n+1} - S_n z_n\| \\ &\leq \left|\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n}\right| \|u - S_n z_n\| + \|x_{n+1} - x_n\| \\ &\quad + M_4(|s_{n+1} - s_n| + |r_{n+1} - r_n| + |\delta_n - \delta_{n+1}|). \end{aligned} \quad (2.16)$$

This in turn implies from the restrictions (a)–(e) that

$$\limsup_{n \rightarrow \infty} (\|l_{n+1} - l_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (2.17)$$

From Lemma 1.4, we obtain that

$$\lim_{n \rightarrow \infty} \|I_n - x_n\| = 0. \quad (2.18)$$

Notice that

$$x_{n+1} - x_n = (1 - \beta_n)(I_n - x_n). \quad (2.19)$$

It follows that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (2.20)$$

This completes Step 2.

Step 3. Show that $\|x_n - Sx_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Since J_{r_n} and J_{s_n} are nonexpansive, we see that

$$\|z_n - p\|^2 \leq \|x_n - p\|^2 - r_n(2\alpha - r_n)\|Ay_n - Ap\|^2, \quad (2.21)$$

$$\|y_n - p\|^2 \leq \|x_n - p\|^2 - s_n(2\beta - s_n)\|Bx_n - Bp\|^2. \quad (2.22)$$

It follows from (2.21) that

$$\begin{aligned} \|x_{n+1} - p\|^2 &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|S_n z_n - p\|^2 \\ &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|z_n - p\|^2 \\ &\leq \alpha_n \|u - p\|^2 + \|x_n - p\|^2 - \gamma_n r_n (2\alpha - r_n) \|Ay_n - Ap\|^2. \end{aligned} \quad (2.23)$$

This in turn implies that

$$\gamma_n r_n (2\alpha - r_n) \|Ay_n - Ap\|^2 \leq \alpha_n \|u - p\|^2 + \|x_n - x_{n+1}\| (\|x_n - p\| + \|x_{n+1} - p\|). \quad (2.24)$$

In view of (2.20), we see from the restrictions (a), (d), and (e) that

$$\lim_{n \rightarrow \infty} \|Ay_n - Ap\| = 0. \quad (2.25)$$

It follows from (2.22) that

$$\begin{aligned}
\|x_{n+1} - p\|^2 &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|S_n z_n - p\|^2 \\
&\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|J_{r_n}(y_n - r_n A y_n) - p\|^2 \\
&\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|y_n - p\|^2 \\
&\leq \alpha_n \|u - p\|^2 + \|x_n - p\|^2 - \gamma_n s_n (2\beta - s_n) \|Bx_n - Bp\|^2.
\end{aligned} \tag{2.26}$$

This in turn implies that

$$\gamma_n s_n (2\beta - s_n) \|Bx_n - Bp\|^2 \leq \alpha_n \|u - p\|^2 + \|x_n - x_{n+1}\| (\|x_n - p\| + \|x_{n+1} - p\|). \tag{2.27}$$

In view of (2.20), we see from the restrictions (a), (d), and (e) that

$$\lim_{n \rightarrow \infty} \|Bx_n - Bp\| = 0. \tag{2.28}$$

Since J_{r_n} is firmly nonexpansive, we obtain that

$$\begin{aligned}
\|z_n - p\|^2 &= \|J_{r_n}(y_n - r_n A y_n) - J_{r_n}(p - r_n A p)\|^2 \\
&\leq \langle z_n - p, (y_n - r_n A y_n) - (p - r_n A p) \rangle \\
&= \frac{1}{2} \left(\|z_n - p\|^2 + \|(y_n - r_n A y_n) - (p - r_n A p)\|^2 \right. \\
&\quad \left. - \|(z_n - p) - ((y_n - r_n A y_n) - (p - r_n A p))\|^2 \right) \\
&\leq \frac{1}{2} \left(\|z_n - p\|^2 + \|y_n - p\|^2 - \|z_n - y_n + r_n(A y_n - A p)\|^2 \right) \\
&= \frac{1}{2} \left(\|z_n - p\|^2 + \|y_n - p\|^2 - \|z_n - y_n\|^2 - r_n^2 \|A y_n - A p\|^2 \right. \\
&\quad \left. - 2r_n \langle z_n - y_n, A y_n - A p \rangle \right) \\
&\leq \frac{1}{2} \left(\|z_n - p\|^2 + \|y_n - p\|^2 - \|z_n - y_n\|^2 + 2r_n \|z_n - y_n\| \|A y_n - A p\| \right) \\
&\leq \frac{1}{2} \left(\|z_n - p\|^2 + \|x_n - p\|^2 - \|z_n - y_n\|^2 + 2r_n \|z_n - y_n\| \|A y_n - A p\| \right).
\end{aligned} \tag{2.29}$$

This in turn implies that

$$\|z_n - p\|^2 \leq \|x_n - p\|^2 - \|z_n - y_n\|^2 + 2r_n \|z_n - y_n\| \|A y_n - A p\|. \tag{2.30}$$

In a similar way, we can obtain that

$$\|y_n - p\|^2 \leq \|x_n - p\|^2 - \|y_n - x_n\|^2 + 2s_n \|y_n - x_n\| \|Bx_n - Bp\|. \quad (2.31)$$

In view of (2.30), we see that

$$\begin{aligned} \|x_{n+1} - p\|^2 &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|S_n z_n - p\|^2 \\ &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|z_n - p\|^2 \\ &\leq \alpha_n \|u - p\|^2 + \|x_n - p\|^2 - \gamma_n \|z_n - y_n\|^2 + 2r_n \|z_n - y_n\| \|Ay_n - Ap\|. \end{aligned} \quad (2.32)$$

It follows that

$$\begin{aligned} \gamma_n \|z_n - y_n\|^2 &\leq \alpha_n \|u - p\|^2 + \|x_n - x_{n+1}\| (\|x_n - p\| + \|x_{n+1} - p\|) \\ &\quad + 2r_n \|z_n - y_n\| \|Ay_n - Ap\|. \end{aligned} \quad (2.33)$$

In view of (2.25), we obtain from the restrictions (d) and (e) that

$$\lim_{n \rightarrow \infty} \|z_n - y_n\| = 0. \quad (2.34)$$

Notice from (2.31), we see that

$$\begin{aligned} \|x_{n+1} - p\|^2 &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|S_n z_n - p\|^2 \\ &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|z_n - p\|^2 \\ &\leq \alpha_n \|u - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|y_n - p\|^2 \\ &\leq \alpha_n \|u - p\|^2 + \|x_n - p\|^2 - \gamma_n \|y_n - x_n\|^2 + 2s_n \|y_n - x_n\| \|Bx_n - Bp\|. \end{aligned} \quad (2.35)$$

It follows that

$$\begin{aligned} \gamma_n \|y_n - x_n\|^2 &\leq \alpha_n \|u - p\|^2 + \|x_n - x_{n+1}\| (\|x_n - p\| + \|x_{n+1} - p\|) \\ &\quad + 2s_n \|y_n - x_n\| \|Bx_n - Bp\|. \end{aligned} \quad (2.36)$$

In view of (2.28), we obtain from the restrictions (d) and (e) that

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \quad (2.37)$$

Combining (2.34) with (2.37) yields that

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0. \quad (2.38)$$

Note that

$$x_{n+1} - x_n = \alpha_n(u - x_n) + \gamma_n(S_n z_n - x_n). \quad (2.39)$$

In view of (2.20), we see from the restriction (d) that

$$\lim_{n \rightarrow \infty} \|S_n z_n - x_n\| = 0. \quad (2.40)$$

Note that

$$S z_n - x_n = \frac{S_n z_n - x_n}{1 - \delta_n} + \frac{\delta_n(x_n - z_n)}{1 - \delta_n}. \quad (2.41)$$

From (2.38) and (2.40), we get from the restriction (c) that

$$\lim_{n \rightarrow \infty} \|S z_n - x_n\| = 0. \quad (2.42)$$

Notice that

$$\|S x_n - x_n\| \leq \|S x_n - S z_n\| + \|S z_n - x_n\|. \quad (2.43)$$

In view of (2.38) and (2.42), we see from Lemma 1.3 that

$$\lim_{n \rightarrow \infty} \|S x_n - x_n\| = 0. \quad (2.44)$$

This completes Step 3.

Step 4. Show that $\limsup_{n \rightarrow \infty} \langle u - q, x_n - q \rangle \leq 0$, where $q = P_{\mathcal{F}} u$.

To show it, we may choose a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle u - q, x_n - q \rangle = \limsup_{i \rightarrow \infty} \langle u - q, x_{n_i} - q \rangle. \quad (2.45)$$

Since $\{x_{n_i}\}$ is bounded, we can choose a subsequence $\{x_{n_{i_j}}\}$ of $\{x_{n_i}\}$ converging weakly to \hat{x} . We may, without loss of generality, assume that $x_{n_{i_j}} \rightharpoonup \hat{x}$, where \rightharpoonup denotes the weak convergence. Next, we prove that $\hat{x} \in \mathcal{F}$. In view of (2.44), we can conclude from Lemma 1.3 that $\hat{x} \in F(S)$ easily. Notice that

$$y_n - r_n A y_n \in z_n + r_n M z_n. \quad (2.46)$$

Let $\mu \in M\nu$. Since M is monotone, we have

$$\left\langle \frac{y_n - z_n}{r_n} - Ay_n - \mu, z_n - \nu \right\rangle \geq 0. \quad (2.47)$$

In view of the restriction (a), we see from (2.34) that

$$\langle -A\bar{x} - \mu, \bar{x} - \nu \rangle \geq 0. \quad (2.48)$$

This implies that $-A\bar{x} \in M\bar{x}$, that is, $\bar{x} \in (A + M)^{-1}(0)$. In similar way, we can obtain that $\bar{x} \in (B + W)^{-1}(0)$. This proves that $\bar{x} \in \mathcal{F}$. It follows from (2.45) that

$$\limsup_{n \rightarrow \infty} \langle u - q, x_n - q \rangle \leq 0. \quad (2.49)$$

This completes Step 4.

Step 5. Show that $x_n \rightarrow q$ as $n \rightarrow \infty$.

Notice that

$$\begin{aligned} \|x_{n+1} - q\|^2 &= \alpha_n \langle u - q, x_{n+1} - q \rangle + \beta_n \langle x_n - q, x_{n+1} - q \rangle \\ &\quad + \gamma_n \langle S_n J_{r_n}(y_n - r_n A y_n) - q, x_{n+1} - q \rangle \\ &\leq \alpha_n \langle u - q, x_{n+1} - q \rangle + \frac{\beta_n}{2} (\|x_n - q\|^2 + \|x_{n+1} - q\|^2) \\ &\quad + \frac{\gamma_n}{2} (\|S_n J_{r_n}(y_n - r_n A y_n) - q\|^2 + \|x_{n+1} - q\|^2) \\ &\leq \alpha_n \langle u - q, x_{n+1} - q \rangle + \frac{\beta_n}{2} (\|x_n - q\|^2 + \|x_{n+1} - q\|^2) \\ &\quad + \frac{\gamma_n}{2} (\|y_n - q\|^2 + \|x_{n+1} - q\|^2) \\ &\leq \alpha_n \langle u - q, x_{n+1} - q \rangle + \frac{1 - \alpha_n}{2} (\|x_n - q\|^2 + \|x_{n+1} - q\|^2). \end{aligned} \quad (2.50)$$

This in turn implies that

$$\|x_{n+1} - q\|^2 \leq (1 - \alpha_n) \|x_n - q\|^2 + 2\alpha_n \langle u - q, x_{n+1} - q \rangle. \quad (2.51)$$

In view of (2.49), we conclude from Lemma 1.5 that

$$\lim_{n \rightarrow \infty} \|x_n - q\| = 0. \quad (2.52)$$

This completes Step 5. This whole proof is completed. \square

If S is a nonexpansive mapping and $\delta_n = 0$, then Theorem 2.1 is reduced to the following.

Corollary 2.2. *Let H be a real Hilbert space H and C a nonempty close and convex subset of H . Let $M : H \rightarrow 2^H$ and $W : H \rightarrow 2^H$ be two maximal monotone operators such that $D(M) \subset C$ and $D(W) \subset C$, respectively. Let $S : C \rightarrow C$ be a nonexpansive mapping, $A : C \rightarrow H$ an α -inverse strongly monotone mapping and $B : C \rightarrow H$ a β -inverse strongly monotone mapping. Assume that $\mathcal{F} := F(S) \cap (A + M)^{-1}(0) \cap (B + W)^{-1}(0) \neq \emptyset$. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{aligned} x_1 &\in C, \\ y_n &= J_{s_n}(x_n - s_n Bx_n), \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n S J_{r_n}(y_n - r_n A y_n), \quad \forall n \geq 1, \end{aligned} \tag{2.53}$$

where $u \in C$ is a fixed element, $J_{r_n} = (I + r_n M)^{-1}$ and $J_{s_n} = (I + s_n W)^{-1}$, $\{r_n\}$ is a sequence in $(0, 2\alpha)$, $\{s_n\}$ is a sequence in $(0, 2\beta)$ and $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ are sequences in $[0, 1]$. Assume that the following restrictions are satisfied:

- (a) $0 < a \leq r_n \leq b < 2\alpha$, $\lim_{n \rightarrow \infty} (r_n - r_{n+1}) = 0$,
- (b) $0 < c \leq s_n \leq d < 2\beta$, $\lim_{n \rightarrow \infty} (s_n - s_{n+1}) = 0$,
- (c) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (d) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \liminf_{n \rightarrow \infty} \beta_n < 1$.

Then, the sequence $\{x_n\}$ converges strongly to $q = P_{\mathcal{F}}u$.

Next, we consider the problem of finding common fixed points of three strict pseudocontractions.

Theorem 2.3. *Let C be a nonempty closed convex subset of a real Hilbert space H and P_C the metric projection from H onto C . Let $S : C \rightarrow C$ be a κ -strict pseudocontraction, $T_A : C \rightarrow H$ an α -strict pseudocontraction, and $B : C \rightarrow H$ a β -strict pseudocontraction. Assume that $\mathcal{F} := F(S) \cap F(T_A) \cap F(T_B) \neq \emptyset$. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{aligned} x_1 &\in C, \\ z_n &= (1 - s_n)x_n + s_n T_B x_n, \\ y_n &= (1 - r_n)z_n + r_n T_A z_n \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n (\delta_n y_n + (1 - \delta_n) S y_n), \quad \forall n \geq 1, \end{aligned} \tag{2.54}$$

where $u \in C$ is a fixed element, $\{r_n\}$ is a sequence in $(0, 1 - \alpha)$, $\{s_n\}$ is a sequence in $(0, 1 - \beta)$, and $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, and $\{\delta_n\}$ are sequences in $[0, 1]$. Assume that the following restrictions are satisfied

- (a) $0 < a \leq r_n \leq b < 1 - \alpha$, $\lim_{n \rightarrow \infty} (r_n - r_{n+1}) = 0$,
- (b) $0 < c \leq s_n \leq d < 1 - \beta$, $\lim_{n \rightarrow \infty} (s_n - s_{n+1}) = 0$,
- (c) $0 \leq \kappa \leq \delta_n < e < 1$, $\lim_{n \rightarrow \infty} (\delta_n - \delta_{n+1}) = 0$,

- (d) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (e) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \liminf_{n \rightarrow \infty} \beta_n < 1$.

Then, the sequence $\{x_n\}$ converges strongly to $q = P_{\overline{C}}u$.

Proof. Putting $A = I - T_A$, we see that A is $((1 - \alpha)/2)$ -inverse strongly monotone. We also have $F(T_A) = VI(C, A)$ and $P_C(x_n - r_n Ax_n) = (1 - r_n)x_n + r_n T x_n$. Putting $B = I - T_B$, we see that B is $(1 - \beta)/2$ -inverse strongly monotone. We also have $F(T_B) = VI(C, B)$ and $P_C(x_n - s_n Bx_n) = (1 - s_n)x_n + s_n Ru_n$. In view of Theorem 2.1, we can obtain the desired results immediately. \square

Acknowledgments

The authors are extremely grateful to the referees for useful suggestions that improved the contents of the paper. This work was supported by the National Natural Science Foundation of China under Grant no. 70871081 and Important Science and Technology Research Project of Henan province, China (102102210022).

References

- [1] F. E. Browder and W. V. Petryshyn, "Construction of fixed points of nonlinear mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 20, pp. 197–228, 1967.
- [2] Y. J. Cho, X. Qin, and J. I. Kang, "Convergence theorems based on hybrid methods for generalized equilibrium problems and fixed point problems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 9, pp. 4203–4214, 2009.
- [3] Y. J. Cho and X. Qin, "Systems of generalized nonlinear variational inequalities and its projection methods," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 12, pp. 4443–4451, 2008.
- [4] Y. J. Cho, S. M. Kang, and X. Qin, "On systems of generalized nonlinear variational inequalities in Banach spaces," *Applied Mathematics and Computation*, vol. 206, no. 1, pp. 214–220, 2008.
- [5] Y. J. Cho, S. M. Kang, and X. Qin, "Approximation of common fixed points of an infinite family of nonexpansive mappings in Banach spaces," *Computers & Mathematics with Applications*, vol. 56, no. 8, pp. 2058–2064, 2008.
- [6] Y. J. Cho, S. M. Kang, and X. Qin, "Some results on k -strictly pseudo-contractive mappings in Hilbert spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 5, pp. 1956–1964, 2009.
- [7] H. Iiduka and W. Takahashi, "Strong convergence theorems for nonexpansive mappings and inverse-strongly monotone mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 61, no. 3, pp. 341–350, 2005.
- [8] S. Park and B. G. Kang, "Generalized variational inequalities and fixed point theorems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 31, no. 1-2, pp. 207–216, 1998.
- [9] S. Park and S. Kum, "An application of a Browder-type fixed point theorem to generalized variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 218, no. 2, pp. 519–526, 1998.
- [10] S. Park, "Acyclic maps, minimax inequalities and fixed points," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 24, no. 11, pp. 1549–1554, 1995.
- [11] S. Park, "Fixed point theory of multimaps in abstract convex uniform spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 7-8, pp. 2468–2480, 2009.
- [12] S. Park, "Recent results in analytical fixed point theory," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 63, no. 5-7, pp. 977–986, 2005.
- [13] S. Park, "The KKM principle implies many fixed point theorems," *Topology and Its Applications*, vol. 135, no. 1-3, pp. 197–206, 2004.
- [14] S. Park, "Fixed point theorems in locally G -convex spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 48, no. 6, pp. 869–879, 2002.
- [15] S. Park, "Fixed point theory of multimaps in abstract convex uniform spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 7-8, pp. 2468–2480, 2009.

- [16] W. Takahashi and M. Toyoda, "Weak convergence theorems for nonexpansive mappings and monotone mappings," *Journal of Optimization Theory and Applications*, vol. 118, no. 2, pp. 417–428, 2003.
- [17] X. Qin, Y. J. Cho, and S. M. Kang, "Convergence theorems of common elements for equilibrium problems and fixed point problems in Banach spaces," *Journal of Computational and Applied Mathematics*, vol. 225, no. 1, pp. 20–30, 2009.
- [18] X. Qin and Y. Su, "Approximation of a zero point of accretive operator in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 415–424, 2007.
- [19] X. Qin and Y. Su, "Strong convergence theorems for relatively nonexpansive mappings in a Banach space," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 6, pp. 1958–1965, 2007.
- [20] X. Qin, M. Shang, and Y. Su, "Strong convergence of a general iterative algorithm for equilibrium problems and variational inequality problems," *Mathematical and Computer Modelling*, vol. 48, no. 7-8, pp. 1033–1046, 2008.
- [21] A. Tada and W. Takahashi, "Weak and strong convergence theorems for a nonexpansive mapping and an equilibrium problem," *Journal of Optimization Theory and Applications*, vol. 133, no. 3, pp. 359–370, 2007.
- [22] K. Aoyama, Y. Kimura, W. Takahashi, and M. Toyoda, "On a strongly nonexpansive sequence in Hilbert spaces," *Journal of Nonlinear and Convex Analysis*, vol. 8, no. 3, pp. 471–489, 2007.
- [23] X. Qin, Y. J. Cho, S. M. Kang, and H. Zhou, "Convergence of a modified Halpern-type iteration algorithm for quasi- ϕ -nonexpansive mappings," *Applied Mathematics Letters*, vol. 22, no. 7, pp. 1051–1055, 2009.
- [24] S. Takahashi, W. Takahashi, and M. Toyoda, "Strong convergence theorems for maximal monotone operators with nonlinear mappings in Hilbert spaces," *Journal of Optimization Theory and Applications*, vol. 147, no. 1, pp. 27–41, 2010.
- [25] G. Marino and H.-K. Xu, "Weak and strong convergence theorems for strict pseudo-contractions in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 336–346, 2007.
- [26] X. Qin, M. Shang, and Y. Su, "A general iterative method for equilibrium problems and fixed point problems in Hilbert spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 11, pp. 3897–3909, 2008.
- [27] X. Qin, S.-S. Chang, and Y. J. Cho, "Iterative methods for generalized equilibrium problems and fixed point problems with applications," *Nonlinear Analysis: Real World Applications*, vol. 11, no. 4, pp. 2963–2972, 2010.
- [28] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter nonexpansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [29] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.
- [30] X. Qin, Y. J. Cho, J. I. Kang, and S. M. Kang, "Strong convergence theorems for an infinite family of nonexpansive mappings in Banach spaces," *Journal of Computational and Applied Mathematics*, vol. 230, no. 1, pp. 121–127, 2009.
- [31] Y. Yao and J.-C. Yao, "On modified iterative method for nonexpansive mappings and monotone mappings," *Applied Mathematics and Computation*, vol. 186, no. 2, pp. 1551–1558, 2007.
- [32] X. Qin, J. I. Kang, and X. Qin, "On quasi-variational inclusions and asymptotically strict pseudocontractions," *Journal of Nonlinear and Convex Analysis*, vol. 11, no. 3, pp. 441–453, 2010.