

ON THE LOCATION OF THE ZEROS OF ANALYTIC FUNCTIONS

K.K. DEWAN

Department of Mathematics
Jamia Millia University
New Delhi - 110025
India

N.K. GOVIL

Department of Algebra, Combinatorics and Analysis
Division of Mathematics
Auburn University
Auburn, AL 36849
U.S.A.

(Received May 1, 1989)

ABSTRACT: In this paper we obtain regions containing all the zeros of a class of analytic functions whose coefficients are subject to certain conditions. Our results sharpen some of the results known in this direction. Also we give some examples to show that in some cases the regions obtained by our results are considerably sharper than the regions obtained by known results.

KEY WORDS AND PHRASES. Zeros of polynomials, rational functions, and other analytic functions.

1980 AMS SUBJECT CLASSIFICATION CODE. 30C15.

1. INTRODUCTION AND STATEMENT OF RESULTS.

In this paper we obtain the regions containing all the zeros of certain classes of analytic functions. Our results improve some of the results known in this direction. We shall in fact be proving the following results which sharpen some of the results of Aziz and Mohammad [1] and of Govil and Rahman [2].

THEOREM 1. Let $f(z) = \sum_{j=0}^{\infty} a_j z^j \neq 0$, be analytic in $|z| \leq t$.
If $|\arg a_j| \leq \alpha \leq \pi/2$, $j = 0, 1, 2, \dots$, and for some finite nonnegative integer
 k , $|a_0| \leq t |a_1| \leq \dots \leq t^k |a_k| \geq t^{k+1} |a_{k+1}| \geq \dots$, then $f(z)$ does not vanish in
 $|z| < R_1$,

where

$$R_1 = \frac{-t^2 |b| (M_1 - |a_0| t) + (t^4 |b|^2 (M_1 - |a_0| t)^2 + 4t^3 |a_0| M_1^3)^{\frac{1}{2}}}{2M_1^2}. \quad (1.1)$$

Here

$$M_1 = t |a_0| \left\{ \left(\frac{2|a_k|}{|a_0|} t^{k-1} \right) \cos \alpha + \sin \alpha + \frac{2 \sin \alpha}{|a_0|} \sum_{j=1}^{\infty} t^j |a_j| \right\} \quad (1.2)$$

and

$$b = a_0 - ta_1. \quad (1.3)$$

The above theorem sharpens Theorem 6 of Aziz and Mohammad [1]. If we take $k = 0$ in Theorem 1, we get the following corollary which sharpens Corollary 5 of Aziz and Mohammad [1].

COROLLARY 1. Let $f(z) = \sum_{j=0}^{\infty} a_j z^j$ be analytic in $|z| \leq t$.

If $|\arg a_j| \leq \alpha \leq \pi/2$, $j = 0, 1, 2, \dots$, and $|a_0| \geq t |a_1| \geq t^2 |a_2| \geq \dots$, then $f(z)$ does not vanish in

$$|z| < R_2,$$

where

$$R_2 = \frac{-t^2 |b| (M_2 - t |a_0|) + (t^4 |b|^2 (M_2 - t |a_0|)^2 + t^3 |a_0| M_2^3)^{\frac{1}{2}}}{2M_2^2}. \quad (1.4)$$

Here

$$M_2 = t |a_0| \left(\cos \alpha + \sin \alpha + \frac{2 \sin \alpha}{|a_0|} \sum_{j=1}^{\infty} t^j |a_j| \right) \quad (1.5)$$

and

$$b = a_0 - ta_1 \quad (1.6)$$

For $t=1$, the above corollary sharpens Theorem 3 of Govil and Rahman [2].

THEOREM 2. Let $f(z) = \sum_{j=0}^{\infty} a_j z^j$ be analytic in $|z| \leq t$. If

$\operatorname{Re} a_j = \alpha_j$, and $\operatorname{Im} a_j = \beta_j$, $j=0, 1, 2, \dots$, and for some finite nonnegative integer
 k , $\alpha_0 \leq \alpha_1 \leq \dots \leq \alpha_k \geq t^{k+1} \alpha_{k+1} \geq \dots$, then $f(z)$ does not vanish in

$$|z| < R_3,$$

where

$$R_3 = \frac{-t^2 |b| (M_3 - |a_0| t) + \{t^4 |b|^2 (M_3 - |a_0| t)^2 + 4t^3 |a_0| M_3\}^{\frac{1}{2}}}{2M_4^2} \tag{1.7}$$

Here

$$M_3 = \alpha_0 \left\{ 2 \binom{\alpha_k}{\alpha_0} t^{k-1} + \frac{2}{\alpha_0} \sum_{j=0}^{\infty} t^j |\beta_j| \right\}, \tag{1.8}$$

and

$$b = a_0 - \alpha_1. \tag{1.9}$$

This theorem sharpens Theorem 7 of Aziz and Mohammad [1]. If we take $k=0$ in the above Theorem 2, we get the following corollary which generalizes and sharpens Theorem 4 of Govil and Rahman [2].

COROLLARY 2. Let $f(z) = \sum_{j=0}^{\infty} a_j z^j$ be analytic in $|z| \leq t$. If $|a_j| \leq \alpha_j + i\beta_j$, $j=0,1,2, \dots$ and $0 < \alpha_0 > t\alpha_1 > t^2\alpha_2 > \dots$, then $f(z)$ does not vanish in

$$|z| < R_4,$$

where

$$R_4 = \frac{-t^2 |b| (M_4 - |a_0| t) + \{t^4 |b|^2 (M_4 - |a_0| t)^2 + 4t^3 |a_0| M_4\}^{\frac{1}{2}}}{2M_4^2}. \tag{1.10}$$

Here

$$M_4 = \alpha_0 \left(1 + \frac{2}{\alpha_0} \sum_{j=0}^{\infty} |\beta_j| t^j \right), \tag{1.11}$$

and

$$b_0 = a_0 - \alpha_1. \tag{1.12}$$

As remarked earlier, Theorems 1 and 2 are respectively the refinements of Theorems 6 and 7 of Aziz and Mohammad [1]. For the sake of completeness we will verify that Theorem 1 sharpens Theorem 6 of Aziz and Mohammad [1], for which it is sufficient to show that

$$\frac{t}{M} \leq R_1 = \frac{-t^2 |b| (M_1 - |a_0| t) + \{t^4 |b|^2 (M_1 - |a_0| t)^2 + 4M_1^3 t^3 |a_0|\}^{\frac{1}{2}}}{2M_1^2} \tag{1.13}$$

$$= \frac{-t^3 |b| |a_0| (M-1) + t^3 |a_0| \{ |b|^2 (M-1)^2 + 4|a_0|^2 M^3 \}^{\frac{1}{2}}}{2t^2 |a_0|^2 M^2}, \tag{1.14}$$

because as can be easily seen, $M_1 = t|a_0|M$. Now (1.14) is equivalent to

$$2|a_0|M < -|b|(M-1) + \{ |b|^2 (M-1)^2 + 4|a_0|^2 M^3 \}^{\frac{1}{2}},$$

which is true if and only if

$$(M-1)(|a_0| - M|b|) \geq 0, \tag{1.15}$$

and since (1.15) is evidently true by Cauchy's inequality [3, p. 84], the inequality (1.13) is verified.

The fact that Theorem 2 is a refinement of Theorem 7 of [1] can be verified on the same lines and we therefore omit the proof.

In general Theorems 1 and 2 give results which are at least as good as obtainable from Theorems 6 and 7 of Aziz and Mohammad [1], but in some cases the results obtained by Theorems 1 and 2 are significantly better than those obtained respectively from Theorem 6 and 7 of Aziz and Mohammad [1]. We illustrate this by the following examples.

EXAMPLE 1.

$$f(z) = 1 + \left(\frac{1}{2} + \frac{i}{2}\right)z + \left(\frac{1}{8} + \frac{i}{8}\right)z^2 + \left(\frac{1}{32} + \frac{i}{32}\right)z^3 + \dots$$

If we take $t = 2, k = 1, \alpha = \pi/4$, then Theorem 6 of Aziz and Mohammad [1] gives that $f(z)$ does not vanish in $|z| < 0.4714045$, while our Theorem 1 gives that $f(z)$ does not vanish in $|z| < 1.1185882$.

EXAMPLE 2.

$$f(z) = (1+i) + \left(2 + \frac{i}{2}\right)z + \left(3 + \frac{i}{2}\right)z^2 + \left(4 + \frac{i}{2}\right)z^3 + \left(1 + \frac{i}{2}\right)z^4 + \left(1 + \frac{i}{2}\right)z^5 + \left(1 + \frac{i}{2}\right)z^6 + \dots$$

If we take $t = 1, k = 3$ then Theorem 7 of Aziz and Mohammad [1] gives that $f(z)$ does not vanish in $|z| < 0.333 \dots$ while by our Theorem 2, $f(z)$ does not vanish in $|z| < 0.791$.

2. LEMMAS.

We need the following lemmas.

LEMMA 1. If $f(z)$ is analytic in $|z| \leq 1, f(0)=a$ where $|a| < 1, f'(0) = b,$
 $|f(z)| \leq 1$ on $|z|=1,$ then for $|z| < 1,$

$$|f(z)| \leq \frac{(1-|a|)|z|^2 + |b||z| + |a|(1-|a|)}{|a|(1-|a|)|z|^2 + |b||z| + (1-|a|)}.$$

The example $f(z) = \frac{a + \frac{b}{1+a}z - z^2}{1 - \frac{b}{1+a}z - az^2}$ shows that the estimate is sharp.

Above lemma is due to Govil et al [4].

One gets easily from the above lemma, the following.

LEMMA 2. If $f(z)$ is analytic in $|z| \leq R, f(0)=0, f'(0)=b$ and $|f(z)| \leq M$
for $|z| = R,$ then for $|z| \leq R,$

$$|f(z)| \leq \frac{M|z|}{R^2} + \frac{M|z| + R^2|b|}{M + |z||b|}.$$

LEMMA 3. Let $f(z) = \sum_{j=0}^{\infty} a_j z^j$ be analytic in $|z| \leq t.$

If $|\arg a_j| \leq \alpha \leq \pi/2, j=0,1,2, \dots,$ then

$$|ta_j - a_{j-1}| \leq |t|a_j| - |a_{j-1}| |\cos \alpha + (t|a_j| + |a_{j-1}|) \sin \alpha.$$

The proof is omitted as it follows immediately from the inequality (6) in [2].

3. PROOFS OF THE THEOREMS.

Proof of Theorem 1. Clearly $\lim_{j \rightarrow \infty} t^j a_j = 0$. We consider the function

$$\begin{aligned} F(z) &= (z-t)f(z) \\ &= -ta_0 + z \sum_{j=1}^{\infty} (a_{j-1} - ta_j) z^{j-1} \\ &= -ta_0 + G(z), \text{ say.} \end{aligned} \tag{3.1}$$

Using Lemma 3, we get for $|z| = t$,

$$\begin{aligned} |G(z)| &\leq t \left(\sum_{j=1}^{\infty} |t|a_j| - |a_{j-1}| |t|^{j-1} \cos \alpha + \sum_{j=1}^{\infty} (t|a_j| + |a_{j-1}|) t^{j-1} \sin \alpha \right) \\ &= t \cos \alpha \sum_1^k (t|a_j| - |a_{j-1}|) t^{j-1} + \sum_{k+1}^{\infty} (|a_{j-1}| - t|a_j|) t^{j-1} \cos \alpha \\ &\quad + 2 \sin \alpha \sum_{j=1}^{\infty} t^j |a_j| + |a_0| \sin \alpha \\ &= t|a_0| \left\{ \left(\frac{2|a_k|t^k}{|a_0|} - 1 \right) \cos \alpha + \sin \alpha + \frac{2 \sin \alpha}{|a_0|} \sum_{j=1}^{\infty} t^j |a_j| \right\} \\ &= t|a_0| M, \text{ say} \\ &= M_1. \end{aligned}$$

Since $G(0) = 0$, $G'(0) = a_0 - ta_1 = b$, we get by Lemma 2, that for $|z| \leq t$,

$$|G(z)| < \frac{M_1 |z|}{t^2} \cdot \frac{M_1 |z| + t^2 |b|}{M_1 + |b| |z|}. \tag{3.2}$$

Combining (3.1) and (3.2), we get for $|z| \leq t$,

$$\begin{aligned} |F(z)| &\geq t|a_0| - |G(z)| \\ &\geq t|a_0| - \frac{M_1 |z|}{t^2} \cdot \frac{M_1 |z| + t^2 |b|}{M_1 + |b| |z|} \\ &= \frac{-1}{t^2 (M_1 + |b| |z|)} \{ |z|^2 M_1^2 + t^2 |b| |z| (M_1 - t|a_0|) - t^3 |a_0| M_1 \} \\ &> 0, \end{aligned}$$

if

$$|z| < \frac{-t^2 |b| (M_1 - t |a_0|) + [t^4 |b|^2 (M_1 - t |a_0|)^2 + 4t^3 |a_0| M_1^3]^{\frac{1}{2}}}{2M_1^2}$$

= R₁.

It is easy to verify that $R_1 \leq t$. Thus in $|z| \leq t$, $|F(z)| > 0$ if $|z| < R_1$. Consequently $f(z)$ does not vanish in $|z| < R_1$ and the proof of Theorem 1 is complete.

Proof of Theorem 2. Clearly $\lim_{j \rightarrow \infty} \alpha_j t^j = 0$, $\lim_{j \rightarrow \infty} \beta_j t^j = 0$. As before we consider the function $F(z) = (z-t) f(z) = -t a_0 + G(z)$, where $G(z)$ is same as in (3.1). Then on $|z| = t$, we have

$$\begin{aligned} |G(z)| &\leq t \sum_{j=1}^{\infty} |a_{j-1} - t a_j| t^{j-1} \\ &\leq t \left(\sum_{j=1}^{\infty} |\alpha_{j-1} - t \alpha_j| t^{j-1} + \sum_{j=1}^{\infty} (|\beta_{j-1}| + t |\beta_j|) \right) \\ &= t \alpha_0 \left\{ 2 \left(\frac{\alpha_k}{\alpha_0} \right) t^k - 1 + \left(\frac{2}{\alpha_0} \right) \sum_{j=0}^{\infty} t^j |\beta_j| \right\} \\ &= M_3. \end{aligned}$$

Since $G(0) = 0$, $G'(0) = a_0 - t a_1 = b$, we can apply Lemma 2 to the function $G(z)$ and then proceeding on the lines of the proof of Theorem 1, the proof of Theorem 2 can be completed.

REFERENCES

1. Aziz, A. and Mohammad, Q. G., On the zeros of a certain class of polynomials and related analytic functions, J. Math. Anal. Appl. 75(1980), 495-502.
2. Govil, N. K. and Rahman, Q. I., On the Eneström-Kakeya theorem, Tôhoku Math. J. 20(1968), 126-136.
3. Titchmarsh, E. C., The Theory of Functions, 2nd edition, Oxford University Press, 1939.
4. Govil, N. K., Rahman, Q. I., and Schmeisser, G., On the derivative of a polynomial, Illinois J. Math. 23(1979), 319-329.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br