

A TOPOLOGY FOR AUTOMATA II

ARUN K. SRIVASTAVA

Department of Mathematics
Banaras Hindu University
Varansi, India

and

WAGISH SHUKLA

Department of Mathematics
Indian Institute of Technology
New Delhi, India

(Received October 6, 1983)

ABSTRACT. A topology on the state set of an automaton is considered and it is shown that under this topology, genetically closed subsets and primaries, in the sense of Bavel [1] turn out to be precisely the regular closed subsets and minimal regular closed subsets respectively. The concept of a compact automaton is introduced and it is indicated that it can be viewed as a generalization of a finite automaton. Included also is an observation showing that our topological considerations can help recover some of the results of Dörfler [2].

KEY WORDS AND PHRASES. Automata, topological concepts and methods, compact automata, product automata, connectedness.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 68D

1. INTRODUCTION.

Shukla and Srivastava [3] have indicated that it is possible to study certain aspects of automata e.g., the connectivity and separation properties, by using topological concepts and methods. They did this by introducing a topology on the state set of an automaton in a natural way. In this note, we consider a topology dual to this and show that a genetically closed set and hence a primary of an automaton have standard topological analogs. We also introduce the concept of a compact automaton and show that it can be viewed as a generalization of a finite automaton. Included also are a few remarks concerning certain preservation properties of product automata.

For completeness we give the relevant definitions first. An X -automaton is a triple $A = (Q, X, \delta)$ where Q is a set (set of states), X is a semigroup (the input alphabet) with identity e , and $\delta: Q \times X \rightarrow Q$ is a mapping (the transition map) satisfying for all $(q, x, y) \in Q \times X \times X$, $\delta(q, e) = q$ and $\delta(q, xy) = \delta(\delta(q, x), y)$. $B = (Q', X, \delta')$ is a subautomaton of A iff $Q' \subseteq Q$ and δ' is δ restricted to $Q' \times X$. (We shall use δ for its restriction δ' when no ambiguity arises). For a subset $R \subseteq Q$, the set $\delta R = \{\delta(q, x) \mid (q, x) \in R \times X\}$ is the set of successors of R , $\sigma R = \{q \in Q \mid \delta(q, x) \in R \text{ for some } x \in X\}$ is the set of sources of R and $\mu R = \{q \in R \mid \sigma q \subseteq R\}$ is the core of R . (We write σq for $\sigma\{q\}$). If $A = (Q, X, \delta)$ and

$B = (P, X, \gamma)$ are X -automata then f is a homomorphism from A to B if $f: Q \rightarrow P$ is a function such that $f(\delta(q, x)) = \gamma(fq, x)$ for all $(q, x) \in Q \times X$.

If $A = (Q, X, \delta)$ is an X -automaton, then a topology t_A on A can be defined by declaring a subset R of Q closed iff $R = \sigma R$ (cf. [3]). It turns out that

THEOREM 1.1([3]). If $B = (Q', X, \delta)$ and $A = (Q, X, \delta)$ are automata then

- (a) B is a subautomaton of A iff Q' is t_A -open.
- (b) B is a separated subautomaton of A iff Q' is both t_A -open and t_A -closed.
- (c) A is connected iff t_A is connected (in topological sense).
- (d) B is a block of A iff Q' is a component of Q .
- (e) A is retrievable iff t_A is an R_0 -topology.
- (f) A is strongly connected iff t_A is indiscrete.

Henceforth, in the absence of any other specifications, we shall assume Q to carry the topology t_A ; this topology will be referred to as the 'state-set topology'.

All undefined concepts used here are either standard or can be found in Shukla and Srivastava [3] and the references cited therein.

2. CORE OPERATOR.

The core operator, as a self-map on the set of subsets of Q , turns out to be an interior operator in topological sense, i.e., the core operator satisfies

- (a) $\mu Q = Q$;
- (b) $\mu R \subseteq R$ for all $R \subseteq Q$;
- (c) $\mu \mu R = \mu R$ for all $R \subseteq Q$;
- (d) $\mu(R_1 \cap R_2) = \mu R_1 \cap \mu R_2$ for all $R_1, R_2 \subseteq Q$.

Indeed, (a) and (b) are obvious. To see (c), let $q \in \mu \mu R$. Observe that $q \in \mu \mu R$ means that $\sigma q \subseteq \mu R$ which in turn means that for all $q \in Q$ such that $\delta(p, x) = q$ (for some $x \in X$), it is the case that $\sigma p \subseteq R$. So let $r \in Q$ be such that $\delta(r, y) = p$ for some $y \in X$. Then $\delta(r, yx) = q$ and so $r \in \sigma q$. Since $q \in \mu R$, $r \in R$. Thus, $\mu R \subseteq \mu \mu R$. But owing to (b), we also see that $\mu \mu R \subseteq \mu R$. Hence $\mu R = \mu \mu R$ and (c) is true. Finally, to verify (d), let $q \in \mu(R_1 \cap R_2)$. Then $\sigma q \subseteq R_1 \cap R_2$ whence $\mu(R_1 \cap R_2) \subseteq \mu R_1 \cap \mu R_2$. Conversely, if $q \in \mu R_1 \cap \mu R_2$ then $\sigma q \subseteq R_1$ and $\sigma q \subseteq R_2$. It follows that (d) is also true.

The core, being an interior operator for Q , defines a topology on Q ; a subset $R \subseteq Q$ is open with respect to this topology iff $R = \mu R$.

REMARK 2.1. (a) The above topology is saturated in the sense that any intersection of its open sets is open. To see this let $\{R_i : i \in I\}$ be any collection of its open sets. Then $R_i = \mu R_i$, for all $i \in I$. It is enough to show that $\bigcap_{i \in I} R_i \subseteq \mu(\bigcap_{i \in I} R_i)$. Let $q \in \bigcap_{i \in I} R_i$. Then $q \in \mu R_i$, for all $i \in I$ which means that $\sigma q \subseteq R_i$, for all $i \in I$. Hence $\sigma q \subseteq \bigcap_{i \in I} R_i$ showing that $q \in \mu(\bigcap_{i \in I} R_i)$. (b) The topology induced on Q by μ is dual to the topology t_A on Q and for this reason we shall denote it by t_A^* . Moreover, with respect to the topology t_A , μR is the largest closed set contained in R for any $R \subseteq Q$. Also, the successor operator δ is the closure operator for t_A^* and σR is the smallest open set containing R .

Throughout by Q^* , we shall understand the topological space (Q, t_A^*) .

3. REGULAR CLOSED SUBSETS.

A subset of a topological space is called regular closed iff it equals the closure of its interior. In Bavel [1] a subset $R \subseteq Q$ is called genetic iff $\sigma R \subseteq \delta R$, genetic for $M \subseteq Q$ iff R is genetic and $\delta R = M$, genetically closed iff there exists a subset $M \subseteq R$ such that $\sigma M \subseteq \delta M$ and $\delta M = R$, and primary iff R is a minimal genetically closed set.

PROPOSITION 3.1. $R \subseteq Q$ is genetically closed iff it is a regular closed subset of Q^* .

PROOF. Suppose R is genetically closed. Then, as shown in Bavel [1], $\delta\mu R = R$, i.e., R is a regular closed subset of Q . Conversely, let R be a regularly closed subset of Q . Then $R = \delta\mu R$. From the definition of μR it is clear that $\sigma\mu R \subseteq R = \delta\mu R$ which shows that R is genetically closed.

COROLLARY 3.2. A primary of an automaton is precisely a minimal regular closed subset of Q^* .

The above observation, aided by Theorem 4.1 of Bavel [1], yields the following:

THEOREM 3.3. If $p \in Q$ then the following statements are equivalent:

- (a) δp is a primary of A ;
- (b) δp is a regular closed subset of Q^* ;
- (c) $\{p\}$ is not a nowhere dense subset of Q^* ;
- (d) δp is a minimal regular closed subset of Q^* ;
- (e) $p \in \mu\delta p$

REMARKS. (a) Since closures of open sets are regular closed, it is clear that if $R \subseteq Q$, then $\delta\sigma R$, as noted in Bavel [1], is genetically closed. (b) It is known that a finite union of regular closed sets stays regular closed. However, under the operation of intersection, this need not be true. Genetically closed sets, thus, do not form closed sets of topology. It may however be noted that genetic sets which are also closed subsets of Q^* constitute a topology on Q for which they are precisely all the closed sets.

4. COMPACT AUTOMATA.

We call an automaton $A = (Q, X, \delta)$ compact (Compact automata were called 'quasi-finite' automata or 'finitely reachable' automata in Shukla and Srivastava [3].) iff tA is a compact topological space (we do not assume a compact space to be necessarily Hausdorff). It was pointed out in Shukla and Srivastava [3] that Q is compact iff Q^* has a finite dense subset. This means that there is a finite subset D of states of Q which every other state of Q can be reached.

Clearly, like finite automata, compact automata also have only a finite number of blocks. In the remaining section we shall observe that finite automata and compact automata are alike in a few more respects.

THEOREM 4.1. A primary of a compact automaton is necessarily a maximal singly generated sub-automaton.

This follows from a result of Bavel ([1], Corollary 4.7) coupled with the fact that compact automata are finitely generated. Thus structure-wise, there is no distinction between the primaries of a finite automaton and of a compact automaton. This paves way for formualting a few theorems on (infinite) compact automata which are analogs of theorems on finite automata. As examples, we give the following two theorems [of Bavel [4]].

THEOREM 4.2. (Primary Decomposition Theorem) Let P_1, P_2, \dots, P_n be the set of primaries of a compact automaton (A, X, δ) then

- (i) $A = \bigcup_{i=1}^n P_i$
- (ii) for any $i, 1 \leq i \leq n, A \neq \bigcup_{i \neq j} P_i$

For the next theorem, we need a notation due to Bavel [4]. If $A_i = (Q_i, X, \delta)$ and $A_j = (Q_j, X, \delta)$ are two subautomata of an automaton $A = (Q, X, \delta)$ and if $f_i: Q_i \rightarrow Q$ and $f_j: Q_j \rightarrow Q$, two functions such that they agree on $Q_i \cap Q_j$ then $f_i \vee f_j$ is defined as

follows: $(f_i \vee f_j)(x)$ equals $f_i(x)$ if $x \in Q_i$ and it equals $f_j(x)$ if $x \in Q_j$.

THEOREM 4.3. (Homomorphism Decomposition Theorem). Let A be a compact automaton and B any other automaton. Let P_1, P_2, \dots, P_n be the primaries of A and $f: A \rightarrow B$ be a homomorphism. Then there exist homomorphisms $f_i: P_i \rightarrow B$, $1 \leq i \leq n$, such that f has a 'decomposition' $f = \bigvee_{i=1}^n f_i$. Furthermore, this decomposition is unique.

5. PRODUCTS OF AUTOMATA. Several definitions of products of automata are available in the literature. We consider here the categorical product of automata (in the category of automata and their homomorphisms) and observe that the state set topology functor from the category of automata to that of topological spaces preserves products. As a consequence certain observations made by Dörfler [2] are recovered.

The following definition of products of automata is easily extendible to any arbitrary family of automata. Let $A_1 = (Q_1, X_1, \delta_1)$ and $A_2 = (Q_2, X_2, \delta_2)$ be two automata. Then $A_1 \times A_2 = (Q_1 \times Q_2, X_1 \times X_2, \delta_1 \times \delta_2)$, where $X_1 \times X_2$ is the direct product of monoids X_1 and X_2 and $\delta_1 \times \delta_2$ is defined by $(\delta_1 \times \delta_2)((q_1, q_2), (x_1, x_2)) = (\delta_1(q_1, x_1), \delta_2(q_2, x_2))$ for any $(q_1, q_2) \in Q_1 \times Q_2$, and $(x_1, x_2) \in X_1 \times X_2$ is an automaton and will be called the product of the automata A_1 and A_2 . Of course this product is nothing but the usual parallel composition or the direct product in the sense of Dörfler [2]. It is easy to verify that it is also the categorical product.

THEOREM 5.1. Let $A_1 = (Q_1, X_1, \delta_1)$ and $A_2 = (Q_2, X_2, \delta_2)$ be two automata and $A_1 \times A_2$ their product. Then $tA_1 \times tA_2 = t(A_1 \times A_2)$, where $tA_1 \times tA_2$ denotes the Tychonoff product of the state set topologies tA_1 and tA_2 .

PROOF. First let U be open in $t(A_1 \times A_2)$. Then $(\delta_1 \times \delta_2)U = U$. Suppose $(q_1, q_2) \in U$. Consider $\delta q_1 \times \delta q_2$ which clearly is basic open in $tA_1 \times tA_2$. If $(q_1', q_2') \in \delta q_1 \times \delta q_2$ then there exist $x_1 \in X_1$ and $x_2 \in X_2$ such that $\delta_1(q_1, x_1) = q_1'$ and $\delta_2(q_2, x_2) = q_2'$. Since $(\delta_1 \times \delta_2)U = U$, it follows that $(\delta_1 \times \delta_2)((q_1, q_2), (x_1, x_2)) = (q_1', q_2') \in U$. Thus U contains a basic open set (viz, $\delta q_1 \times \delta q_2$) around its each point (q_1, q_2) and so must be open in $tA_1 \times tA_2$. Conversely, let U be open in $tA_1 \times tA_2$ and let $(q_1, q_2) \in U$. Then there must exist sets U_1 and U_2 open respectively in tA_1 and tA_2 with $(q_1, q_2) \in U_1 \times U_2 \subseteq U$. Since $\delta_1 U_1 = U_1$ and $\delta_2 U_2 = U_2$, it follows that $(\delta_1 \times \delta_2)(q_1, q_2) \in U$ whence $(\delta_1 \times \delta_2)U = U$ showing openness of U in $t(A_1 \times A_2)$.

It is known from topology that if T_1, T_2 are two simultaneously indiscrete, connected, R_0 or compact topologies then so is their Tychonoff product and also conversely. Thus in view of Theorem 1.1, we have the following.

COROLLARY 5.2. Let P be any of the properties connectedness, strong connectedness, retrievability or compactness of an automaton. Let A_1 and A_2 be two automata. Then $A_1 \times A_2$ has the property P iff both A_1 and A_2 have P .

The above observation in the case of P being connectedness or strong connectedness was made by Dörfler [2].

REFERENCES

1. BAVEL, Z. On the Decomposibility of Monadic Algebras and Automata, In. "Proceedings, Eighth Annual Symposium on Switching and Automata Theory, October 1967", (1967), 322-335.
2. DÖRFLER, W. The Direct Product of Automata and Quasiautomata, In. "Mathematical Foundation of Computer Science 1976", Lecture Notes in Computer Science No. 45, Springer-Verlag Berlin (1976), 270-276.
3. SHUKLA, W. and SRIVASTAVA, A.K. A Topology for Automata: A Note, *Inform. Contr.* **32**, (1976), 163-168.
4. BAVEL, Z. Structure and Transition-preserving Functions of Finite Automata, *J. Assoc. Comput. Mach.* **15** (1968), 135-158.