

# NULL EXACT CONTROLLABILITY OF THE PARABOLIC EQUATIONS WITH EQUIVALUED SURFACE BOUNDARY CONDITION

ZHONGQI YIN

Received 25 January 2005; Revised 21 April 2005; Accepted 26 April 2005

This paper is devoted to showing the null exact controllability for a class of parabolic equations with equivalued surface boundary condition. Our method is based on the duality argument and global Carleman-type estimate for a parabolic operator.

Copyright © 2006 Zhongqi Yin. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let  $T > 0$ ,  $\Omega \subset \mathbb{R}^n$  ( $n \in \mathbb{N}$ ) be a given bounded domain,  $\partial\Omega = \Gamma_0 \cup \Gamma_1$  ( $\Gamma_1 \neq \emptyset$ ) (where  $\Gamma_0$  is the interior boundary and  $\Gamma_1$  the outer boundary),  $\Gamma_0 \cap \Gamma_1 = \emptyset$ . For simplicity, we assume that  $\Gamma_0, \Gamma_1 \in C^\infty$  and  $\omega \neq \emptyset$  is a given subdomain of  $\Omega$ . Denote the characteristic function of  $\omega$  by  $\chi_\omega$ , and the unit outward normal vector of  $\Omega$  by  $(n_1, \dots, n_n)$ . Put  $Q = \Omega \times (0, T)$ ,  $Q^\omega = \omega \times (0, T)$ , and  $\Sigma = \partial\Omega \times (0, T)$ . Let  $a_{ij}(x) \in C^2(\bar{\Omega})$  satisfy  $a_{ij} = a_{ji}$ , and for some  $\Lambda > 0$ , it holds that

$$\sum_{i,j} a_{ij} \xi_i \xi_j \geq \Lambda |\xi|^2, \quad \forall (x, \xi) \in \Omega \times \mathbb{R}^n. \quad (1.1)$$

Here and henceforth, we denote  $\sum_{i,j=1}^n$  simply by  $\sum_{i,j}$ .

We consider the following controlled parabolic equation with equivalued surface boundary condition:

$$\begin{aligned} \frac{\partial y}{\partial t} - \sum_{i,j} \frac{\partial}{\partial x_i} \left( a_{ij}(x) \frac{\partial y}{\partial x_j} \right) &= \chi_\omega(x) b \quad \text{in } Q, \\ y|_{\Gamma_1} &= 0, \quad y|_{\Gamma_0} = m(t), \\ \int_{\Gamma_0} \frac{\partial y}{\partial n_A} ds &= 0, \quad y(x, 0) = y_0(x) \quad \text{in } \Omega, \end{aligned} \quad (1.2)$$

## 2 Null controllability of a nonlocal parabolic equation

where  $y = y(x, t)$  is the *state*,  $b = b(x, t)$  is the *control*,  $m(t) \in L^2(0, T)$  is an unknown function which depends only on the time variable  $t$ ,  $y_0$  is the initial state, and

$$\frac{\partial y}{\partial n_A} = \sum_{i,j} a_{ij}(x) \frac{\partial y}{\partial x_j} n_i. \quad (1.3)$$

In system (1.2), the *state space* is chosen as  $L^2(\Omega)$ , and the *control space* is  $L^2(\omega)$ . Let

$$Y \triangleq \left\{ y \in C([0, T]; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega)) \mid y|_{\Gamma_1} = 0, y|_{\Gamma_0} = m(\cdot) \in L^2(0, T) \right\}. \quad (1.4)$$

It can be shown that for any  $y_0 \in L^2(\Omega)$  and  $b \in L^2(\omega \times (0, T))$ , system (1.2) admits one and only one weak solution  $y \in Y$  (cf. [5, 6, 8]).

The null exact controllability problem of (1.2) is formulated as follows: for any given  $y_0 \in L^2(\Omega)$ , find a control  $b(x, t) \in L^2(\omega \times (0, T))$  (if possible) such that the weak solution  $y(\cdot) \in Y$  satisfies  $y(T) = 0$ .

There are many concrete physical backgrounds for problem (1.2), for example, the problem of resistivity well logging, the unstable temperature field around an underground electric cable, and so on (cf. [5, 6]).

In recent years, great progress has been made in the exact controllability problem of the linear and semilinear partial differential equations with Dirichlet or Neumann boundary condition, or other sorts of pointwise boundary value conditions ([1–4, 7, 9], and the references cited therein). However, to the author's best knowledge, there is no reference devoted to the same problem for the parabolic equations but with a spatial nonlocal boundary condition. In this paper, we will show the null exact controllability for system (1.2). By duality, the problem is reduced to the obtention of an observability inequality for the corresponding adjoint equation, which in turn is derived by means of a global Carleman-type estimate. Our method is stimulated by that in [4].

The rest of this paper is organized as follows. In Section 2, we state some preliminary results and our main results. The final section, Section 3, is devoted to the proof of our main theorem.

### 2. Main result

Throughout this paper,  $C$  denotes a positive constant depending only on  $\lambda, \mu, \Omega, T$ , and  $\omega$ , which may change from line to line.

To begin with, we fix  $\omega_0$  to be a nonempty open subset of  $\Omega$  such that  $\bar{\omega}_0 \subset \omega$ . Let  $\psi \in C^\infty(\bar{\Omega})$  satisfy  $\psi > 0$  in  $\Omega$ ,  $\psi = 0$  on  $\partial\Omega$ , and  $|\nabla\psi(x)| > 0$  for all  $x \in \Omega_0 = \Omega \setminus \omega_0$ . The existence of function  $\psi$  was proved in [7]. In this paper, we further assume that the following technical condition holds:

$$\left( \frac{\partial\psi}{\partial n} \right)^2 \sum_{i,j} a_{ij} n_i n_j |_{\Gamma_0} = \text{Const}. \quad (2.1)$$

This technical condition admits several interesting cases such as  $\Omega = \{x \in \mathbb{R}^n \mid r < |x| < R\}$

for some  $0 < r < R < \infty$ ,  $a_{ij}(x) = \delta_i^j$ , and  $\partial\psi/\partial n = \text{Const}$  on  $\Gamma_0$ . In this case, one may choose  $\psi(x) = (|x|^2 - r^2)(R^2 - |x|^2)$ .

The main result in this paper is stated as follows.

**THEOREM 2.1.** *Under the assumption (2.1), system (1.2) is null exact controllable.*

By means of the usual duality argument (see, e.g., [3, 9]), the proof of Theorem 2.1 is easily reduced to the obtention of an observability inequality for the following adjoint system of system (1.2):

$$\begin{aligned}
 -Lu \equiv -\frac{\partial u}{\partial t} - \sum_{i,j} \frac{\partial}{\partial x_i} \left( a_{ij}(x) \frac{\partial u}{\partial x_j} \right) &= 0 \quad \text{in } Q, \\
 u|_{\Gamma_1} &= 0, \quad u|_{\Gamma_0} = c(t), \\
 \int_{\Gamma_0} \frac{\partial u}{\partial n_A} ds &= 0, \quad u(x, T) = u(T) \quad \text{in } \Omega,
 \end{aligned} \tag{2.2}$$

where  $u(T) \in L^2(\Omega)$ , and similarly to system (1.2),  $c(\cdot) \in L^2(0, T)$  is an unknown function. More precisely, we need to show the following.

**THEOREM 2.2.** *Under the assumption (2.1), there is a constant  $C > 0$  such that solutions  $u \in Y$  of system (2.2) satisfy*

$$\|u(0)\|_{L^2(\Omega)} \leq C \|u(x, t)\|_{L^2(\omega \times (0, T))}. \tag{2.3}$$

*Remark 2.3.* It would be quite interesting to drop the technical condition (2.1). But this is by now an unsolved problem.

### 3. Proof of the main theorem

It suffices to prove Theorem 2.2. To this end, for any given parameters  $\lambda$  and  $\mu$ , we set

$$\begin{aligned}
 \alpha(x, t) &= (t(T - t))^{-1} (e^{\mu\psi(x)} - e^{2\mu\|\psi\|_{C(\bar{\Omega})}}), \\
 \varphi(x, t) &= (t(T - t))^{-1} e^{\mu\psi(x)}, \quad \theta(x, t) = e^{\lambda\alpha(x, t)}.
 \end{aligned} \tag{3.1}$$

Clearly, Theorem 2.2 is an easy consequence of the following global Carleman-type estimate for solutions of (2.2).

**THEOREM 3.1.** *Let (2.1) hold. Then there exist a constant  $\mu_1$  and a function  $\lambda_1 : \mathbb{R}^+ \rightarrow (1, \infty)$  such that for any  $\mu > \mu_1$  and  $\lambda > \lambda_1(\mu)$ , solutions  $u \in Y$  of system (2.2) satisfy*

$$\begin{aligned}
 \lambda^3 \int_Q \theta^2 \varphi^3 u^2 dx dt + \lambda \int_Q \theta^2 \varphi |\nabla u|^2 dx dt + \lambda^{-1} \int_Q \theta^2 \varphi^{-1} [u_t^2 + (\Delta u)^2] dx dt \\
 \leq C \lambda^3 \int_{Q^\omega} \theta^2 \varphi^3 u^2 dx dt.
 \end{aligned} \tag{3.2}$$

The rest of this section is devoted to prove Theorem 3.1. For this, we need the following pointwise estimate for parabolic operator  $Lu$ , which is a special case of [4, Lemma 3.1].

#### 4 Null controllability of a nonlocal parabolic equation

LEMMA 3.2. *Let  $u, \alpha \in C^2(\mathbb{R}^{n+1})$  and  $\lambda > 0$  be given. Put  $\theta = e^{\lambda\alpha}$  and  $v = \theta u$ . Then it holds that*

$$\begin{aligned}
\theta^2 |Lu|^2 \geq & \left[ -\lambda\alpha_t v^2 - \sum_{i,j} \left( a_{ij} v_{x_i} v_{x_j} + \lambda (a_{ij})_{x_j} \alpha_{x_i} v^2 + \lambda a_{ij} \alpha_{x_i x_j} v^2 - \lambda^2 a_{ij} \alpha_{x_i} \alpha_{x_j} v^2 \right) \right]_t \\
& + \sum_j \left[ 2 \sum_i \left( a_{ij} v_{x_i} v_t + \lambda^2 a_{ij} \alpha_{x_i} \alpha_t v^2 \right) \right. \\
& \quad + 2\lambda \sum_i \left( a_{ij} a_{\ell m} \alpha_{x_i} v_{x_\ell} v_{x_m} + a_{ij} (a_{\ell m} \alpha_{x_\ell x_m})_{x_i} v^2 - 2a_{ij} a_{\ell m} \alpha_{x_\ell} v_{x_i} v_{x_m} \right. \\
& \quad \quad - 2a_{ij} a_{\ell m} \alpha_{x_\ell x_m} v v_{x_i} + \lambda a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_\ell x_m} v^2 \\
& \quad \quad \left. \left. + \lambda a_{ij} (a_{\ell m})_{x_m} \alpha_{x_\ell} \alpha_{x_m} v^2 - \lambda^2 a_{ij} a_{\ell m} \alpha_{x_\ell} \alpha_{x_m} v^2 \right) \right]_{x_j} \\
& + \left\{ \lambda \alpha_{tt} - \lambda \sum_{i,j} \left[ ((a_{ij})_{x_j} \alpha_{x_i})_t + (a_{ij} \alpha_{x_i x_j})_t - \lambda (\alpha_{ij} \alpha_{x_i} \alpha_{x_j})_t \right. \right. \\
& \quad \quad \left. - 2\lambda (a_{ij} \alpha_{x_i} \alpha_t)_{x_j} + 4\lambda a_{ij} \alpha_{x_i x_j} \alpha_t \right] \\
& \quad + 2\lambda \sum_{i,j,\ell,m} \left[ 2\lambda (a_{ij})_{x_j} a_{\ell m} \alpha_{x_i} \alpha_{x_\ell x_m} - (a_{ij} (a_{\ell m} \alpha_{x_\ell x_m})_{x_j})_{x_i} - \lambda ((a_{\ell m})_{x_m} a_{ij} \alpha_{x_i} \alpha_{x_\ell})_{x_j} \right. \\
& \quad \quad \left. - \lambda (a_{ij} a_{\ell m} \alpha_{x_i x_j} \alpha_{x_\ell})_{x_m} - 2\lambda a_{ij} a_{\ell m} \alpha_{x_i x_j} \alpha_{x_\ell x_m} \right. \\
& \quad \quad \left. \left. + \lambda^2 (a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_j} \alpha_{x_\ell})_{x_m} - 2\lambda^2 a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_j} \alpha_{x_\ell x_m} \right] \right\} v^2 \\
& + 2\lambda \sum_{i,j,\ell,m} \left[ 2a_{ij} (a_{\ell m} \alpha_{x_\ell})_{x_j} v_{x_i} v_{x_m} - (a_{ij} a_{\ell m})_{x_m} \alpha_{x_\ell} v_{x_i} v_{x_j} + a_{ij} a_{\ell m} \alpha_{x_\ell x_m} v_{x_i} v_{x_j} \right].
\end{aligned} \tag{3.3}$$

*Proof of Theorem 3.1.* The main idea of our proof is to use the pointwise estimate in Lemma 3.2. The proof is divided into several steps.

*Step 1.* Recall that  $\theta = e^{\lambda\alpha}$ ,  $v = \theta u$ . We claim that

$$\begin{aligned}
& \int_Q \theta^2 (Lu)^2 dx dt - \int_Q \operatorname{div} V dx dt + C\lambda^3 \mu^3 \int_Q \varphi^3 v^2 dx dt + C\lambda \mu^2 \int_Q \varphi |\nabla v|^2 dx dt \\
& \geq 2\Lambda^2 \lambda^3 \mu^4 \int_Q \varphi^3 |\nabla \psi|^4 v^2 dx dt + 2\Lambda^2 \lambda \mu^2 \int_Q \varphi |\nabla v|^2 |\nabla \psi|^2 dx dt,
\end{aligned} \tag{3.4}$$

where  $V = (V_1, V_2, \dots, V_n)$ , and

$$\begin{aligned}
V_j = & -2 \sum_i \left( a_{ij} v_{x_i} v_t + \lambda^2 a_{ij} \alpha_{x_i} \alpha_t v^2 \right) \\
& + 2\lambda \sum_{i,\ell,m} \left( a_{ij} a_{\ell m} \alpha_{x_i} v_{x_\ell} v_{x_m} - 2a_{ij} a_{\ell m} \alpha_{x_\ell} v_{x_i} v_{x_m} + \lambda a_{ij} (a_{\ell m})_{x_m} \alpha_{x_i} \alpha_{x_m} v^2 \right. \\
& \quad \left. - \lambda^2 a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_\ell} \alpha_{x_m} v^2 + \lambda a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_\ell x_m} v^2 \right. \\
& \quad \left. - 2a_{ij} a_{\ell m} \alpha_{x_\ell x_m} v v_{x_i} + a_{ij} (a_{\ell m} \alpha_{x_\ell x_m})_{x_i} v^2 \right).
\end{aligned} \tag{3.5}$$

By the definition of  $\alpha$  and  $\varphi$  in (3.1), it is easy to see that

$$\int_Q \left[ -\lambda \alpha_t v^2 + \sum_{i,j} (a_{ij} v_{x_i} v_{x_j} + \lambda (a_{ij})_{x_j} \alpha_{x_i} v^2 - \lambda^2 a_{ij} \alpha_{x_i} \alpha_{x_j}^2 + \lambda a_{ij} \alpha_{x_i x_j} v^2) \right] dt = 0. \quad (3.6)$$

Let us estimate the last three “energy” terms of first order in the right-hand side of (3.3). First,

$$\begin{aligned} & \left| \int_Q 4\lambda \sum_{i,j,\ell,m} a_{ij} (a_{\ell m} \alpha_{x_\ell})_{x_j} v_{x_i} v_{x_m} dx dt \right| \\ &= \left| \int_Q 4\lambda \sum_{i,j,\ell,m} (a_{ij} (a_{\ell m})_{x_j} \mu \psi_{x_\ell} \varphi v_{x_i} v_{x_m} + a_{ij} a_{\ell m} (\mu \psi_{x_\ell x_j} \varphi + \mu^2 \psi_{x_\ell} \psi_{x_j} \varphi) v_{x_i} v_{x_m}) dx dt \right| \\ &\leq C\lambda \mu \int_Q \varphi |\nabla v|^2 dx dt + C\lambda \mu^2 \int_Q \varphi |\nabla \psi|^2 dx dt, \end{aligned} \quad (3.7)$$

where  $C$  is a positive constant for  $\lambda$  large enough.

Next, it is easy to see that

$$\begin{aligned} & \left| -\int_Q 2\lambda \sum_{i,j,\ell,m} (a_{ij} a_{\ell m})_{x_m} \alpha_{x_\ell} v_{x_i} v_{x_j} dx dt \right| \leq C\lambda \mu \int_Q \varphi |\nabla v|^2 dx dt, \\ & \int_Q 2\lambda \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \alpha_{x_\ell x_m} v_{x_i} v_{x_j} dx dt \\ &= \int_Q 2\lambda \sum_{i,j,\ell,m} (\mu a_{ij} a_{\ell m} \psi_{x_\ell x_m} \varphi v_{x_i} v_{x_j} + \mu^2 a_{ij} a_{\ell m} \psi_{x_\ell} \psi_{x_m} \varphi v_{x_i} v_{x_j}) dx dt \\ &\geq -C\lambda \mu \int_Q \varphi |\nabla v|^2 dx dt + 2\lambda \mu^2 \Lambda^2 \int_Q \varphi |\nabla \psi|^2 |\nabla v|^2 dx dt. \end{aligned} \quad (3.8)$$

It remains to deal with the “energy” term of zero order, that is,  $\int_Q \{\dots\} v^2 dx dt$ , in the right-hand side of (3.3). Similarly to [4], we have

$$\begin{aligned} & -2 \int_{\underline{Q}} \lambda^3 \sum_{i,j,\ell,m} (2a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_j} \alpha_{x_\ell x_m} v^2 - (a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_j} \alpha_{x_\ell})_{x_m} v^2) dx dt \\ &= -\int_{\underline{Q}} 2\lambda^3 \mu^3 \sum_{i,j,\ell,m} (2a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_j} \psi_{x_\ell x_m} \varphi^3 v^2 + 2\mu a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_j} \psi_{x_\ell} \psi_{x_m} \varphi^3 v^2 \\ &\quad - (a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_j} \psi_{x_\ell})_{x_m} \varphi^3 v^2 - 3\mu a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_j} \psi_{x_\ell} \psi_{x_m} \varphi^3 v^2) dx dt \\ &\geq -C\lambda^3 \mu^3 \int_{\underline{Q}} \varphi^3 v^2 dx dt + 2\lambda^3 \mu^4 \Lambda^2 \int_{\underline{Q}} \varphi^3 |\nabla \psi|^4 v^2 dx dt. \end{aligned} \quad (3.9)$$

## 6 Null controllability of a nonlocal parabolic equation

Therefore, for large  $\lambda$ , the following estimate holds:

$$\begin{aligned}
& \int_Q \left\{ \lambda \alpha_{tt} - \lambda \sum_{i,j} [((a_{ij})_{x_j} \alpha_{x_i})_t - \lambda (a_{ij} \alpha_{x_i} \alpha_{x_j})_t + (a_{ij} \alpha_{x_i x_j})_t - 2\lambda (a_{ij} \alpha_{x_i} \alpha_t)_{x_j} + 4\lambda a_{ij} \alpha_{x_i x_j} \alpha_t] \right. \\
& \quad + 2\lambda \sum_{i,j,\ell,m} \left[ \lambda^2 (a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_j} \alpha_{x_\ell})_{x_m} - \lambda (a_{ij} a_{\ell m} \alpha_{x_i x_j} \alpha_{x_\ell})_{x_m} - \lambda ((a_{\ell m})_{x_m} a_{ij} \alpha_{x_i} \alpha_{x_j})_{x_j} \right. \\
& \quad \quad + 2\lambda (a_{ij})_{x_j} a_{\ell m} \alpha_{x_i} \alpha_{x_\ell x_m} - 2\lambda^2 a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_j} \alpha_{x_\ell x_m} \\
& \quad \quad \left. \left. - 2\lambda a_{ij} a_{\ell m} \alpha_{x_i x_j} \alpha_{x_\ell x_m} - (a_{ij} (a_{\ell m} \alpha_{x_\ell x_m})_{x_j})_{x_i} \right] \right\} v^2 dx dt \\
& \geq -C\lambda^3 \mu^3 \int_Q \varphi^3 v^2 dx dt + 2\lambda^3 \mu^4 \Lambda^2 \int_Q \varphi^3 |\nabla \psi|^4 v^2 dx dt.
\end{aligned} \tag{3.10}$$

*Step 2.* From (3.4), one finds

$$\begin{aligned}
& C \left( \int_Q \theta^2 (Lu)^2 dx dt - \int_Q \operatorname{div} V dx dt + \lambda^3 \mu^3 \int_Q \varphi^3 v^2 dx dt + \lambda \mu^2 \int_Q \varphi |\nabla v|^2 dx dt \right) \\
& \geq \lambda^3 \mu^4 \int_Q \varphi^3 |\nabla \psi|^4 v^2 dx dt + \lambda \mu^2 \int_Q \varphi |\nabla \psi|^2 |\nabla v|^2 dx dt.
\end{aligned} \tag{3.11}$$

Set

$$Q^{\omega_0} = \omega_0 \times (0, T). \tag{3.12}$$

Noting that  $|\nabla \psi(x)| > 0$  for all  $x \in \Omega_0 = \bar{\Omega} \setminus \omega_0$ , by (3.11), it is easy to see that

$$\begin{aligned}
& C \left( \int_Q \theta^2 (Lu)^2 dx dt - \int_Q \operatorname{div} V dx dt + \lambda^3 \mu^3 \int_{Q^{\omega_0}} \varphi^3 v^2 dx dt + \lambda \mu^2 \int_{Q^{\omega_0}} \varphi |\nabla v|^2 dx dt \right) \\
& \geq \lambda^3 \mu^4 \int_Q \varphi^3 v^2 dx dt + \lambda \mu^2 \int_Q \varphi |\nabla v|^2 dx dt.
\end{aligned} \tag{3.13}$$

Returning  $v$  to  $e^{\lambda \alpha} u$  in (3.13), we arrive at

$$\begin{aligned}
& C \left( \int_Q \theta^2 (Lu)^2 dx dt - \int_Q \operatorname{div} V dx dt + \lambda^3 \int_{Q^{\omega_0}} \theta^2 \varphi^3 u^2 dx dt + \lambda \int_{Q^{\omega_0}} \theta^2 \varphi |\nabla u|^2 dx dt \right) \\
& \geq \lambda^3 \int_Q \theta^2 \varphi^3 u^2 dx dt + \lambda \int_Q \theta^2 \varphi |\nabla u|^2 dx dt.
\end{aligned} \tag{3.14}$$

*Step 3.* The purpose of this step is to get rid of the fourth term in the left-hand side of (3.14). To this end, we multiply  $Lu$  by  $\tilde{\chi} \theta^2 \varphi u$  and then integrate it over  $Q$ , where  $\tilde{\chi} \in C_0^\infty(\omega)$ ,  $\tilde{\chi} = 1$  in  $\omega_0$ , and  $\tilde{\chi} = 0$  in  $\Omega \setminus \omega$ . Then, we obtain

$$\int_Q \frac{\partial u}{\partial t} \tilde{\chi} \theta^2 \varphi u dx dt + \int_Q \sum_{i,j} \frac{\partial}{\partial x_i} \left( a_{ij}(x) \frac{\partial u}{\partial x_j} \right) \tilde{\chi} \theta^2 \varphi u dx dt = \int_Q (Lu) \tilde{\chi} \theta^2 \varphi u dx dt. \tag{3.15}$$

Using integration by parts, it is easy to deduce from (3.15) that

$$\lambda \int_{Q^{\omega_0}} \theta^2 \varphi |\nabla u|^2 dx dt \leq C \left( \int_Q \theta^2 (Lu)^2 dx dt + \lambda^3 \int_{Q^\omega} \theta^2 \varphi^3 u^2 dx dt \right). \quad (3.16)$$

Combining (3.13) and (3.16), we end up with

$$\begin{aligned} & C \left( \int_Q \theta^2 (Lu)^2 dx dt + \lambda^3 \int_{Q^\omega} \theta^2 \varphi^3 u^2 dx dt - \int_Q \operatorname{div} V dx dt \right) \\ & \geq \lambda^3 \int_Q \theta^2 \varphi^3 u^2 dx dt + \lambda \int_Q \theta^2 \varphi |\nabla u|^2 dx dt. \end{aligned} \quad (3.17)$$

*Step 4.* This step is to estimate the “divergence” term  $\operatorname{div} V$ . Denote the terms on the right-hand side of (3.5) by  $I_i$ ,  $i = 1, 2, \dots, 9$ . First,

$$\begin{aligned} I_1 + I_2 &= \int_Q \sum_j \left( 2 \sum_i (a_{ij} v_{x_i} v_t + \lambda^2 a_{ij} \alpha_{x_i} \alpha_t v^2) \right)_{x_j} \\ &= \lambda^2 \mu \int_0^T \theta^2 \varphi_t c^2(t) dt \int_{\Gamma_0} \sum_{i,j} a_{ij} \psi_{x_i} n_j ds - 2\lambda^2 \mu \int_0^T \theta^2 \varphi \alpha_t c^2(t) dt \int_{\Gamma_0} \sum_{i,j} a_{ij} \psi_{x_i} n_j ds. \end{aligned} \quad (3.18)$$

Next,

$$\begin{aligned} I_3 + I_5 + I_9 &= \int_Q \sum_j \left( 2\lambda \sum_{i,\ell,m} (a_{ij} a_{\ell m} \alpha_{x_i} v_{x_\ell} v_{x_m} - 2a_{ij} a_{\ell m} \alpha_{x_\ell} v_{x_i} v_{x_m} - \lambda^2 a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_m} v^2) \right)_{x_j} dx dt \\ &= -4\lambda^3 \mu^3 \int_0^T \varphi^3 c^2(t) \theta^2 dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_\ell} \psi_{x_m} n_j ds \\ &\quad - 4\lambda^2 \mu^2 \int_0^T \varphi^2 \theta^2 c(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_\ell} \psi_{x_i} u_{x_m} n_j ds \\ &\quad - 2 \int_0^T \lambda \mu \varphi \theta^2 dt \int_{\Gamma} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_\ell} u_{x_i} u_{x_m} n_j ds. \end{aligned} \quad (3.19)$$

Further,

$$\begin{aligned} I_4 &= \int_Q \sum_j \left( 2\lambda \sum_i a_{ij} (a_{\ell m} \alpha_{x_\ell x_m})_{x_i} v^2 \right)_{x_j} dx dt \\ &= 2\lambda \mu^2 \int_0^T \theta^2 \varphi c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} (a_{\ell m})_{x_i} \psi_{x_\ell} \psi_{x_m} n_j ds \\ &\quad + 2\lambda \mu \int_0^T \theta^2 \varphi c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} (a_{\ell m})_{x_i} \psi_{x_\ell x_m} n_j ds \\ &\quad + 2\lambda \mu^3 \int_0^T \theta^2 \varphi c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_\ell} \psi_{x_m} n_j ds \end{aligned}$$

## 8 Null controllability of a nonlocal parabolic equation

$$\begin{aligned}
& + 2\lambda\mu^2 \int_0^T \theta^2 \varphi c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i x_\ell} \psi_{x_m} n_j ds \\
& + 2\lambda\mu^2 \int_0^T \theta^2 \varphi c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_\ell} \psi_{x_i x_m} n_j ds \\
& + 2\lambda\mu \int_0^T \theta^2 \varphi c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i x_\ell x_\ell} n_j ds \\
& \geq C\lambda\mu^3 \int_0^T \theta^2 \varphi c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_\ell} \psi_{x_m} n_j ds.
\end{aligned} \tag{3.20}$$

Further,

$$\begin{aligned}
I_6 & = - \int_Q \sum_j \left( 4\lambda \sum_{i,\ell,m} a_{ij} a_{\ell m} \alpha_{x_\ell x_m} \nu \nu_{x_i} \right)_{x_j} dx dt \\
& = -4\lambda^2 \mu^3 \int_0^T \theta^2 \varphi^2 c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_\ell} \psi_{x_m} n_j ds \\
& \quad - 4\lambda\mu^2 \int_0^T \theta^2 \varphi^2 c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_\ell x_m} n_j ds \\
& \quad - 4\lambda\mu^2 \int_0^T \theta^2 \varphi c(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_\ell} \psi_{x_m} u_{x_i} n_j ds \\
& \quad - 4\lambda\mu \int_0^T \theta^2 \varphi c(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_\ell x_m} u_{x_i} n_j ds.
\end{aligned} \tag{3.21}$$

Finally,

$$\begin{aligned}
I_7 + I_8 & = \int_Q \sum_j \left( 2\lambda^2 \sum_{i,\ell,m} (a_{ij} a_{\ell m} \alpha_{x_i} \alpha_{x_\ell x_m} v^2 + a_{ij} (a_{\ell m})_{x_m} \alpha_{x_i} \alpha_{x_m} v^2) \right)_{x_j} dx dt \\
& = 2\lambda^2 \mu^3 \int_0^T \theta^2 \varphi^2 c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_\ell} \psi_{x_m} n_j ds \\
& \quad + 2\lambda^2 \mu^2 \int_0^T \theta^2 \varphi^2 c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} a_{\ell m} \psi_{x_i} \psi_{x_\ell x_m} n_j ds \\
& \quad + 2\lambda^2 \mu^2 \int_0^T \theta^2 \varphi^2 c^2(t) dt \int_{\Gamma_0} \sum_{i,j,\ell,m} a_{ij} (a_{\ell m})_{x_m} \psi_{x_i} \psi_{x_m} n_j ds.
\end{aligned} \tag{3.22}$$

Now, combining (3.18)–(3.22) and using the technical condition (2.1), we conclude that, for large  $\lambda$  and  $\mu$ , it holds that

$$\int_Q \operatorname{div} V dx dt \geq C\lambda^3 \mu^3 \int_0^T \theta^2 \varphi^3 c^2(t) dt - 4\lambda\mu\Lambda \int_0^T \theta^2 \varphi dt \int_\Gamma |\nabla u|^2 \sum_{i,j} a_{ij} \psi_{x_i} n_j ds. \tag{3.23}$$



Step 5. It remains to estimate  $\int_Q (\lambda\varphi)^{-1}\theta^2(u_t^2 + \sum_{i,j}(a_{ij}u_{x_i})_{x_j})^2 dx dt$ . For this, we observe that

$$\begin{aligned} \left( v_t + \sum_{i,j} (a_{ij}v_{x_i})_{x_j} \right)^2 &\leq C \left[ \lambda^2 \alpha_t^2 v^2 + \sum_{i,j} (\lambda^2 (a_{ij})_{x_j}^2 \alpha_{x_i}^2 v^2 + \lambda^2 a_{ij}^2 \alpha_{x_i x_j}^2 v^2 \right. \\ &\quad \left. + \lambda^4 a_{ij}^2 \alpha_{x_i}^2 \alpha_{x_j}^2 v^2 + \lambda^2 a_{ij}^2 \alpha_{ij}^2 \alpha_{x_i}^2 v_{x_j}^2 + (a_{ij})_{x_j}^2 v_{x_i}^2 \right]. \end{aligned} \quad (3.24)$$

This implies

$$\begin{aligned} (\lambda\varphi)^{-1} \left( v_t + \sum_{i,j} (a_{ij}v_{x_i})_{x_j} \right)^2 &\leq C(\lambda\varphi)^{-1} \left[ \lambda^2 \alpha_t^2 v^2 + \sum_{i,j} (\lambda^2 (a_{ij})_{x_j}^2 \alpha_{x_i}^2 v^2 + \lambda^2 a_{ij}^2 \alpha_{x_i x_j}^2 v^2 \right. \\ &\quad \left. + \lambda^4 a_{ij}^2 \alpha_{x_i}^2 \alpha_{x_j}^2 v^2 + \lambda^2 a_{ij}^2 \alpha_{ij}^2 \alpha_{x_i}^2 v_{x_j}^2 + (a_{ij})_{x_j}^2 v_{x_i}^2 \right] \\ &\leq C(\lambda^3 \mu^4 \varphi^3 v^2 + \lambda\mu\varphi|\nabla v|^2). \end{aligned} \quad (3.25)$$

Noting that

$$\begin{aligned} 2 \int_Q \sum_{i,j} (\lambda\varphi)^{-1} v_t (a_{ij}v_{x_i})_{x_j} dx dt &= 2 \sum_{i,j} \left( \int_Q (a_{ij}(\lambda\varphi)^{-1} v_{x_i} v_t)_{x_j} dx dt - 2 \int_Q a_{ij}(\lambda\varphi)^{-1} v_{x_i} v_t dx dt + \int_Q a_{ij}(\lambda\varphi)^{-1} v_{x_i} v_{x_j} dx dt \right. \\ &\quad \left. + \int_Q (a_{ij})_t (\lambda\varphi)^{-1} v_{x_i} v_{x_j} dx dt + \int_Q a_{ij} ((\lambda\varphi)^{-1} v_t)_{x_i} v_{x_j} dx dt \right), \end{aligned} \quad (3.26)$$

we get

$$\begin{aligned} 2 \int_Q (\lambda\varphi)^{-1} \sum_{i,j} (a_{ij}v_{x_i})_{x_j} v_t dx dt &\leq \frac{1}{2} \int_Q (\lambda\varphi)^{-1} v_t^2 dx dt + C\lambda \int_Q \varphi |\nabla v|^2 dx dt \\ &\quad + \lambda\mu \int_0^T \theta \alpha_t c^2(t) dt \int_{\Gamma_0} \sum_{i,j} a_{ij} \psi_{x_i} n_j ds. \end{aligned} \quad (3.27)$$

By (3.24)–(3.27), we obtain that

$$\begin{aligned} \int_Q (\lambda\varphi)^{-1} [v_t^2 + (v_{x_i x_j})^2] dx dt &\leq C \left( \int_Q (Lu)^2 \theta^2 dx dt + \lambda^3 \int_{Q^{\omega_0}} \mu^3 \varphi^3 v^2 dx dt + \lambda \int_{Q^{\omega_0}} \mu\varphi |\nabla v|^2 dx dt \right) \\ &\quad + \lambda\mu \int_0^T \theta \alpha_t c^2(t) dt \int_{\Gamma_0} \sum_{i,j} a_{ij} \psi_{x_i} n_j ds. \end{aligned} \quad (3.28)$$

This gives

$$\begin{aligned} & \int_Q (\lambda\varphi)^{-1} [v_t^2 + (\Delta v)^2] dx dt \\ & \leq C \left( \int_Q (Lu)^2 \theta^2 dx dt + \lambda^3 \int_{Q^{w_0}} \mu^3 \varphi^3 v^2 dx dt + \lambda \int_{Q^{w_0}} \mu \varphi |\nabla v|^2 dx dt \right) \\ & \quad + \lambda \mu \int_0^T \theta \alpha_t c^2(t) dt \int_{\Gamma_0} \sum_{i,j} a_{ij} \psi_{x_i} n_j ds. \end{aligned} \quad (3.29)$$

It is easy to see that the last term of the above inequality can be absorbed by (3.23) for large  $\lambda$  and  $\mu$ . Finally, replacing  $v$  by  $e^{\lambda\alpha}u$  and combining (2.2), (3.16), (3.17), (3.23), and (3.29), we get the desired estimate (3.2). This completes the proof of Theorem 3.1.  $\square$

### Acknowledgments

This work is partially supported by FANEDD of China (Project No 200119), NCET of China under Grant NCET-04-0882, and the NSF of China under Grants 10371084 and 10525105. The author gratefully acknowledges Professor Xu Zhang for his guidance, encouragement, and suggestions.

### References

- [1] V. Barbu, *Exact controllability of the superlinear heat equation*, Applied Mathematics and Optimization. An International Journal with Applications to Stochastics **42** (2000), no. 1, 73–89.
- [2] A. Doubova, E. Fernández-Cara, M. González-Burgos, and E. Zuazua, *On the controllability of parabolic systems with a nonlinear term involving the state and the gradient*, SIAM Journal on Control and Optimization **41** (2002), no. 3, 798–819.
- [3] A. V. Fursikov and O. Yu. Imanuvilov, *Controllability of Evolution Equations*, Lecture Notes Series, vol. 34, Seoul National University, Research Institute of Mathematics, Global Analysis Research Center, Seoul, 1996.
- [4] W. Li, *Observability estimate for the ultra-parabolic equations*, Master's thesis, School of Mathematics, Sichuan University, Chengdu, 2004.
- [5] T. Li, S. Zheng, Y. Tan, and W. Shen, *Boundary Value Problems with Equivalued Surface and Resistivity Well-Logging*, Pitman Research Notes in Mathematics Series, vol. 382, Longman, Harlow, 1998.
- [6] W. Shen, *Initial-boundary problem for the second order parabolic equations with equivalued surface boundary condition*, Journal of Fudan University Natural Science **4** (1978), 15–23 (Chinese).
- [7] G. Wang and L. Wang, *The Carleman inequality and its application to periodic optimal control governed by semilinear parabolic differential equations*, Journal of Optimization Theory and Applications **118** (2003), no. 2, 429–461.
- [8] X. Zhang, *Solvability of non-linear parabolic boundary value problem with equivalued surface*, Mathematical Methods in the Applied Sciences **22** (1999), no. 3, 259–265.
- [9] ———, *Exact Controllability of Semi-linear Distributed Parameter Systems*, Chinese Higher Education Press, Beijing, 2004.

Zhongqi Yin: Department of Engineering and Technology of Caotang School,  
Sichuan Normal University, Chengdu 610072, China  
E-mail address: zhongqiyin@sohu.com