

Research Article

Precise Asymptotics in the Law of Iterated Logarithm for Moving Average Process under Dependence

Jie Li^{1,2}

¹ Wang Yanan Institute for Studies in Economics, Xiamen University, Xiamen 361005, China

² School of Mathematics and Statistics, Zhejiang University of Finance and Economics, Hangzhou 310018, China

Correspondence should be addressed to Jie Li, lijiezufe@gmail.com

Received 10 November 2010; Revised 2 February 2011; Accepted 3 March 2011

Academic Editor: Ulrich Abel

Copyright © 2011 Jie Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $\{\xi_i, -\infty < i < \infty\}$ be a doubly infinite sequence of identically distributed and ϕ -mixing random variables, and let $\{a_i, -\infty < i < \infty\}$ be an absolutely summable sequence of real numbers. In this paper, we get precise asymptotics in the law of the logarithm for linear process $\{X_k = \sum_{i=-\infty}^{+\infty} a_{i+k} \xi_i, k \geq 1\}$, which extend Liu and Lin's (2006) result to moving average process under dependence assumption.

1. Introduction and Main Results

Let $\{\xi_i, -\infty < i < \infty\}$ be a doubly infinite sequence of identically distributed random variables with zero means and finite variances, and let $\{a_i, -\infty < i < \infty\}$ be an absolutely summable sequence of real numbers. Let

$$X_k = \sum_{i=-\infty}^{+\infty} a_{i+k} \xi_i, \quad k \geq 1, \tag{1.1}$$

be the moving average process based on $\{\xi_i, -\infty < i < \infty\}$. As usual, we denote $S_n = \sum_{k=1}^n X_k$, $n \geq 1$ as the sequence of partial sums.

Under the assumption that $\{\xi_i, -\infty < i < \infty\}$ is a sequence of independent identically distributed random variables, many limiting results have been obtained. Ibragimov [1] established the central limit theorem; Burton and Dehling [2] obtained a large deviation principle; Yang [3] established the central limit theorem and the law of the iterated logarithm;

Li et al. [4] obtained the complete convergence result for $\{X_k, k \geq 1\}$. As we know, X_k ($k \geq 1$) are dependent even if $\{\xi_i, -\infty < i < \infty\}$ is a sequence of i.i.d. random variables. Therefore, we introduce the definition of ϕ -mixing,

$$\phi(m) := \sup_{k \geq 1} \left\{ |P(B | A) - P(B)|, A \in \mathcal{F}_{-\infty}^k, P(A) \neq 0, B \in \mathcal{F}_{k+m}^\infty \right\} \rightarrow 0, \quad m \rightarrow \infty, \quad (1.2)$$

where $\mathcal{F}_a^b = \sigma(\xi_i, a \leq i \leq b)$. Many limiting results of moving average for ϕ -mixing have been obtained. For example, Zhang [5] got complete convergence.

Theorem A. Suppose that $\{\xi_i, -\infty < i < \infty\}$ is a sequence of identically distributed and ϕ -mixing random variables with $\sum_{m=1}^{\infty} \phi^{1/2}(m) < \infty$, and $\{X_k, k \geq 1\}$ is defined as (1.1). Let $h(x) > 0$ ($x > 0$) be a slowly varying function and $1 \leq t < 2, r \geq 1$, then $E\xi_1 = 0$ and $E|\xi_1|^{rt}h(|Y_t^t|) < \infty$ imply

$$\sum_{n=1}^{\infty} n^{r-2} h(n) P(|S_n| \geq n^{1/t} \epsilon) < \infty, \quad \forall \epsilon > 0. \quad (1.3)$$

Li and Zhang [6] achieved precise asymptotics in the law of the iterated logarithm.

Theorem B. Suppose that $\{\xi_i, -\infty < i < \infty\}$ is a sequence of identically distributed and ϕ -mixing random variables with mean zeros and finite variances, $\sum_{m=1}^{\infty} \phi^{1/2}(m) < \infty$, and $0 < \sigma^2 = E\xi_1^2 + 2 \sum_{k=2}^{\infty} E\xi_1 \xi_k < \infty$, $E\xi_1^2 (\log^+ |\xi_1|)^{\delta-1} < \infty$, for $\delta > 0$. Suppose that $\{X, X_k, k \geq 1\}$ is defined as in (1.1), where $\{a_i, -\infty < i < \infty\}$ is a sequence of real number with $\sum_{i=-\infty}^{\infty} |a_i| < \infty$, then one has

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta+2} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta}}{n} P(|S_n| \geq \sqrt{n \log n} \tau \epsilon) = \frac{\tau^{2\delta+2}}{\delta+1} E|N|^{2\delta+2}, \quad (1.4)$$

where $\tau =: \sigma \sum_{i=-\infty}^{\infty} a_i$, N is a standard normal random variable.

On the other hand, since Hsu and Robbins [7] introduced the concept of the complete convergence, there have been extensions in some directions. For the case of i.i.d. random variables, Davis [8] proved $\sum_{n=1}^{\infty} (\log n/n) P(|S_n| \geq \sqrt{n \log n} \epsilon) < \infty$, for $\epsilon > 0$ if and only if $EX_1 = 0$, $EX_1^2 < \infty$. Gut and Spătaru [9] gave the precise asymptotics of $\sum_{n=2}^{\infty} ((\log n)^{\delta}/n) P(|S_n| \geq \sqrt{n \log n} \epsilon)$. We know that complete convergence can be derived from complete moment convergence. Liu and Lin [10] introduced a new kind of convergence of $\sum_{n=2}^{\infty} ((\log n)^{\delta-1}/n^2) E|S_n|^2 I\{|S_n| \geq \sqrt{n \log n} \epsilon\}$. In this note, we show that the precise asymptotics for the moment convergence hold for moving-average process when $\{\xi_i, -\infty < i < \infty\}$ is a strictly stationary ϕ -mixing sequences. Now, we state the main results.

Theorem 1.1. Suppose that $\{X, X_k, k \geq 1\}$ is defined as in (1.1), where $\{a_i, -\infty < i < \infty\}$ is a sequence of real number with $\sum_{i=-\infty}^{\infty} |a_i| < \infty$, and $\{\xi_i, -\infty < i < \infty\}$ is a sequence of identically

distributed ϕ -mixing random variables with mean zeros and finite variances, $\sum_{m=1}^{\infty} \phi^{1/2}(m) < \infty$ and $0 < \sigma^2 = E\xi_1^2 + 2 \sum_{k=2}^n E\xi_1\xi_k < \infty$, $E\xi_1^2(\log^+|\xi_1|)^{\delta} < \infty$, for $0 < \delta \leq 1$, then one has

$$\lim_{\epsilon \searrow 0} e^{2\delta} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-1}}{n^2} E|S_n|^2 I\left\{ |S_n| \geq \sqrt{n \log n} \tau \epsilon \right\} = \frac{\tau^{2\delta+2}}{\delta} E|N|^{2\delta+2}, \quad (1.5)$$

where $\tau =: \sigma \sum_{i=-\infty}^{\infty} a_i$.

Theorem 1.2. Under the conditions in Theorem 1.1, one has

$$\lim_{\epsilon \searrow 0} e^{2\delta} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-1}}{n^2 \log n} E S_n^2 I\left\{ |S_n| \geq \sqrt{n \log \log n} \tau \epsilon \right\} = \frac{\tau^{2\delta+2} E|N|^{2(\delta+1)}}{\delta}. \quad (1.6)$$

Remark 1.3. In this paper, we generate the results of Liu and Lin [10] to linear process under dependence based on Theorem B by using the technique of dealing with the innovation process in Zhang [5].

We first proceed with some useful lemmas.

Lemma 1.4. Let $\{X, X_k, k \geq 1\}$ be defined as in (1.1), and let $\{\xi_i, -\infty < i < \infty\}$ be a sequence of identically distributed ϕ -mixing random variables with $E\xi_1 = 0$, $E\xi_1^2 < \infty$, $0 < \sigma^2 = E\xi_1^2 + 2 \sum_{k=2}^{\infty} E\xi_1\xi_k < \infty$, $\sum_{m=1}^{\infty} \phi^{1/2}(2^m) < \infty$, then

$$\frac{S_n}{\tau\sqrt{n}} \xrightarrow{D} N(0, 1). \quad (1.7)$$

The proof is similar to Theorem 1 in [11]. Set $\Delta_n = \sup_x |P(|S_n| \geq \sqrt{n}x) - P(|N| \geq x)|$. From Lemma 1.4, one can get $\Delta_n \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 1.5 (see [2]). Let $\sum_{i=-\infty}^{+\infty} a_i$ be an absolutely convergent series of real numbers with $a = \sum_{i=-\infty}^{+\infty} a_i$ and $k \geq 1$, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=-\infty}^{+\infty} \left| \sum_{j=i+1}^{i+n} a_j \right|^k = |a|^k. \quad (1.8)$$

Lemma 1.6 (see [12]). Let $\{X_i, i \geq 1\}$ be a sequence of ϕ -mixing random variables with zero means and finite second moments. Let $S_n = \sum_{i=1}^n X_i$. If exists C_n such that $\max_{1 \leq i \leq n} E S_i^2 \leq C_n$, then for all $q \geq 2$, there exists $C = C(q, \phi(\cdot))$ such that

$$E \max_{1 \leq i \leq n} |S_i|^q \leq C \left(C_n^{q/2} + E \max_{1 \leq i \leq n} |X_i|^q \right). \quad (1.9)$$

2. Proofs

Proof of Theorem 1.1. Without loss of generality, we assume that $\tau = 1$. We have

$$\begin{aligned} & \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-1}}{n^2} E S_n^2 I\left\{|S_n| \geq \sqrt{n \log n \epsilon}\right\} \\ &= \epsilon^2 \sum_{n=2}^{\infty} \frac{(\log n)^{\delta}}{n} P\left\{|S_n| \geq \sqrt{n \log n \epsilon}\right\} + \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-1}}{n^2} \int_{\sqrt{n \log n \epsilon}}^{\infty} 2x P(|S_n| \geq x) dx. \end{aligned} \quad (2.1)$$

Set $d(\epsilon) = \exp(M\epsilon^{-2})$, where $M > 1$. By Theorem B, we need to show

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-1}}{n^2} \int_{\sqrt{n \log n \epsilon}}^{\infty} 2x P(|S_n| \geq x) dx = \frac{1}{\delta(\delta+1)} E|N|^{2\delta+2}. \quad (2.2)$$

By Proposition 5.1 in [10], we have

$$\lim_{\epsilon \searrow 0} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-1}}{n^2} \int_{\sqrt{n \log n \epsilon}}^{\infty} 2x P\left(|N| \geq \frac{x}{\sqrt{n}}\right) dx = \frac{1}{\delta(\delta+1)} E|N|^{2\delta+2}. \quad (2.3)$$

Hence, Theorem 1.1 will be proved if we show the following two propositions. \square

Proposition 2.1. *One has*

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta} \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-1}}{n^2} \left| \int_{\sqrt{n \log n \epsilon}}^{\infty} 2x P(|S_n| \geq x) dx - \int_{\sqrt{n \log n \epsilon}}^{\infty} 2x P\left(|N| \geq \frac{x}{\sqrt{n}}\right) dx \right| = 0. \quad (2.4)$$

Proof. Write

$$\begin{aligned} & \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-1}}{n^2} \left| \int_{\sqrt{n \log n \epsilon}}^{\infty} 2x P(|S_n| \geq x) dx - \int_{\sqrt{n \log n \epsilon}}^{\infty} 2x P\left(|N| \geq \frac{x}{\sqrt{n}}\right) dx \right| \\ & \leq \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-1}}{n} \log n \\ & \quad \times \int_0^{\infty} 2(x+\epsilon) \left| P\left(|S_n| \geq \sqrt{n \log n}(x+\epsilon)\right) - P\left(|N| \geq \sqrt{\log n}(x+\epsilon)\right) \right| dx \\ & \leq \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-1}}{n} (\Delta_{n1} + \Delta_{n2} + \Delta_{n3}), \end{aligned} \quad (2.5)$$

where

$$\begin{aligned}
\Delta_{n1} &= \log n \int_0^{1/\sqrt{\log n \Delta_n^{1/4}}} 2(x+\epsilon) \left| P(|S_n| \geq \sqrt{n \log n}(x+\epsilon)) - P(|N| \geq \sqrt{\log n}(x+\epsilon)) \right| dx, \\
\Delta_{n2} &= \log n \int_{1/\sqrt{\log n \Delta_n^{1/4}}}^{\infty} 2(x+\epsilon) P(|S_n| \geq \sqrt{n \log n}(x+\epsilon)) dx, \\
\Delta_{n3} &= \log n \int_{1/\sqrt{\log n \Delta_n^{1/4}}}^{\infty} 2(x+\epsilon) P(|N| \geq \sqrt{\log n}(x+\epsilon)) dx.
\end{aligned} \tag{2.6}$$

Since $n \leq d(\epsilon)$ implies $\sqrt{\log n \epsilon} \leq \sqrt{M}$, we have

$$\Delta_{n1} \leq C \log n \Delta_n \left(\frac{1}{\sqrt{\log n \Delta_n^{1/4}}} + \epsilon \right)^2 \leq C (\Delta_n^{1/4} + \sqrt{M \Delta_n})^2. \tag{2.7}$$

For Δ_{n3} , by Markov's inequality, we get

$$\Delta_{n3} \leq C \log n \int_{1/\sqrt{\log n \Delta_n^{1/4}}}^{\infty} \frac{1}{(\log n)^{3/2} (x+\epsilon)^2} dx \leq C \Delta_n^{1/4}. \tag{2.8}$$

From (2.7) and (2.8), we can get

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta} \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-1}}{n} (\Delta_{n1} + \Delta_{n3}) = 0. \tag{2.9}$$

Note that $\sum_{k=1}^n X_k = \sum_{i=-\infty}^{\infty} \sum_{k=1}^n a_{k+i} \xi_i = \sum_{i=-\infty}^{\infty} a_{ni} \xi_i$, where $a_{ni} = \sum_{k=1}^n a_{k+i}$. By Lemma 1.5, we can assume that

$$\sum_{i=-\infty}^{\infty} |a_{ni}|^t \leq n, \quad t \geq 1, \quad \tilde{a} = \sum_{i=-\infty}^{\infty} |a_i| \leq 1. \tag{2.10}$$

Set $S'_n = \sum_{i=-\infty}^{\infty} a_{ni} \xi_i I\{|a_{ni} \xi_i| \leq \sqrt{n \log n}(x+\epsilon)\}$. As $E \xi_1 = 0$, by (2.10), we have

$$\begin{aligned}
|ES'_n| &\leq C \left| E \sum_{i=-\infty}^{\infty} a_{ni} \xi_i I\{|a_{ni} \xi_i| > \sqrt{n \log n}(x+\epsilon)\} \right| \\
&\leq C \sum_{i=-\infty}^{\infty} |a_{ni}| E |\xi_1| I\{|a_{ni} \xi_1| > \sqrt{n \log n}(x+\epsilon)\} \\
&\leq C n E |\xi_1| I\{\tilde{a} |\xi_1| > \sqrt{n \log n}(x+\epsilon)\}
\end{aligned}$$

$$\begin{aligned}
&\leq CnE|\xi_1|I\left\{|\xi_1| > \sqrt{n \log n}(x + \epsilon)\right\} \\
&\leq Cn\left(E\xi_1^2\right)^{1/2}\left(P\left(|\xi_1| > \sqrt{n \log n}(x + \epsilon)\right)\right)^{1/2} \\
&\leq C\frac{\sqrt{n}E\xi_1^2}{\sqrt{\log n}(x + \epsilon)}. \\
\end{aligned} \tag{2.11}$$

So, when $x \in (1/\sqrt{\log n}\Delta_n^{1/4}, \infty)$,

$$\frac{|ES'_n|}{\sqrt{n \log n}(x + \epsilon)} \leq C\frac{E\xi_1^2}{\log n\left(1/\sqrt{\log n}\Delta_n^{1/4} + \epsilon\right)^2} < \epsilon, \quad \text{for } n \text{ large enough.} \tag{2.12}$$

By (2.12), we have

$$\begin{aligned}
&\sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-1}}{n} \Delta_{n2} \\
&\leq \sum_{n=2}^{d(\epsilon)} \int_{1/\sqrt{\log n}\Delta_n^{1/4}}^{\infty} \frac{(\log n)^\delta(x + \epsilon)}{n} \\
&\quad \times \left[P\left(\sup_i |a_{ni}\xi_i| > \sqrt{n \log n}(x + \epsilon)\right) + P\left(|S'_n - ES'_n| \geq \frac{\sqrt{n \log n}(x + \epsilon)}{2}\right) \right] dx \\
&=: H_1 + H_2. \\
\end{aligned} \tag{2.13}$$

Set $I_{nj} = \{j \in \mathcal{L}, 1/(j+1) < |a_{nj}| \leq 1/j, j = 1, 2, \dots\}$, then $\bigcup_{j \geq 1} I_{nj} = \mathcal{L}$ (referred by [4]). We can get

$$\sum_{j=1}^k \#I_{nj} \leq n(k+1). \tag{2.14}$$

Then,

$$\begin{aligned}
&P\left\{\sup_i |a_{ni}\xi_i| > \sqrt{n \log n}(x + \epsilon)\right\} \\
&\leq \sum_{i=-\infty}^{\infty} P\left\{|a_{ni}\xi_i| > \sqrt{n \log n}(x + \epsilon)\right\}
\end{aligned}$$

$$\begin{aligned}
&\leq \sum_{j=1}^{\infty} \sum_{i \in I_{nj}} P\left\{ |\xi_1| \geq \sqrt{n \log n j} (x + \epsilon) \right\} \leq \sum_{j=1}^{\infty} (\#I_{nj}) P\left\{ |\xi_1| \geq \sqrt{n \log n j} (x + \epsilon) \right\} \\
&\leq \sum_{j=1}^{\infty} \sum_{k \geq j} (\#I_{nj}) P\left\{ \sqrt{n \log n k} (x + \epsilon) \leq |\xi_1| < \sqrt{n \log n} (k+1) (x + \epsilon) \right\} \\
&\leq \sum_{k=1}^{\infty} \sum_{j=1}^k (\#I_{nj}) P\left\{ \sqrt{n \log n k} (x + \epsilon) \leq |\xi_1| < \sqrt{n \log n} (k+1) (x + \epsilon) \right\} \\
&\leq \sum_{k=1}^{\infty} n(k+1) P\left\{ \sqrt{n \log n k} (x + \epsilon) \leq |\xi_1| < \sqrt{n \log n} (k+1) (x + \epsilon) \right\} \\
&\leq \frac{\sqrt{n} E|\xi_1| I\left\{ |\xi_1| \geq \sqrt{n \log n} (x + \epsilon) \right\}}{\sqrt{\log n} (x + \epsilon)}. \tag{2.15}
\end{aligned}$$

So, we get

$$\begin{aligned}
H_1 &\leq CE|\xi_1| \int_{1/\sqrt{\log n} \Delta_n^{1/4}}^{\infty} \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-1/2}}{n^{1/2}} \\
&\quad \times \sum_{k=n}^{\infty} I\left\{ \sqrt{k \log k} (x + \epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x + \epsilon) \right\} dx \\
&\leq C \int_0^{\infty} \sum_{k=2}^{\infty} E|\xi_1| I\left\{ \sqrt{k \log k} (x + \epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x + \epsilon) \right\} dx \sum_{n=2}^k \frac{(\log n)^{\delta-1/2}}{n^{1/2}} \\
&\leq CE\xi_1^2 \int_0^{\infty} (x + \epsilon)^{-1} \sum_{k=2}^{\infty} (\log k)^{\delta-1} I\left\{ \sqrt{k \log k} (x + \epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x + \epsilon) \right\} dx \\
&\leq CE\xi_1^2 \int_0^{\infty} |\log^+ |\xi_1| - \log(x + \epsilon)|^{(\delta-1)} (x + \epsilon)^{-1} I\{|\xi_1| > (x + \epsilon)\} dx \\
&\leq CE\xi_1^2 |\log^+ |\xi_1| - \log \epsilon|^{\delta} \leq CE\xi_1^2 (\log^+ |\xi_1|)^{\delta} + CE\xi_1^2 (-\log \epsilon)^{\delta}. \tag{2.16}
\end{aligned}$$

Therefore,

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta} H_1 = 0. \tag{2.17}$$

By Lemma 1.6, noting that $\sum_{m=1}^{\infty} \phi^{1/2}(m) < \infty$, for $q > 2$,

$$\begin{aligned}
H_2 &\leq C \sum_{n=2}^{d(\epsilon)} \int_0^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} (x+\epsilon)^{1-q} \\
&\quad \times \left\{ \left(\sum_{i=-\infty}^{\infty} E(a_{ni}\xi_1)^2 I\left\{ |a_{ni}\xi_1| \leq \sqrt{n \log n} (x+\epsilon) \right\} \right)^{q/2} \right. \\
&\quad + \sum_{i=-\infty}^{\infty} E|a_{ni}\xi_1|^q I\left\{ |a_{ni}\xi_1| \leq \sqrt{n \log n} \epsilon \right\} \\
&\quad \left. + \sum_{i=-\infty}^{\infty} E|a_{ni}\xi_1|^q I\left\{ \sqrt{n \log n} \epsilon < |a_{ni}\xi_1| \leq \sqrt{n \log n} (x+\epsilon) \right\} \right\} dx \\
&=: H_{21} + H_{22} + H_{23}.
\end{aligned} \tag{2.18}$$

For H_{21} , we have

$$\begin{aligned}
H_{21} &\leq \sum_{n=2}^{d(\epsilon)} \int_0^{\infty} \frac{(\log n)^{\delta-q/2}}{n} (x+\epsilon)^{1-q} \left(E\xi_1^2 I\left\{ |a_{ni}\xi_1| \leq \sqrt{n \log n} (x+\epsilon) \right\} \right)^{q/2} dx \\
&\leq C\epsilon^{-2\delta} M^{\delta+1-q/2}.
\end{aligned} \tag{2.19}$$

Then, for $0 < \delta \leq 1$, $q > 2$, we have

$$\lim_{M \rightarrow \infty} \limsup_{\epsilon \searrow 0} \epsilon^{2\delta} H_{21} = 0. \tag{2.20}$$

For H_{22} , we decompose it into two parts,

$$\begin{aligned}
H_{22} &\leq \sum_{n=2}^{d(\epsilon)} \int_0^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} (x+\epsilon)^{1-q} \sum_{j=1}^{\infty} \sum_{i \in I_{nj}} |a_{ni}|^q E|\xi_1|^q I\left\{ |a_{ni}\xi_1| \leq \sqrt{n \log n} \epsilon \right\} dx \\
&\leq \epsilon^{2-q} \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} \\
&\quad \times \sum_{j=1}^{\infty} (\#I_{nj}) j^{-q} \left\{ \sum_{k=0}^{2n} E|\xi_1|^q I\left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \right. \\
&\quad \left. + \sum_{k=2n+1}^{(j+1)n} E|\xi_1|^q I\left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \right\} \\
&=: H_{221} + H_{222}.
\end{aligned} \tag{2.21}$$

It is easy to see that

$$\sum_{j=m}^{\infty} (\#I_{nj}) (j+1)^{-q} (m+1)^{q-1} \leq \sum_{j=1}^{\infty} (\#I_{nj}) (j+1)^{-1} \leq \sum_{i=-\infty}^{\infty} |a_{ni}| = \sum_{j=1}^{\infty} \sum_{i \in I_{nj}} |a_{ni}| \leq n. \quad (2.22)$$

So,

$$\sum_{j=m}^{\infty} (\#I_{nj}) j^{-q} \leq Cnm^{-(q-1)}. \quad (2.23)$$

Now, we estimate H_{221} , by (2.23),

$$\begin{aligned} H_{221} &\leq \epsilon^{2-q} \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{q/2}} \sum_{k=0}^{2n} E|\xi_1|^q I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq \epsilon^{2-q} \sum_{k=2}^{d(\epsilon)} E|\xi_1|^q I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \sum_{n=[k/2]}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{q/2}} \\ &\leq \epsilon^{2-q} \sum_{k=2}^{d(\epsilon)} \frac{(\log k)^{\delta-q/2}}{k^{q/2-1}} E|\xi_1|^q I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq \sum_{k=2}^{d(\epsilon)} (\log k)^{\delta-1} E\xi_1^2 I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq \sum_{k=2}^{d(\epsilon)} (\log k)^{-1} |\log^+ |\xi_1| - \log \epsilon|^{\delta} E\xi_1^2 I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq CE\xi_1^2 (\log^+ |\xi_1|)^{\delta} + CE\xi_1^2 (-\log \epsilon)^{\delta}. \end{aligned} \quad (2.24)$$

For H_{222} , we have

$$\begin{aligned} H_{222} &\leq \epsilon^{2-q} \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} \\ &\times \sum_{k=2n+1}^{\infty} \sum_{j \geq k/n-1} (\#I_{nj}) j^{-q} E|\xi_1|^q I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq \epsilon^{2-q} \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{1-q/2}} \sum_{k=2n+1}^{\infty} k^{1-q} E|\xi_1|^q I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq \sum_{k=2}^{d(\epsilon)} (\log k)^{\delta-1} E\xi_1^2 I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq \sum_{k=2}^{d(\epsilon)} (\log k)^{-1} |\log^+ |\xi_1| - \log \epsilon|^{\delta} E\xi_1^2 I \left\{ \sqrt{k \log k} \epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\leq CE\xi_1^2 (\log^+ |\xi_1|)^{\delta} + CE\xi_1^2 (-\log \epsilon)^{\delta}. \end{aligned} \quad (2.25)$$

From (2.24) and (2.25), we can get

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta} H_{22} = \lim_{\epsilon \searrow 0} \epsilon^{2\delta} H_{221} + \lim_{\epsilon \searrow 0} \epsilon^{2\delta} H_{222} = 0. \quad (2.26)$$

Finally, $q > 2$, and we will get

$$\begin{aligned} H_{23} &\leq C \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} E|\xi_1|^q I\left\{ |\tilde{a}\xi_1| > \sqrt{n \log n \epsilon} \right\} \\ &\quad \times \int_0^\infty (x+\epsilon)^{1-q} \sum_{j=1}^\infty (\#I_{nj}) j^{-q} \left\{ \sum_{k=0}^{2n} + \sum_{k=2n+1}^{(j+1)n} \right\} \\ &\quad \times I\left\{ \sqrt{k \log k} (x+\epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x+\epsilon) \right\} dx \\ &\leq C \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{q/2}} E|\xi_1|^q I\left\{ |\tilde{a}\xi_1| > \sqrt{n \log n \epsilon} \right\} \\ &\quad \times \int_0^\infty (x+\epsilon)^{1-q} I\left\{ |\xi_1| \leq \sqrt{2n \log(2n)} (x+\epsilon) \right\} dx + C \sum_{n=2}^{d(\epsilon)} \frac{(\log n)^{\delta-q/2}}{n^{1-q/2}} E|\xi_1|^q \\ &\quad \times \int_0^\infty \sum_{k=2n+1}^\infty \frac{k^{1-q}}{(x+\epsilon)^{q-1}} I\left\{ \sqrt{k \log k} (x+\epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x+\epsilon) \right\} dx \\ &\leq C \sum_{k=2}^\infty E\xi_1^2 I\left\{ \sqrt{k \log k} \epsilon < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} \epsilon \right\} \sum_{n=2}^k \frac{(\log n)^{\delta-1}}{n} \\ &\quad + CE\xi_1^2 \int_0^\infty (x+\epsilon)^{-1} \\ &\quad \times \sum_{k=4}^\infty (\log k)^{\delta-1} I\left\{ \sqrt{k \log k} (x+\epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x+\epsilon) \right\} dx \\ &\leq C \sum_{k=2}^\infty (\log k)^\delta E\xi_1^2 I\left\{ \sqrt{k \log k} \epsilon < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\ &\quad + CE\xi_1^2 \int_0^\infty |\log^+ |\xi_1| - \log(x+\epsilon)|^{(\delta-1)} (x+\epsilon)^{-1} I\{|\xi_1| > (x+\epsilon)\} dx \\ &\leq CE\xi_1^2 |\log^+ |\xi_1| - \log \epsilon|^\delta \leq CE\xi_1^2 (\log^+ |\xi_1|)^\delta + CE\xi_1^2 (-\log \epsilon)^\delta, \end{aligned} \quad (2.27)$$

then

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta} H_{23} = 0. \quad (2.28)$$

Hence, (2.4) can be referred from (2.9), (2.17), (2.20), (2.26), and (2.28). \square

Proposition 2.2. *One has*

$$\lim_{\epsilon \searrow 0} \epsilon^{2\delta} \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-1}}{n^2} \left| \int_{\sqrt{n \log n \epsilon}}^{\infty} 2xP(|S_n| \geq x)dx - \int_{\sqrt{n \log n \epsilon}}^{\infty} 2xP\left(|N| \geq \frac{x}{\sqrt{n}}\right)dx \right| = 0. \quad (2.29)$$

Proof. Consider the following:

$$\begin{aligned} & \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-1}}{n^2} \left| \int_{\sqrt{n \log n \epsilon}}^{\infty} 2xP(|S_n| \geq x)dx - \int_{\sqrt{n \log n \epsilon}}^{\infty} 2xP\left(|N| \geq \frac{x}{\sqrt{n}}\right)dx \right| \\ & \leq \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta}}{n} \int_0^{\infty} 2(x+\epsilon)P\left(|N| \geq \sqrt{\log n}(x+\epsilon)\right) \\ & \quad + \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta}}{n} \int_0^{\infty} 2(x+\epsilon)P\left(|S_n| \geq \sqrt{n \log n}(x+\epsilon)\right)dx \\ & =: G_1 + G_2. \end{aligned} \quad (2.30)$$

We first estimate G_1 , for $\theta > 2\delta$, by Markov's inequality,

$$\begin{aligned} G_1 & \leq \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta}}{n} \int_0^{\infty} \frac{1}{(\log n)^{(\theta+2)/2} (x+\epsilon)^{\theta+1}} dx \\ & \leq CM^{\delta-\theta/2} \epsilon^{-2\delta}. \end{aligned} \quad (2.31)$$

Hence,

$$\lim_{M \rightarrow \infty} \limsup_{\epsilon \searrow 0} \epsilon^{2\delta} G_1 = 0. \quad (2.32)$$

Now, we estimate G_2 . Here, $n > M\epsilon^{-2}$, so

$$\frac{|ES'_n|}{\sqrt{n \log n}(x+\epsilon)} < \frac{E\xi_1^2}{(\log n)(x+\epsilon)^2} < \frac{E\xi_1^2}{M} < \epsilon, \quad \text{for } M \rightarrow \infty. \quad (2.33)$$

We have

$$\begin{aligned}
G_2 &\leq \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta}}{n} \int_0^{\infty} (x+\epsilon) \left[P\left(\sup_i |a_{ni}\xi_i| > \sqrt{n \log n}(x+\epsilon) \right) \right. \\
&\quad \left. + P\left(|S'_n - ES'_n| \geq \frac{\sqrt{n \log n}(x+\epsilon)}{2} \right) \right] dx \\
&=: G_{21} + G_{22}.
\end{aligned} \tag{2.34}$$

We estimate G_{21} first. Similar to the proof of (2.16), we have

$$\begin{aligned}
G_{21} &\leq CE|\xi_1| \int_0^{\infty} \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-1/2}}{n^{1/2}} \\
&\quad \times \sum_{k=n}^{\infty} I\left\{ \sqrt{k \log k}(x+\epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)}(x+\epsilon) \right\} dx \\
&\leq CE|\xi_1| \int_0^{\infty} \sum_{k=d(\epsilon)+1}^{\infty} I\left\{ \sqrt{k \log k}(x+\epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)}(x+\epsilon) \right\} dx \\
&\quad \times \sum_{n=d(\epsilon)+1}^k \frac{(\log n)^{\delta-1/2}}{n^{1/2}} \\
&\leq CE\xi_1^2 \int_0^{\infty} \sum_{k=d(\epsilon)+1}^{\infty} (\log n)^{\delta-1} (x+\epsilon)^{-1} \\
&\quad \times I\left\{ \sqrt{k \log k}(x+\epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)}(x+\epsilon) \right\} dx \\
&\leq CE\xi_1^2 \int_0^{\infty} |\log |\xi_1| - \log(x+\epsilon)|^{(\delta-1)} (x+\epsilon)^{-1} I\{|\xi_1| > (x+\epsilon)\} dx \\
&\leq CE\xi_1^2 |\log^+ |\xi_1| - \log \epsilon|^{\delta} \leq CE\xi_1^2 (\log^+ |\xi_1|)^{\delta} + CE\xi_1^2 (-\log \epsilon)^{\delta},
\end{aligned} \tag{2.35}$$

then

$$\lim_{M \rightarrow \infty} \limsup_{\epsilon \searrow 0} \epsilon^{2\delta} G_{21} = 0. \tag{2.36}$$

By Lemma 1.6, for $q > 2$, we have

$$\begin{aligned}
G_{22} &= \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta}}{n} \int_0^{\infty} (x + \epsilon) P \left\{ |S'_n - ES'_n| \geq \frac{\sqrt{n \log n}(x + \epsilon)}{2} \right\} dx \\
&\leq \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} \\
&\quad \times \int_0^{\infty} (x + \epsilon)^{1-q} \left\{ \left(\sum_{i=-\infty}^{\infty} E(a_{ni}\xi_1)^2 I \left\{ |a_{ni}\xi_1| \leq \sqrt{n \log n}(x + \epsilon) \right\} \right)^{q/2} \right. \\
&\quad + \sum_{i=-\infty}^{\infty} E|a_{ni}\xi_1|^q I \left\{ |a_{ni}\xi_1| \leq \sqrt{n \log n}\epsilon \right\} \\
&\quad \left. + \sum_{i=-\infty}^{\infty} E|a_{ni}\xi_1|^q I \left\{ \sqrt{n \log n}\epsilon < |a_{ni}\xi_1| \leq \sqrt{n \log n}(x + \epsilon) \right\} \right\} dx \\
&=: G_{221} + G_{222} + G_{223}.
\end{aligned} \tag{2.37}$$

For G_{221} , we have

$$\begin{aligned}
G_{221} &\leq \sum_{n=d(\epsilon)+1}^{\infty} \int_0^{\infty} \frac{(\log n)^{\delta-q/2}}{n} (x + \epsilon)^{1-q} \left(E\xi_1^2 I \{ |a_{ni}\xi_1| \leq n(x + \epsilon) \} \right)^{q/2} dx \\
&\leq Ce^{2-q} \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n} \leq CM^{\delta+1-q/2}e^{-2\delta}.
\end{aligned} \tag{2.38}$$

Next, turning to G_{222} , it follows that

$$\begin{aligned}
G_{222} &\leq \epsilon^{2-q} \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} \\
&\quad \times \sum_{j=1}^{\infty} (\#I_{nj}) j^{-q} \left\{ \sum_{k=2}^{2n} E|\xi_1|^q I \left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \right. \\
&\quad \left. + \sum_{k=2n+1}^{(j+1)n} E|\xi_1|^q I \left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \right\} \\
&=: G_{2221} + G_{2222},
\end{aligned} \tag{2.39}$$

then

$$\begin{aligned}
G_{2221} &\leq C\epsilon^{2-q} \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{q/2}} \sum_{k=2}^{2n} E|\xi_1|^q I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \\
&\leq C\epsilon^{2-q} \sum_{k=2}^{\infty} E|\xi_1|^q I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \sum_{n=[k/2]}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{q/2}} \\
&\leq C\epsilon^{2-q} \sum_{k=2}^{\infty} \frac{(\log k)^{\delta-q/2}}{k^{q/2-1}} E|\xi_1|^q I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \\
&\leq C \sum_{k=2}^{\infty} (\log k)^{\delta-1} E\xi_1^2 I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \\
&\leq C \sum_{k=2}^{\infty} (\log k)^{-1} |\log^+|\xi_1| - \log \epsilon|^{\delta} E\xi_1^2 I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \\
&\leq CE\xi_1^2 (\log^+|\xi_1|)^{\delta} + CE\xi_1^2 (-\log \epsilon)^{\delta}. \tag{2.40}
\end{aligned}$$

For G_{2222} , it follows that

$$\begin{aligned}
G_{2222} &\leq C\epsilon^{2-q} \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} \\
&\quad \times \sum_{k=2n+1}^{\infty} \sum_{j \geq k/n-1} (\#I_{nj}) j^{-q} E|\xi_1|^q I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \\
&\leq C\epsilon^{2-q} \sum_{k=d(\epsilon)+1}^{\infty} k^{1-q} E|\xi_1|^q I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \sum_{n=d(\epsilon)+1}^{[k/2]} \frac{(\log n)^{\delta-q/2}}{n^{1-q/2}} \\
&\leq C \sum_{k=d(\epsilon)+1}^{\infty} (\log k)^{\delta-1} E\xi_1^2 I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \\
&\leq C \sum_{k=d(\epsilon)+1}^{\infty} (\log k)^{-1} |\log^+|\xi_1| - \log \epsilon|^{\delta} E\xi_1^2 I\left\{ \sqrt{k \log k}\epsilon \leq |\xi_1| < \sqrt{(k+1) \log(k+1)}\epsilon \right\} \\
&\leq CE\xi_1^2 (\log^+|\xi_1|)^{\delta} + CE\xi_1^2 (-\log \epsilon)^{\delta}. \tag{2.41}
\end{aligned}$$

Finally, $q > 2$, we have

$$\begin{aligned}
G_{223} &\leq C \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{1+q/2}} E|\xi_1|^q I\left\{ |\tilde{a}\xi_1| > \sqrt{n \log n} \epsilon \right\} \\
&\quad \times \int_0^{\infty} (x + \epsilon)^{1-q} \sum_{j=1}^{\infty} (\#I_{nj}) j^{-q} \left\{ \sum_{k=0}^{2n} + \sum_{k=2n+1}^{(j+1)n} \right\} \\
&\quad \times I\left\{ \sqrt{k \log k} (x + \epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x + \epsilon) \right\} dx \\
&\leq C \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{q/2}} E|\xi_1|^q I\left\{ |\tilde{a}\xi_1| > \sqrt{n \log n} \epsilon \right\} \\
&\quad \times \int_0^{\infty} (x + \epsilon)^{1-q} I\left\{ |\xi_1| \leq \sqrt{2n \log(2n)} (x + \epsilon) \right\} dx + C \sum_{n=d(\epsilon)+1}^{\infty} \frac{(\log n)^{\delta-q/2}}{n^{1-q/2}} E|\xi_1|^q \\
&\quad \times \int_0^{\infty} \sum_{k=2n+1}^{\infty} \frac{k^{1-q}}{(x + \epsilon)^{q-1}} I\left\{ \sqrt{k \log k} (x + \epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x + \epsilon) \right\} dx \\
&\leq C \sum_{k=d(\epsilon)+1}^{\infty} E\xi_1^2 I\left\{ \sqrt{k \log k} \epsilon < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} \epsilon \right\} \sum_{n=d(\epsilon)+1}^k \frac{(\log n)^{\delta-1}}{n} \\
&\quad + CE\xi_1^2 \int_0^{\infty} (x + \epsilon)^{-1} \\
&\quad \times \sum_{k=d(\epsilon)+1}^{\infty} (\log k)^{\delta-1} I\left\{ \sqrt{k \log k} (x + \epsilon) < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} (x + \epsilon) \right\} dx \\
&\leq C \sum_{k=2}^{\infty} (\log k)^{\delta} E\xi_1^2 I\left\{ \sqrt{k \log k} \epsilon < |\xi_1| \leq \sqrt{(k+1) \log(k+1)} \epsilon \right\} \\
&\quad + CE\xi_1^2 \int_0^{\infty} |\log^+ |\xi_1| - \log(x + \epsilon)|^{(\delta-1)} (x + \epsilon)^{-1} I\{|\xi_1| > (x + \epsilon)\} dx \\
&\leq CE\xi_1^2 |\log^+ |\xi_1| - \log \epsilon|^\delta \leq CE\xi_1^2 (\log^+ |\xi_1|)^\delta + CE\xi_1^2 (-\log \epsilon)^\delta.
\end{aligned} \tag{2.42}$$

From (2.38) to (2.42), we can get

$$\lim_{M \rightarrow \infty} \lim_{\epsilon \searrow 0} e^{2\delta} G_{22} = 0. \tag{2.43}$$

(2.29) can be derived by (2.32), (2.36), and (2.43). \square

Proof of Theorem 1.2. Without loss of generality, we set $\tau = 1$. It is easy to see that

$$\begin{aligned} & \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-1}}{n^2 \log n} E S_n^2 I\left\{|S_n| \geq \sqrt{n \log \log n \epsilon}\right\} \\ &= \epsilon^2 \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta}}{n \log n} P\left\{|S_n| \geq \sqrt{n \log \log n \epsilon}\right\} \\ &+ \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-1}}{n^2 \log n} \int_{\sqrt{n \log \log n \epsilon}}^{\infty} 2xP(|S_n| \geq x)dx. \end{aligned} \quad (2.44)$$

So, we only prove the following two propositions:

$$\epsilon^{2\delta+2} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta}}{n \log n} P\left\{|S_n| \geq \sqrt{n \log \log n \epsilon}\right\} = \frac{E|N|^{2(\delta+1)}}{\delta+1}, \quad (2.45)$$

$$\epsilon^{2\delta} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-1}}{n^2 \log n} \int_{\sqrt{n \log \log n \epsilon}}^{\infty} 2xP(|S_n| \geq x)dx = \frac{E|N|^{2(\delta+1)}}{\delta(\delta+1)}. \quad (2.46)$$

The proof of (2.45) can be referred to [6], and the proof of (2.46) is similar to Propositions 2.1 and 2.2. \square

Acknowledgments

The author would like to thank the referee for many valuable comments. This research was supported by Humanities and Social Sciences Planning Fund of the Ministry of Education of PRC. (no. 08JA790118)

References

- [1] I. A. Ibragimov, "Some limit theorems for stationary processes," *Theory of Probability and Its Applications*, vol. 7, pp. 349–382, 1962.
- [2] R. M. Burton and H. Dehling, "Large deviations for some weakly dependent random processes," *Statistics & Probability Letters*, vol. 9, no. 5, pp. 397–401, 1990.
- [3] X. Y. Yang, "The law of the iterated logarithm and the central limit theorem with random indices for B-valued stationary linear processes," *Chinese Annals of Mathematics Series A*, vol. 17, no. 6, pp. 703–714, 1996.
- [4] D. L. Li, M. B. Rao, and X. C. Wang, "Complete convergence of moving average processes," *Statistics & Probability Letters*, vol. 14, no. 2, pp. 111–114, 1992.
- [5] L.-X. Zhang, "Complete convergence of moving average processes under dependence assumptions," *Statistics & Probability Letters*, vol. 30, no. 2, pp. 165–170, 1996.
- [6] Y. X. Li and L. X. Zhang, "Precise asymptotics in the law of the iterated logarithm of moving-average processes," *Acta Mathematica Sinica (English Series)*, vol. 22, no. 1, pp. 143–156, 2006.
- [7] P. L. Hsu and H. Robbins, "Complete convergence and the law of large numbers," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 33, pp. 25–31, 1947.
- [8] J. A. Davis, "Convergence rates for probabilities of moderate deviations," *Annals of Mathematical Statistics*, vol. 39, pp. 2016–2028, 1968.

- [9] A. Gut and A. Spătaru, "Precise asymptotics in the law of the iterated logarithm," *The Annals of Probability*, vol. 28, no. 4, pp. 1870–1883, 2000.
- [10] W. Liu and Z. Lin, "Precise asymptotics for a new kind of complete moment convergence," *Statistics & Probability Letters*, vol. 76, no. 16, pp. 1787–1799, 2006.
- [11] T.-S. Kim and J.-I. Baek, "A central limit theorem for stationary linear processes generated by linearly positively quadrant-dependent process," *Statistics & Probability Letters*, vol. 51, no. 3, pp. 299–305, 2001.
- [12] Q. M. Shao, "A moment inequality and its applications," *Acta Mathematica Sinica*, vol. 31, no. 6, pp. 736–747, 1988 (Chinese).