


# Journal of Inequalities in Pure and Applied Mathematics

## NOTE ON THE PERTURBED TRAPEZOID INEQUALITY

XIAO-LIANG CHENG AND JIE SUN

Department of Mathematics  
Zhejiang University,  
Xixi Campus, Zhejiang 310028,  
The People's Republic of China.  
EMail: [xlcheng@mail.hz.zj.cn](mailto:xlcheng@mail.hz.zj.cn)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
046-01



---

volume 3, issue 2, article 29,  
2002.

*Received 21 May, 2001;  
accepted 01 February, 2002.*

*Communicated by: N.S. Barnett*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

In this paper, we utilize a variant of the Grüss inequality to obtain some new perturbed trapezoid inequalities. We improve the error bound of the trapezoid rule in numerical integration in some recent known results. Also we give a new Iyengar's type inequality involving a second order bounded derivative for the perturbed trapezoid inequality.


In the literature [2], [4] – [8], [11], [12] on numerical integration, the following estimation is well known as the trapezoid inequality:

$$(1) \quad \left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) \right| \leq \frac{1}{12} M_2 (b-a)^3,$$

where the mapping  $f : [a, b] \rightarrow \mathbb{R}$  is supposed to be twice differentiable on the interval  $(a, b)$ , with the second derivative bounded on  $(a, b)$  by  $M_2 = \sup_{x \in (a, b)} |f''(x)| < +\infty$ . In [5], the authors derived the error bounds for the trapezoid inequality (1) by different norm of mapping  $f$ . In [2, 7, 11], the authors obtained the trapezoid inequality by the difference of sup and inf bound of the first derivative, that is,

$$\left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) \right| \leq \frac{1}{8} (\Gamma_1 - \gamma_1) (b-a)^2,$$

where  $\Gamma_1 = \sup_{x \in (a, b)} f'(x) < +\infty$  and  $\gamma_1 = \inf_{x \in (a, b)} f'(x) > -\infty$ .



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 2 of 16

For the perturbed trapezoid inequality, S. Dragomir et al. [5] obtained the following inequality by an application of the Grüss inequality:


$$(2) \quad \left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{12}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{1}{32}(\Gamma_2 - \gamma_2)(b-a)^3,$$

where  $f$  is supposed to be twice differentiable on the interval  $(a, b)$ , with the second derivative bounded on  $(a, b)$  by  $\Gamma_2 = \sup_{x \in (a,b)} f''(x) < +\infty$  and  $\gamma_2 = \inf_{x \in (a,b)} f''(x) > -\infty$ . The constant  $\frac{1}{32}$  is smaller than  $\frac{1}{6\sqrt{5}}$  given in [11] and  $\frac{1}{18\sqrt{3}}$  given in [2].

In this note we first improve the constant  $\frac{1}{32}$  in the inequality (2) to the best possible one of  $\frac{1}{36\sqrt{3}}$ . Then we give two new perturbed trapezoid inequalities for high-order differentiable mappings. We need the following variant of the Grüss inequality:

**Theorem 1.** *Let  $h, g : [a, b] \rightarrow \mathbb{R}$  be two integrable functions such that  $\phi \leq g(x) \leq \Phi$  for some constants  $\phi, \Phi$  for all  $x \in [a, b]$ , then*

$$(3) \quad \left| \frac{1}{b-a} \int_a^b h(x)g(x) dx - \frac{1}{(b-a)^2} \int_a^b h(x) dx \int_a^b g(x) dx \right| \leq \frac{1}{2} \left( \int_a^b \left| h(x) - \frac{1}{b-a} \int_a^b h(y) dy \right| dx \right) (\Phi - \phi).$$



Littlewood's Inequality for  $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

*Proof.* We write the left hand of inequality (3) as

$$\begin{aligned} \int_a^b h(x)g(x) dx - \frac{1}{b-a} \int_a^b h(x) dx \int_a^b g(x) dx \\ = \int_a^b \left( h(x) - \frac{1}{b-a} \int_a^b h(y) dy \right) g(x) dx. \end{aligned}$$

Denote

$$I^+ = \int_a^b \max\left(h(x) - \frac{1}{b-a} \int_a^b h(y) dy, 0\right) dx$$

and

$$I^- = \int_a^b \min\left(h(x) - \frac{1}{b-a} \int_a^b h(y) dy, 0\right) dx.$$

Obviously  $I^+ + I^- = 0$ . For  $\phi \leq g(x) \leq \Phi$ , then


$$\int_a^b \left( h(x) - \frac{1}{b-a} \int_a^b h(y) dy \right) g(x) dx \leq I^+ \Phi + I^- \phi$$

and

$$- \int_a^b \left( h(x) - \frac{1}{b-a} \int_a^b h(y) dy \right) g(x) dx \leq -I^+ \phi - I^- \Phi$$

and hence the obtained result (3) follows.  $\square$

**Theorem 2.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a twice differentiable mapping on  $(a, b)$  with  $\Gamma_2 = \sup_{x \in (a,b)} f''(x) < +\infty$  and  $\gamma_2 = \inf_{x \in (a,b)} f''(x) > -\infty$ , then we have



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 4 of 16

the estimation

$$(4) \quad \left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{12}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{1}{36\sqrt{3}}(\Gamma_2 - \gamma_2)(b-a)^3,$$


where the constant  $\frac{1}{36\sqrt{3}}$  is the best one in the sense that it cannot be replaced by a smaller one.

*Proof.* We choose in (3),  $h(x) = -\frac{1}{2}(x-a)(b-x)$  and  $g(x) = f''(x)$ , we get

$$\begin{aligned} \frac{1}{2} \int_a^b \left| h(x) - \frac{1}{b-a} \int_a^b h(y) dy \right| dx &= \left| \int_{x_1}^{x_2} \left( h(x) + \frac{1}{12}(b-a)^2 \right) dx \right| \\ &= \frac{1}{36\sqrt{3}}(b-a)^3, \end{aligned}$$

where  $x_1 = a + \frac{3-\sqrt{3}}{6}(b-a)$  and  $x_2 = a + \frac{3+\sqrt{3}}{6}(b-a)$ . Thus from (3), we derive

$$\begin{aligned} &\left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{12}(b-a)^2(f'(b) - f'(a)) \right| \\ &= \left| \int_a^b -\frac{1}{2}(x-a)(b-x)f''(x) dx \right. \\ &\quad \left. - \frac{1}{b-a} \int_a^b -\frac{1}{2}(x-a)(b-x) dx \int_a^b f''(x) dx \right| \\ &\leq \frac{1}{36\sqrt{3}}(\Gamma_2 - \gamma_2)(b-a)^3. \end{aligned}$$



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 5 of 16

To explain the best constant  $\frac{1}{36\sqrt{3}}$  in the inequality (4), we can construct the function  $f(x) = \int_a^x \left( \int_a^y j(z) dz \right) dy$  to attain the inequality in (4),

$$j(x) = \begin{cases} \gamma_2, & a \leq x < x_1 = a + \frac{3 - \sqrt{3}}{6}(b - a), \\ \Gamma_2, & x_1 \leq x < x_2 = a + \frac{3 + \sqrt{3}}{6}(b - a), \\ \gamma_2, & x_2 \leq x \leq b. \end{cases}$$

The proof is complete.  $\square$


**Theorem 3.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a third-order differentiable mapping on  $(a, b)$  with  $\Gamma_3 = \sup_{x \in (a,b)} f'''(x) < +\infty$  and  $\gamma_3 = \inf_{x \in (a,b)} f'''(x) > -\infty$ , then we have the estimation

$$(5) \quad \left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{12}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{1}{384}(\Gamma_3 - \gamma_3)(b-a)^4,$$

where the constant  $\frac{1}{384}$  is the best one in the sense that it cannot be replaced by a smaller one.

*Proof.* We choose in (3),  $h(x) = \frac{1}{12}(x-a)(2x-a-b)(b-x)$ ,  $g(x) = f'''(x)$ , to get

$$\frac{1}{2} \int_a^b |h(x) - \frac{1}{b-a} \int_a^b h(y) dy| dx = \int_a^{\frac{a+b}{2}} h(x) dx = \frac{1}{384}(b-a)^4,$$



Littlewood's Inequality for  $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

Thus from (3) in Theorem 1, we can derive the inequality (5) immediately. Finally, we construct the function  $f(x) = \int_a^x \left( \int_a^y \left( \int_a^z j(s) ds \right) dz \right) dy$ , where  $j(x) = \Gamma_3$  for  $a \leq x < \frac{a+b}{2}$  and  $j(x) = \gamma_3$  for  $\frac{a+b}{2} \leq x \leq b$ , then the equality holds in (5).  $\square$

**Theorem 4.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a fourth-order differentiable mapping on  $(a, b)$  with  $M_4 = \sup_{x \in (a,b)} |f^{(4)}(x)| < +\infty$ , then

$$(6) \quad \left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{12}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{1}{720} M_4 (b-a)^5,$$


where  $\frac{1}{720}$  is the best possible constant.

*Proof.* We may write the remainder of the perturbed trapezoid inequality in the kernel form

$$(7) \quad \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{12}(b-a)^2(f'(b) - f'(a)) = \int_a^b f^{(4)}(x) k_4(x) dx,$$

where  $k_4(x) = \frac{1}{24}(x-a)^2(b-x)^2$ . Then we get

$$(8) \quad \int_a^b |k_4(x)| dx = \frac{1}{24} \int_0^1 x^2(1-x)^2 dx = \frac{1}{720}.$$



Littlewood's Inequality for  $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

Then (7) – (8) imply (6). The equality holds for  $f(x) = x^4$ ,  $a \leq x \leq b$  in inequality (6).  $\square$

**Remark 0.1.** We also can prove Theorem 2 and 3 in the kernel form

$$(9) \quad \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{12}(b-a)^2(f'(b) - f'(a)) \\ = \int_a^b f^{(n)}(x)k_n(x) dx,$$


where  $k_2(x) = -\frac{1}{2}(x-a)(b-x) + \frac{1}{12}$  and  $k_3(x) = \frac{1}{12}(x-a)(2x-a-b)(b-x)$ . By the formula (7) and (9), and derive the perturbed trapezoid inequality for different norms as shown in [5].

Now we present the composite perturbed trapezoid quadrature for an equidistant partitioning of interval  $[a, b]$  into  $n$  subintervals. Applying Theorems 2 – 4, we obtain

$$\int_a^b f(x) dx = T_n(f) + R_n(f),$$

where

$$T_n(f) = \frac{b-a}{2n} \sum_{i=0}^{n-1} \left[ f\left(a + i\frac{b-a}{n}\right) + f\left(a + (i+1)\frac{b-a}{n}\right) \right] \\ - \frac{(b-a)^2}{12n^2}(f'(b) - f'(a)),$$



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16



and the remainder  $R_n(f)$  satisfies the error estimate

$$(10) \quad |R_n(f)| \leq \begin{cases} \frac{(b-a)^3}{36\sqrt{3}n^2}(\Gamma_2 - \gamma_2), & \text{if } \gamma_2 \leq f''(x) \leq \Gamma_2, \forall x \in (a, b), \\ \frac{(b-a)^4}{384n^3}(\Gamma_3 - \gamma_3), & \text{if } \gamma_3 \leq f'''(x) \leq \Gamma_3, \forall x \in (a, b) \\ \frac{(b-a)^5}{720n^4}M_4, & \text{if } |f^{(4)}(x)| \leq M_4, \forall x \in (a, b). \end{cases}$$


Then we can use (10) to get different error estimates of the composite perturbed trapezoid quadrature.

As in [5], we may also apply the Theorems 2, 3 and 4 to special means. In this case we may improve some of the bounds related to inequalities about special means as given in [5, p. 492-494].

Furthermore, we discuss the Iyengar's type inequality for the perturbed trapezoidal quadrature rule for functions whose first and second order derivatives are bounded. In [1, 3, 9, 10] they proved the following interesting inequality involving bounded derivatives.

If  $f$  is a differentiable function on  $(a, b)$  and  $|f'(x)| \leq M_1$ , then

$$\left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) \right| \leq \frac{M_1(b-a)^2}{4} - \frac{(f(b) - f(a))^2}{4M_1}.$$



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 9 of 16

If  $|f''(x)| \leq M_2$ ,  $x \in [a, b]$  for positive constant  $M_2 \in \mathbb{R}$ , then

$$\left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{8}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{M_2}{24} \left( (b-a)^3 - \left( \frac{|\Delta|}{M_2} \right)^3 \right),$$

$$\left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{8}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{M_2}{24}(b-a)^3 - \frac{\Delta_1^2(b-a)}{16M_2},$$


where

$$(11) \quad \Delta = f'(a) - 2f' \left( \frac{a+b}{2} \right) + f'(b),$$

$$\Delta_1 = f'(a) - 2 \frac{(f(b) - f(a))}{b-a} + f'(b).$$

We will prove the following inequality.

**Theorem 5.** Let  $f : I \rightarrow \mathbb{R}$ , where  $I \subseteq \mathbb{R}$  is an interval. Suppose that  $f$  is twice differentiable in the interior  $\overset{\circ}{I}$  of  $I$ , and let  $a, b \in \overset{\circ}{I}$  with  $a < b$ . If



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

$|f''(x)| \leq M_2, x \in [a, b]$  for positive constant  $M_2 \in \mathbb{R}$ . Then

$$(12) \quad \left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{8}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{1}{24}M_2(b-a)^3 - \sqrt{\frac{|\Delta_1|^3(b-a)^3}{72M_2}},$$

where  $\Delta_1$  is defined as (11).

*Proof.* Denote


$$J_f = \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{8}(b-a)^2(f'(b) - f'(a)).$$

It is easy to see that

$$J_f = \int_a^b \frac{1}{2} \left( x - \frac{a+b}{2} \right)^2 f''(x) dx,$$

and

$$(13) \quad \begin{aligned} \Delta_1 &= f'(a) - \frac{2(f(b) - f(a))}{b-a} + f'(b) \\ &= \frac{1}{b-a} \int_a^b 2 \left( x - \frac{a+b}{2} \right) f''(x) dx. \end{aligned}$$



**Littlewood's Inequality for  
 $p$ -Bimeasures**

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 11 of 16

For any  $|\varepsilon| \leq \frac{1}{8}$ , we get for  $|f''(x)| \leq M_2$ ,  $x \in [a, b]$ ,


$$\begin{aligned} & J_f + \varepsilon(b-a)^2 \Delta_1 \\ &= \int_a^b \left( \frac{1}{2} \left( x - \frac{a+b}{2} \right)^2 + 2\varepsilon(b-a) \left( x - \frac{a+b}{2} \right) \right) f''(x) dx \\ &\leq F(\varepsilon) M_2 (b-a)^3, \end{aligned}$$

where

$$\begin{aligned} F(\varepsilon) &= \frac{1}{(b-a)^3} \int_a^b \left| \frac{1}{2} \left( x - \frac{a+b}{2} \right)^2 + 2\varepsilon(b-a) \left( x - \frac{a+b}{2} \right) \right| dx \\ &= \int_0^1 \left| \frac{1}{2} \left( x - \frac{1}{2} \right)^2 + 2\varepsilon \left( x - \frac{1}{2} \right) \right| dx. \end{aligned}$$

For the case  $0 \leq \varepsilon \leq \frac{1}{8}$ , we have

$$\begin{aligned} F(\varepsilon) &= \int_0^1 \left| \frac{1}{2} \left( x - \frac{1}{2} \right)^2 + 2\varepsilon \left( x - \frac{1}{2} \right) \right| dx \\ &= \left\{ \int_0^{\frac{1}{2}-4\varepsilon} \left( \frac{1}{2} \left( x - \frac{1}{2} \right)^2 + 2\varepsilon \left( x - \frac{1}{2} \right) \right) dx \right. \\ &\quad \left. - \int_{\frac{1}{2}-4\varepsilon}^{\frac{1}{2}} \left( \frac{1}{2} \left( x - \frac{1}{2} \right)^2 + 2\varepsilon \left( x - \frac{1}{2} \right) \right) dx \right. \end{aligned}$$



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 12 of 16

$$\begin{aligned}
 & + \int_{\frac{1}{2}}^1 \left( \frac{1}{2} \left( x - \frac{1}{2} \right)^2 + 2\varepsilon \left( x - \frac{1}{2} \right) \right) dx \Big\} \\
 & = \frac{1}{24} + \frac{32}{3}\varepsilon^3.
 \end{aligned}$$

For the case  $-\frac{1}{8} \leq \varepsilon \leq 0$ , we have similarly

$$F(\varepsilon) = \frac{1}{24} - \frac{32}{3}\varepsilon^3.$$

We can prove  $|\Delta_1| \leq \frac{1}{2}(b-a)M_2$  easily from (13). Thus we choose the parameter

$$\varepsilon_* = \text{sign}(\Delta_1) \sqrt{\frac{|\Delta_1|}{32(b-a)M_2}}, \quad |\varepsilon_*| \leq \frac{1}{8}.$$


By the above inequalities, we obtain

$$J_f \leq F(\varepsilon_*)M_2 - \varepsilon_*(b-a)^2\Delta_1 \leq \frac{1}{24}(b-a)^3M_2 - \sqrt{\frac{(b-a)^3|\Delta_1|^3}{72M_2}}.$$

Replacing  $f$  with  $-f$ , we have

$$J_{-f} = -J_f \leq \frac{1}{24}(b-a)^3M_2 - \sqrt{\frac{(b-a)^3|\Delta_1|^3}{72M_2}}.$$

Thus we obtain bounds for  $|J_f|$  and prove the inequality (12). □



**Littlewood's Inequality for  
 $p$ -Bimeasures**

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 13 of 16

**Remark 0.2.** As  $|\Delta_1| \leq \frac{1}{2}M_2(b-a)$ , we have


$$\sqrt{\frac{|\Delta_1|}{M_2}} \geq \sqrt{\frac{2}{b-a}} \frac{|\Delta_1|}{M_2},$$

$$\left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) + \frac{1}{8}(b-a)^2(f'(b) - f'(a)) \right| \leq \frac{M_2}{24}(b-a)^3 - \frac{\Delta_1^2(b-a)}{6M_2}.$$

For the case  $f'(a) = f'(b) = 0$ , we have

$$(14) \quad \left| \int_a^b f(x) dx - \frac{1}{2}(b-a)(f(a) + f(b)) \right| \leq \frac{M_2}{24}(b-a)^3 - \frac{2}{3} \frac{|f(b) - f(a)|^2}{M_2(b-a)}.$$

The inequality (14) is sharper than that stated in [9, p. 69].



Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

Title Page

Contents



Go Back

Close

Quit

Page 14 of 16

## References

- [1] R.P. AGARWAL, V. ČULJAK AND J. PEČARIĆ, Some integral inequalities involving bounded higher order derivatives, *Mathl. Comput. Modelling*, **28**(3) (1998), 51–57.
- [2] X.L. CHENG, Improvement of some Ostrowski-Grüss type inequalities, *Computers Math. Applic.*, **42** (2001), 109–114.
- [3] X.L. CHENG, The Iyengar type inequality, *Appl. Math. Lett.*, **14** (2001), 975–978.
- [4] S.S. DRAGOMIR AND R.P. AGARWAL, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, *Appl. Math. Lett.*, **11**(5) (1998), 91–95.
- [5] S.S. DRAGOMIR, P. CERONE AND A. SOFO, Some remarks on the trapezoid rule in numerical integration, *Indian J. Pure Appl. Math.*, **31**(5) (2000), 475–494.
- [6] S.S. DRAGOMIR, Y.J. CHO AND S.S. KIM, Inequalities of Hadamard's type for Lipschitzian mappings and their applications, *J. Math. Anal. Appl.*, **245** (2000), 489–501.
- [7] S.S. DRAGOMIR AND S. WANG, An inequality of Ostrowski-Grüss' type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules, *Computers Math. Applic.*, **33**(11) (1997), 15–20.



---

Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

---

Title Page

Contents




Go Back

Close

Quit

Page 15 of 16

- [8] S.S. DRAGOMIR AND S. WANG, Applications of Ostrowski' inequality to the estimation of error bounds for some special means and for some numerical quadrature rules, *Appl. Math. Lett.*, **11**(1) (1998), 105–109.
- [9] N. ELEZOVIĆ AND J. PEČARIĆ, Steffensen's inequality and estimates of error in trapezoidal rule, *Appl. Math. Lett.*, **11**(6) (1998), 63–69.
- [10] K.S.K. IYENGAR, Note on an inequality, *Math. Student*, **6** (1938), 75–76.
- [11] M. MATIĆ, J. PEČARIĆ AND N. UJEVIĆ, Improvement and further generalization of inequalities of Ostrowski-Grüss type, *Computers Math. Applic.*, **39**(3-4) (2000), 161–175.
- [12] D.S. MITRINOVIĆ, J. PEČARIĆ AND A.M. FINK, *Inequalities for Functions and Their Integrals and Derivatives*, Kluwer Academic Publishers, Dordrecht, (1994).




---

Littlewood's Inequality for  
 $p$ -Bimeasures

Xiao-liang Cheng and Jie Sun

---

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16