

Journal of Inequalities in Pure and Applied Mathematics

POINCARÉ INEQUALITIES FOR THE HEISENBERG GROUP TARGET

GAO JIA

College of Science
University of Shanghai for Science and Technology
Shanghai 200093, China.

EMail: gaojia79@sina.com.cn

©2000 Victoria University
ISSN (electronic): 1443-5756
057-05



volume 6, issue 2, article 33,
2005.

*Received 24 November, 2004;
accepted 24 February, 2005.*

Communicated by: J. Sándor

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

In this paper some Poincaré type inequalities are obtained for the maps of the Heisenberg group target.


2000 Mathematics Subject Classification: 22E20, 26D10, 46E35.

Key words: Heisenberg group, Sobolev space, Poincaré type inequality.

Foundation item: the National Natural Science Foundation of China (19771048).

Contents

1	Introduction and Preliminaries	3
2	The Poincaré Type Inequalities for the Heisenberg Group Target	6
	References	



Poincaré Inequalities for the Heisenberg Group Target

Gao Jia

Title Page

Contents



Go Back

Close

Quit

Page 2 of 10

1. Introduction and Preliminaries

Let \mathbf{H}^m ([1]) denote a Heisenberg group which is a Lie group that has algebra $\mathfrak{g} = \mathbb{R}^{2m+1}$, with a non-abelian group law:

$$(1.1) \quad (x_1, y_1, t_1) \cdot (x_2, y_2, t_2) = (x_1 + x_2, y_1 + y_2, t_1 + t_2 + 2(y_2x_1 - x_2y_1)),$$

for every $u_1 = (x_1, y_1, t_1), u_2 = (x_2, y_2, t_2) \in \mathbf{H}^m$. The Lie algebra is generated by the left invariant vector fields

$$(1.2) \quad X_i = \frac{\partial}{\partial x_i} + 2y_i \frac{\partial}{\partial t}, \quad Y_i = \frac{\partial}{\partial y_i} - 2x_i \frac{\partial}{\partial t}, \quad i = 1, 2, \dots, m,$$


and $T = \frac{\partial}{\partial t}$. For every $u_1 = (x_1, y_1, t_1), u_2 = (x_2, y_2, t_2) \in \mathbf{H}^m$, the metric $d(u_1, u_2)$ in the Heisenberg group \mathbf{H}^m is defined as ([2])

$$(1.3) \quad \begin{aligned} d(u_1, u_2) &= |u_2 u_1^{-1}| \\ &= [((x_2 - x_1)^2 + (y_2 - y_1)^2)^2 \\ &\quad + (t_2 - t_1 + 2(x_2y_1 - x_1y_2))^2]^{\frac{1}{4}}. \end{aligned}$$

We see that \mathbf{H}^m possesses the nonlinear structure of group laws. It is one of the differences between \mathbf{H}^m and general Riemann manifolds.

Let $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) be a bounded and connected Lipschitz domain . Let $2 \leq p < \infty$.

L. Capogna and Fang-Hua Lin [3] have provided the characterizations for the Sobolev space $W^{1,p}(\Omega, \mathbf{H}^m)$, proved the existence theorem for the minimizer, and established that all critical points for the energy are Lipschitz continuous in



Title Page

Contents



Go Back

Close

Quit

Page 3 of 10

the 2-dimensional case. However, the higher dimensional regularity problem is still open.

In this paper, we shall give some Poincaré type inequalities for the maps of the Heisenberg group target. The statements of these results are similar to the ones in the classical case. However, since the metric possesses the nonlinear structure of the group law, we require the use of a few techniques in the proofs for our Poincaré type inequalities.

Definition 1.1. Let $2 \leq p < \infty$. A function $u = (z, t) : \Omega \rightarrow \mathbf{H}^m$ is in $L^p(\Omega, \mathbf{H}^m)$ if for some $h_0 \in \Omega$, one has

$$(1.4) \quad \int_{\Omega} (d(u(h), u(h_0)))^p dh < \infty.$$

A function $u = (z, t) : \Omega \rightarrow \mathbf{H}^m$ is in the Sobolev space $W^{1,p}(\Omega, \mathbf{H}^m)$ if $u \in L^p(\Omega, \mathbf{H}^m)$ and

$$(1.5) \quad E_{p,\Omega}(u) = \sup_{f \in C_c(\Omega), 0 \leq f \leq 1} \limsup_{\epsilon \rightarrow 0} \int_{\Omega} f(h) e_{u,\epsilon}(h) dh < \infty,$$


where

$$e_{u,\epsilon}(h) = \int_{|h-q|=\epsilon} \left(\frac{d(u(h), u(q))}{\epsilon} \right)^p \frac{d\sigma_{\epsilon}(q)}{\epsilon^{n-1}}.$$

$E_{p,\Omega}(u)$ is called the p -energy of u on Ω .

Lemma 1.1. If $u = (x, y, t) \in W^{1,p}(\Omega, \mathbf{H}^m)$, then

$$(1.6) \quad \nabla t = 2(y \nabla x - x \nabla y) \quad \text{in} \quad L^{\frac{p}{2}}(\Omega).$$



Poincaré Inequalities for the Heisenberg Group Target

Gao Jia

Title Page

Contents



Go Back

Close

Quit

Page 4 of 10

The maps satisfying (1.6) are called Legendrian maps.

Lemma 1.2. *If $u = (z, t) = (x, y, t) \in W^{1,p}(\Omega, \mathbf{H}^m)$, then*

$$E_{p,\Omega}(u) = \omega_{n-1} \int_{\Omega} |\nabla z|^p(q) dq.$$

Lemma 1.1 and Lemma 1.2 are due to L. Capogna and Fang-Hua Lin [3].

Lemma 1.3 ([4]). (*C_p-inequality*) *Let $p > 0$. Then for any $a_i \in \mathbb{R}$,*


$$\left(\sum_{i=1}^n |a_i| \right)^p \leq C_p \sum_{i=1}^n |a_i|^p,$$

where $C_p = 1$ if $0 < p < 1$ and $C_p = n^{p-1}$ if $p \geq 1$.

Lemma 1.4 ([5]). (*Poincaré Inequality in the classical case*) *Let Ω be a bounded and connected Lipschitz domain in \mathbb{R}^m . Let $p > 1$. Then there exists a constant C depending only on Ω , m and p , such that for every function $u \in W^{1,p}(\Omega, \mathbb{R})$, we have*

$$\int_{\Omega} |u(x) - \lambda_u|^p dx \leq C \int_{\Omega} |\nabla u|^p dx,$$

where $\lambda_u = \frac{1}{|\Omega|} \int_{\Omega} u(x) dx$.



Poincaré Inequalities for the
Heisenberg Group Target

Gao Jia

Title Page

Contents



Go Back

Close

Quit

Page 5 of 10

2. The Poincaré Type Inequalities for the Heisenberg Group Target

Theorem 2.1 (Poincaré type inequality). *Let Ω be a bounded and connected Lipschitz domain in \mathbb{R}^n . Then there exists a constant C depending only on Ω , n , m and p , such that for every function $u = (x, y, t) = (z, t) \in W^{1,p}(\Omega, \mathbf{H}^m)$,*

$$(2.1) \quad \int_{\Omega} (d(u(q), \lambda_u))^p dq \leq C_{\Omega} E_{p,\Omega}(u) = C_{\Omega} \int_{\Omega} |\nabla z|^p(q) dq.$$


Here $\lambda_u = (\lambda_x, \lambda_y, \lambda_t)$ and $\lambda_f = \frac{1}{|\Omega|} \int_{\Omega} f(q) dq$.

Proof. Obviously, $\lambda_u \in W^{1,p}(\Omega, \mathbf{H}^m)$. From (1.3), using the C_p -inequality, we have

$$(2.2) \quad \begin{aligned} & (d(u(q), \lambda_u))^p \\ &= [|z(q) - \lambda_z|^4 + (t(q) - \lambda_t + 2(\lambda_x y(q) - \lambda_y x(q)))^2]^{\frac{p}{4}} \\ &\leq C_p \left[|x(q) - \lambda_x|^p + |y(q) - \lambda_y|^p \right. \\ &\quad \left. + |t(q) - \lambda_t + 2(\lambda_x y(q) - \lambda_y x(q))|^{\frac{p}{2}} \right], \end{aligned}$$

where C_p depends on p . By the Poincaré inequality in the classical case, noting that

$$2(\lambda_x y(q) - \lambda_y x(q)) = 2\lambda_x(y(q) - \lambda_y) - 2\lambda_y(x(q) - \lambda_x),$$



Poincaré Inequalities for the Heisenberg Group Target

Gao Jia

Title Page

Contents



Go Back

Close

Quit


Page 6 of 10

we obtain

$$\begin{aligned} & \int_{\Omega} (d(u(q), \lambda_u))^p dq \\ & \leq C_p \left[\int_{\Omega} \left(|x(q) - \lambda_x|^p + |y(q) - \lambda_y|^p + |t(q) - \lambda_t + 2(\lambda_x y(q) - \lambda_y x(q))|^{\frac{p}{2}} \right) dq \right] \\ & \leq C_1 \int_{\Omega} |\nabla x|^p(q) dq + C_2 \int_{\Omega} |\nabla y|^p(q) dq + C_3 \int_{\Omega} |\nabla t + 2(\lambda_x \nabla y - \lambda_y \nabla x)|^{\frac{p}{2}} dq. \end{aligned}$$

By virtue of the Legendrian condition $\nabla t = 2(y\nabla x - x\nabla y)$, using the Hölder inequality, noting that $|\nabla x| \leq |\nabla z|$ and $|\nabla y| \leq |\nabla z|$, we have

$$\begin{aligned} & \int_{\Omega} (d(u(q), \lambda_u))^p dq \\ & \leq C_1 \int_{\Omega} |\nabla x|^p dq + C_2 \int_{\Omega} |\nabla y|^p(q) \\ & \quad + C_3 2^{\frac{p}{2}} \int_{\Omega} |\nabla y(x - \lambda_x) - \nabla x(y - \lambda_y)|^{\frac{p}{2}} dq \\ & \leq C_1 \int_{\Omega} |\nabla x|^p dq + C_2 \int_{\Omega} |\nabla y|^p dq \\ & \quad + C_4 \left(\int_{\Omega} |\nabla x|^{\frac{p}{2}} |y - \lambda_y|^{\frac{p}{2}} dq + \int_{\Omega} |\nabla y|^{\frac{p}{2}} |x - \lambda_x|^{\frac{p}{2}} dq \right) \\ & \leq C_1 \int_{\Omega} |\nabla x|^p dq + C_2 \int_{\Omega} |\nabla y|^p dq + C_5 \left(\int_{\Omega} |\nabla x|^p dq \int_{\Omega} |\nabla y|^p dq \right)^{\frac{1}{2}} \end{aligned}$$



Poincaré Inequalities for the Heisenberg Group Target

Gao Jia

Title Page

Contents




Go Back

Close

Quit

Page 7 of 10



Title Page

Contents



Go Back

Close

Quit

Page 8 of 10

$$\begin{aligned} &\leq C_1 \int_{\Omega} |\nabla x|^p dq + C_2 \int_{\Omega} |\nabla y|^p dq + C_6 \left(\int_{\Omega} |\nabla x|^p dq + \int_{\Omega} |\nabla y|^p dq \right) \\ &\leq C \int_{\Omega} |\nabla z|^p(q) dq, \end{aligned}$$

where $C_1, C_2, C_3, C_4, C_5, C_6$ and C are dependent on Ω, n, m and p . □

Corollary 2.2. *If $u \in W^{1,p}(B(h_0, r), \mathbf{H}^m)$, then*

$$\begin{aligned} (2.3) \quad \int_{B(h_0, r)} (d(u(q), \lambda_u))^p dq &\leq Cr^p E_{p, B(h_0, r)}(u) \\ &= Cr^p \int_{B(h_0, r)} |\nabla z|^p(q) dq. \end{aligned}$$

Proof. Observe that

$$\begin{aligned} &(d(u(q), \lambda_u))^p \\ &= [|z(q) - \lambda_z|^4 + (t(q) - \lambda_t + 2(\lambda_x y(q) - \lambda_y x(q)))^2]^{\frac{p}{4}} \\ &\leq C_p \left[|x(q) - \lambda_x|^p + |y(q) - \lambda_y|^p + |t(q) - \lambda_t + 2(\lambda_x y(q) - \lambda_y x(q))|^{\frac{p}{2}} \right]. \end{aligned}$$

Here C_p depends on p . By the Poincaré inequality in the classical case, noting that

$$2(\lambda_x y(q) - \lambda_y x(q)) = 2\lambda_x(y(q) - \lambda_y) - 2\lambda_y(x(q) - \lambda_x),$$

we deduce


$$\int_{B_r(h_0)} (d(u(q), \lambda_u))^p dq$$

$$\begin{aligned}
&\leq C_p \left[\int_{B_r(h_0)} (|x - \lambda_x|^p + |y - \lambda_y|^p + |t - \lambda_t + 2(\lambda_x y - \lambda_y x)|^{\frac{p}{2}}) dq \right] \\
&\leq C_1 r^p \int_{B_r(h_0)} |\nabla x|^p(q) dq + C_2 r^p \int_{B_r(h_0)} |\nabla y|^p(q) dq \\
&\quad + C_3 r^{\frac{p}{2}} \int_{B_r(h_0)} |\nabla t + 2(\lambda_x \nabla y - \lambda_y \nabla x)|^{\frac{p}{2}} dq.
\end{aligned}$$

By virtue of the Legendrian condition $\nabla t = 2(y\nabla x - x\nabla y)$, using Hölder's inequality, noting that $|\nabla x| \leq |\nabla z|$ and $|\nabla y| \leq |\nabla z|$, we can obtain

$$\begin{aligned}
&\int_{B_r(h_0)} (d(u(q), \lambda_u))^p dq \\
&\leq C_1 r^p \int_{B_r(h_0)} |\nabla x|^p(q) dq + C_2 r^p \int_{B_r(h_0)} |\nabla y|^p(q) dq \\
&\quad + C_3 r^{\frac{p}{2}} \int_{B_r(h_0)} |\nabla y(q)(x(q) - \lambda_x) - \nabla x(q)(y(q) - \lambda_y)|^{\frac{p}{2}} dq \\
&\leq C_1 r^p \int_{B_r(h_0)} |\nabla x|^p(q) dq + C_2 r^p \int_{B_r(h_0)} |\nabla y|^p(q) dq \\
&\quad + C_4 r^p \left(\int_{B_r(h_0)} |\nabla x|^p dq \int_{B_r(h_0)} |\nabla y|^p dq \right)^{\frac{1}{2}} \\
&\leq C r^p \int_{\Omega} |\nabla z|^p(q) dq,
\end{aligned}$$

where C_1, C_2, C_3, C_4 and C depend on Ω, n, m and p . □



Poincaré Inequalities for the Heisenberg Group Target

Gao Jia

Title Page

Contents



Go Back


Close

Quit

Page 9 of 10

References

- [1] D. CHRISTODOULOU, On the geometry and dynamics of crystalline continua, *Ann. Inst. H. Poincaré*, **13** (1998), 335–358.
- [2] A. VENERUSO, Hömander multipliers on the Heisenberg group, *Monatsh. Math.*, **130** (2000), 231–252.
- [3] L. CAPOGNA AND FANG-HUA LIN, Legendrian energy minimizers, Part I: Heisenberg group target, *Calc. Var.*, **12** (2001), 145–171.
- [4] B.F. BECKENBACH AND R. BELLMAN, *Inequalities*, Springer-Verlag, 1961.
- [5] W.P. ZIEMER, *Weakly Differentiable Function*, Springer-Verlag, New York, 1989.
- [6] Z.Q. WANG AND M. WILLEM, Singular minimization problems, *J. Diff. Equa.*, **161** (2000), 307–370.



Poincaré Inequalities for the Heisenberg Group Target

Gao Jia

Title Page

Contents



Go Back

Close

Quit

Page 10 of 10