

# Journal of Inequalities in Pure and Applied Mathematics

## A NOTE ON AN INDIVIDUAL BIOEQUIVALENCE SETTING


C. SURULESCU AND N. SURULESCU

Institut für Angewandte Mathematik  
INF 294, 69120 Heidelberg  
Germany.

*E-Mail:* [Christina.Surulescu@iwr.uni-heidelberg.de](mailto:Christina.Surulescu@iwr.uni-heidelberg.de)

*E-Mail:* [Nicolae.Surulescu@iwr.uni-heidelberg.de](mailto:Nicolae.Surulescu@iwr.uni-heidelberg.de)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
079-06



---

volume 7, issue 3, article 100,  
2006.

*Received 07 April, 2005;*  
*accepted 07 April, 2006.*

*Communicated by: S.S. Dragomir*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract


We give a new simpler proof along with a generalization for the inequality of Yao and Iyer [10] arising in bioequivalence studies and by using a nonparametric approach we also discuss an extension of the individual bioequivalence setting to the case where the data are not necessarily normally distributed.

*2000 Mathematics Subject Classification:* 60E15, 62F03, 62P10.

*Key words:* Bioequivalence, Probabilistic inequalities

## Contents

1	Introduction .....	3
2	A Generalization of the Inequality of Yao and Iyer .....	6
3	A Nonparametric Approach for the Bioequivalence Setting ..	10
4	Conclusions .....	15
A	Proofs .....	16
References		



### A Note on an Individual Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 2 of 21


# 1. Introduction

Bioequivalence testing is required when trying to get the approval for manufacturing and selling of a generic drug having mainly the same properties with a (more expensive) reference (brand-name) drug. Establishing bioequivalence saves the generic drug manufacturer from performing expensive clinical trials to demonstrate the quality of his product. Two drugs are considered bioequivalent if they are absorbed into the blood and become active at about the same rate and concentration. Bioequivalent drugs are supposed to provide the same therapeutic effect.

For explaining the notions and notations we use, we recall the problem setting in [10] for the individual bioequivalence. Thus, the amount of the chemical absorbed by a patient's bloodstream when using a reference drug is a random variable  $X$ , which has mean  $\mu_R$  and standard deviation  $\sigma_R$ . The corresponding variable for the same patient when using the generic drug has mean  $\mu_T$  and standard deviation  $\sigma_T$ . The *therapeutic window* of a patient is defined to be the interval in which must lie the concentration of the chemical in the bloodstream, in order for the drug to be classified as beneficial for that patient. Usually, the therapeutic window is assumed to be an interval centered at the mean  $\mu_R$ , namely the range  $(\mu_R - z\sigma_R, \mu_R + z\sigma_R)$ . The drug will be ineffective if the amount absorbed in the bloodstream is too low and it could cause severe side effects if too much of the chemical substance is absorbed.

Denoting by  $p_R$  and  $p_T$  the probabilities that the subject will have benefit from using drug  $R$ , respectively  $T$ , the regulatory agency might approve the marketing of drug  $T$  provided that  $\frac{p_T}{p_R} \geq \gamma$ , where  $\gamma$  is about 1 or even larger.

The therapeutic window of a patient is generally unknown, therefore a quan-



---

A Note on an Individual  
Bioequivalence Setting

C. Surulescu and N. Surulescu

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 21

tity of interest for the approval procedure will be  $\inf_{z \geq 0} \frac{p_T}{p_R}$ .

Usually, to derive this quantity the assumption that  $X$  and  $T$  have normal distributions is made, though it is well known that in practice this is rarely the case. Under this assumption, Yao and Iyer [10] have shown that

$$(1.1) \quad \inf_{z > 0} \frac{p_T}{p_R} = \inf_{z > 0} \frac{\Phi\left(\frac{\mu_R + z\sigma_R - \mu_T}{\sigma_T}\right) - \Phi\left(\frac{\mu_R - z\sigma_R - \mu_T}{\sigma_T}\right)}{\Phi(z) - \Phi(-z)} \\ = \min \left\{ 1, \frac{\sigma_R \sqrt{2\pi}}{\sigma_T} \phi\left(\frac{\mu_T - \mu_R}{\sigma_T}\right) \right\},$$

where  $\phi$  and  $\Phi$  are respectively the probability density function and the cumulative density function of a standard normal variable.


Thus, the approval of manufacturing the generic drug  $T$  may be granted if from the statistical analysis of experimental data it can be proven that

$$(1.2) \quad \frac{\sigma_R \sqrt{2\pi}}{\sigma_T} \phi\left(\frac{\mu_T - \mu_R}{\sigma_T}\right) \geq \gamma.$$

Alternatively to (1.2), a more flexible approval criterion can be used, namely one of the type

$$(1.3) \quad \ell(\mathcal{Z}_\gamma) \geq b_\gamma,$$

for some large enough given bound  $b_\gamma$ , where  $\mathcal{Z}_\gamma := \{z > 0 : \frac{p_T}{p_R} > \gamma\}$  and  $\ell$  is for instance the Lebesgue measure. We will discuss in the next sections sufficient conditions for (1.3) to be satisfied.



A Note on an Individual  
Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close


Quit

Page 4 of 21

For more information about individual bioequivalence see [1], [7], [8], [4], [5], [6] and the references therein.

In this paper we give a simpler proof and a generalization of the inequality of Yao and Iyer and we also give a nonparametric extension of the above bioequivalence setting.

For the sake of clarity, we put all our proofs in the Appendix.



---

## A Note on an Individual Bioequivalence Setting

C. Surulescu and N. Surulescu

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 21

## 2. A Generalization of the Inequality of Yao and Iyer

The main result of Yao and Iyer in [10] was the proof of the following inequality:

$$(2.1) \quad \frac{\Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right)}{\Phi(z) - \Phi(-z)} > \min \left\{ 1, \frac{\sqrt{2\pi}}{\sigma} \phi\left(\frac{\mu}{\sigma}\right) \right\},$$


for all  $z > 0$ ,  $\mu \in \mathbb{R} \setminus \{0\}$  and  $\sigma \in (0, \infty) \setminus \{1\}$ ; this inequality comes from (1.1) after some changes of notations.

As in [10], observe that it is enough to treat the case  $\mu \geq 0$ . Here we will prove the following generalization:

**Proposition 2.1.** *In the above settings, we have:*

(i) *If  $\sigma > 1$ , then*

$$\begin{aligned} \frac{\Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right)}{\Phi(z) - \Phi(-z)} &> e^{-\frac{\mu^2}{2\sigma^2}} \frac{\Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right)}{\Phi(z) - \Phi(-z)} \\ &> e^{-\frac{\mu^2}{2\sigma^2}} \frac{1}{\sigma} \left[ 1 + \frac{1}{6} \left( 1 - \frac{1}{\sigma^2} \right) \cdot z^2 e^{-\frac{z^2}{2}} \right] \\ &> \frac{1}{\sigma} e^{-\frac{\mu^2}{2\sigma^2}} = \min \left\{ 1, \frac{\sqrt{2\pi}}{\sigma} \phi\left(\frac{\mu}{\sigma}\right) \right\}, \quad \forall z > 0. \end{aligned}$$



A Note on an Individual  
Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 6 of 21

(ii) If  $\sigma \in (0, 1)$  then

$$(2.2) \quad \frac{\Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right)}{\Phi(z) - \Phi(-z)} \geq \min \left\{ 1, \frac{\sqrt{2\pi}}{\sigma} \phi\left(\frac{\mu}{\sigma}\right) \right\} \frac{\Phi\left(\frac{z-\mu(\sigma)}{\sigma}\right) - \Phi\left(\frac{-z-\mu(\sigma)}{\sigma}\right)}{\Phi(z) - \Phi(-z)}$$

$$> \min \left\{ 1, \frac{\sqrt{2\pi}}{\sigma} \phi\left(\frac{\mu}{\sigma}\right) \right\}, \quad \forall z > 0, \forall \mu > 0,$$

where  $\mu(\sigma) := \sigma \sqrt{2 \ln \frac{1}{\sigma}}$ .

This result is based on Lemma 2.2, generalising Lemma 3 in [10], (which was the most difficult part of the proof therein) and on Lemmas 2.3 and 2.4 below.


**Lemma 2.2.** *The function  $\sigma \mapsto F(\sigma) := \Phi\left(\frac{z-\mu(\sigma)}{\sigma}\right) - \Phi\left(\frac{-z-\mu(\sigma)}{\sigma}\right)$  is strictly decreasing on  $(0, 1)$ , with  $\mu(\sigma)$  as above.*

**Lemma 2.3.** *For every  $\sigma > 0$ ,  $\sigma \neq 1$ , we have:*

$$(2.3) \quad \frac{\Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right)}{\Phi(z) - \Phi(-z)} - \min \left\{ 1, \frac{1}{\sigma} \right\} > \frac{1}{6\sigma} \left( 1 - \min \left\{ 1, \frac{1}{\sigma} \right\}^2 \right) \cdot z^2 e^{-\frac{z^2}{2}}$$

$$+ \left( \frac{1}{\sigma} - \min \left\{ 1, \frac{1}{\sigma} \right\} \right) e^{-\frac{z^2}{2\sigma^2}} > 0, \quad \forall z > 0.$$

The following lemma generalises Lemma 1 and Lemma 4 in [10]:



A Note on an Individual Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 7 of 21

**Lemma 2.4.**

$$(2.4) \quad \frac{\Phi(z - \mu_1) - \Phi(-z - \mu_1)}{\Phi(z - \mu_2) - \Phi(-z - \mu_2)} > \min \left\{ 1, e^{-\frac{1}{2}[(\mu_1)^2 - (\mu_2)^2]} \right\}$$

$\forall z > 0, \mu_1, \mu_2 \in \mathbb{R}, \mu_1 \neq \mu_2.$

**Remark 1.** Replacing  $z$  with  $\frac{z}{\sigma}$ ,  $\mu_1$  by  $\frac{\mu_1}{\sigma}$  and  $\mu_2$  by  $\frac{\mu_2}{\sigma}$  then

- for  $\mu_1 > \mu_2 > 0$  one has

$$\frac{\Phi\left(\frac{z-\mu_1}{\sigma}\right) - \Phi\left(\frac{-z-\mu_1}{\sigma}\right)}{\Phi\left(\frac{z-\mu_2}{\sigma}\right) - \Phi\left(\frac{-z-\mu_2}{\sigma}\right)} > \min \left\{ 1, e^{-\frac{1}{2}\left[\left(\frac{\mu_1}{\sigma}\right)^2 - \left(\frac{\mu_2}{\sigma}\right)^2\right]} \right\} = e^{-\frac{1}{2}\left[\left(\frac{\mu_1}{\sigma}\right)^2 - \left(\frac{\mu_2}{\sigma}\right)^2\right]},$$

thus the function

$$(0, \infty) \ni \mu \mapsto B_1(\mu) := e^{\frac{1}{2}\left(\frac{\mu}{\sigma}\right)^2} \left[ \Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right) \right]$$

is increasing, i.e. Lemma 1 in [10] is obtained;


- for  $\mu_2 > \mu_1 > 0$  one has

$$\frac{\Phi\left(\frac{z-\mu_1}{\sigma}\right) - \Phi\left(\frac{-z-\mu_1}{\sigma}\right)}{\Phi\left(\frac{z-\mu_2}{\sigma}\right) - \Phi\left(\frac{-z-\mu_2}{\sigma}\right)} > 1,$$

thus the function

$$(0, \infty) \ni \mu \mapsto B_2(\mu) := \Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right)$$

is proved to be decreasing, i.e. we obtained Lemma 4 in [10].



**A Note on an Individual  
Bionequivalence Setting**

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit


Page 8 of 21



**Remark 2.** Sufficient conditions for (1.3) to be satisfied can be easily derived upon using for  $\sigma > 1$  the monotonicity of the function  $z \mapsto z^2 e^{-\frac{z^2}{2}}$  (increasing on  $(0, \sqrt{2})$ , decreasing on  $[\sqrt{2}, \infty)$ ) and for  $0 < \sigma < 1$  the fact that

$$\begin{aligned} \frac{\Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right)}{\Phi(z) - \Phi(-z)} &> \min \left\{ 1, \frac{\sqrt{2\pi}}{\sigma} \phi\left(\frac{\mu}{\sigma}\right) \right\} \frac{\Phi\left(\frac{z-\mu(\sigma)}{\sigma}\right) - \Phi\left(\frac{-z-\mu(\sigma)}{\sigma}\right)}{\Phi(z) - \Phi(-z)} \\ &\geq \sigma \min \left\{ 1, \frac{\sqrt{2\pi}}{\sigma} \phi\left(\frac{\mu}{\sigma}\right) \right\} \frac{\Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right)}{\Phi(z) - \Phi(-z)}, \quad \forall z > 0 \end{aligned}$$

and this term can be minorated by applying Lemma 2.3.



A Note on an Individual  
Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 9 of 21

### 3. A Nonparametric Approach for the Bioequivalence Setting

Consider now that the random variables  $X, T$  from the Introduction have continuous univariate distributions, with the densities  $f_X$ , respectively  $f_T$ , which are not necessarily Gaussian, and assume that we correspondingly have the independent observations:  $x_1, \dots, x_m$ , respectively  $t_1, \dots, t_n$ . Then  $f_X$  and  $f_T$  can be estimated by using the classical nonparametric estimators:

$$\hat{f}_X(x) = \frac{1}{mh_m^X} \cdot \sum_{i=1}^m K\left(\frac{x - x_i}{h_m^X}\right)$$


and

$$\hat{f}_T(t) = \frac{1}{nh_n^T} \cdot \sum_{i=1}^n K\left(\frac{t - t_i}{h_n^T}\right),$$

where  $K$  is a kernel,  $h_m^X$  and  $h_n^T$  are the bandwidths (with the usual properties:  $h_m^X \rightarrow 0$ ,  $h_n^T \rightarrow 0$  for  $m, n \rightarrow \infty$ ;  $mh_m^X \rightarrow 0$ ,  $nh_n^T \rightarrow \infty$  for  $m, n \rightarrow \infty$  and, of course,  $h_m^X$  and  $h_n^T$  have to be chosen in practice by a corresponding criterion, see [9]).

With the above notations, the fractions of interest for bioequivalence studies are

$$(3.1) \quad \frac{p_T}{p_R} = R(z) := \frac{\int_{\mu_R - z\sigma_R}^{\mu_R + z\sigma_R} f_T(t) dt}{\int_{\mu_R - z\sigma_R}^{\mu_R + z\sigma_R} f_X(x) dx}$$



A Note on an Individual Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit


Page 10 of 21

$$\approx \hat{R}(z) := \frac{\int_{\mu_R - z\sigma_R}^{\mu_R + z\sigma_R} \hat{f}_T(t) dt}{\int_{\mu_R - z\sigma_R}^{\mu_R + z\sigma_R} \hat{f}_X(x) dx}, \quad z > 0,$$

and for the approval procedure it will be important to find the quantity of interest  $\inf_{z>0} \hat{R}(z)$ , but this is clearly more difficult than for the case presented in the previous section and an analytic treatment is hardly possible. Even when considering the Gaussian kernel, the available nonlinear optimization procedures are surprisingly very time consuming. However, a great (and also easy to implement) simplification can be achieved when using one of the following kernels (see e.g., [9]):

- the rectangular kernel:  $K(u) := \frac{1}{2}\chi_{(-1,1)}(u)$ ;
- the triangular kernel:  $K(u) := (1 - |u|)\chi_{(-1,1)}(u)$ ;
- the Epanechnikov kernel:  $K(u) := \frac{3}{4}(1 - u^2)\chi_{(-1,1)}(u)$ ;
- the triangle kernel:  $K(u) := (1 - |u|)\chi_{(-1,1)}(u)$ ;
- the double Epanechnikov kernel:  $K(u) := 3|u|(1 - |u|)\chi_{(-1,1)}(u)$ .

Moreover, the procedure can also be used for finding the largest therapeutic window under which the fraction of interest exceeds some given positive constant  $\gamma$ . Since all kernels above are supported on  $(-1, 1)$ , then the global minimum can be found in the following way:



### A Note on an Individual Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 11 of 21

- construct  $A_n^{T,+} := A_n^T \cap (0, \infty)$ , where


$$A_n^T := \bigcup_{i=1}^n \left\{ -\frac{t_i - h_n^T - \mu_R}{\sigma_R}, \frac{t_i + h_n^T - \mu_R}{\sigma_R}, \frac{t_i - h_n^T - \mu_R}{\sigma_R}, -\frac{t_i + h_n^T - \mu_R}{\sigma_R} \right\};$$

- construct  $A_m^{X,+} := A_m^X \cap (0, \infty)$ , where

$$A_m^X := \bigcup_{j=1}^m \left\{ -\frac{x_j - h_m^X - \mu_R}{\sigma_R}, \frac{x_j + h_m^X - \mu_R}{\sigma_R}, \frac{x_j - h_m^X - \mu_R}{\sigma_R}, -\frac{x_j + h_m^X - \mu_R}{\sigma_R} \right\};$$

- if  $K$  is not the rectangular kernel, then construct  $A_{T,deriv}^+$  as the set constituted by the critical points of  $\hat{R}(z)$  on the union of the subintervals of  $\mathbb{R}_+$  where  $\hat{R}$  is differentiable (which in this case are roots of some polynomials, thus easy to be handled by the computer); if  $K$  is the rectangular kernel, set  $A_{T,deriv}^+ = \emptyset$ .
- denote  $\mathcal{A} := A_n^{T,+} \cup A_m^{X,+} \cup A_{T,deriv}^+$ .

Then we have:



**A Note on an Individual  
Bioequivalence Setting**

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 12 of 21

**Proposition 3.1.** *With the notations above and with  $K$  being one of the kernels enumerated before,*

$$\hat{R}(z) = \frac{\frac{1}{nh_n^T} \sum_{i=1}^n \int_{\mu_R - z\sigma_R}^{\mu_R + z\sigma_R} K\left(\frac{t-t_i}{h_n^T}\right) dt}{\frac{1}{mh_m^X} \sum_{i=1}^m \int_{\mu_R - z\sigma_R}^{\mu_R + z\sigma_R} K\left(\frac{x-x_i}{h_m^X}\right) dx} \geq \min \left\{ \hat{R}(z_{max}), \lim_{z \rightarrow 0} \hat{R}(z), \hat{R}(\mathcal{A}) \right\},$$

where  $z_{max} = \max\{|\zeta| : \zeta \in \mathcal{A}\}$ ,


$$\lim_{z \rightarrow 0} \hat{R}(z) = \frac{mh_m^X \sum_{i=1}^n K\left(\frac{\mu_R - t_i}{h_n^T}\right)}{nh_n^T \sum_{i=1}^m K\left(\frac{\mu_R - x_i}{h_m^X}\right)} \text{ for the continuous kernels above,}$$

and

$$\lim_{z \rightarrow 0} \hat{R}(z) = \frac{mh_m^X \sum_{i=1}^n \left[ K_+ \left( \frac{\mu_R - t_i}{h_n^T} \right) + K_- \left( \frac{\mu_R - t_i}{h_n^T} \right) \right]}{nh_n^T \sum_{i=1}^m \left[ K_+ \left( \frac{\mu_R - x_i}{h_m^X} \right) + K_- \left( \frac{\mu_R - x_i}{h_m^X} \right) \right]} \text{ for the rectangular kernel,}$$

with the notations  $K_+(\xi) := \lim_{t \searrow \xi} K(t)$ , respectively  $K_-(\xi) := \lim_{t \nearrow \xi} K(t)$ .

**Remark 3.** *If  $K$  is one of the kernels presented above, then for each  $\gamma > 0$  and using an algorithm similar to the one described above one can easily determine  $\mathcal{Z}_\gamma$ , in order to check the approval condition (1.3).*



**A Note on an Individual  
Bionequivalence Setting**

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 13 of 21

**Simulation result:** Figure 1 illustrates a simulation with  $m = 150$ ,  $n = 160$  for the case where  $X$  and  $T$  are normally distributed and the nonparametric estimators are constructed with the Epanechnikov kernel. The continuous lines represent  $\hat{R}(z)$  and its minimum, which is 0.012; the lines of circles represent  $R(z)$  and its minimum, which is 0.0088. This shows that the Gaussian case can be well recovered with this nonparametric procedure.





Figure 1: The continuous line is  $\hat{R}(z)$ ; the line of circles is  $R(z)$ ; the horizontal lines are the corresponding minimums



## A Note on an Individual Bioequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back


Close

Quit

Page 14 of 21

## 4. Conclusions

In this paper we gave a generalization and a simpler proof of the inequality of Yao and Iyer [10] concerning an individual bioequivalence setting when the data were supposed to be normally distributed. Our generalization can be used to develop a more flexible approval criterion (of the type (1.3)) for manufacturing and selling a drug which is supposed to be bioequivalent with a reference one. Finally, we extended these settings to the more general situation when the data are not necessarily normally distributed, upon using the nonparametric estimation technique. This idea can also be useful in the context of global nonlinear optimization problems.



---

### A Note on an Individual Bioequivalence Setting

C. Surulescu and N. Surulescu

---

Title Page

Contents



Go Back

Close

Quit

Page 15 of 21

## A. Proofs

*Proof of Lemma 2.2.* Denoting  $\theta := \frac{1}{\sigma} \in (1, \infty)$ , we have

$$\Phi\left(\frac{z - \mu(\sigma)}{\sigma}\right) - \Phi\left(\frac{-z - \mu(\sigma)}{\sigma}\right) = \Phi(z\theta - \sqrt{2\ln\theta}) - \Phi(-z\theta - \sqrt{2\ln\theta}).$$

Consider  $g : (1, \infty) \rightarrow \mathbb{R}$ ,  $g(\theta) := \Phi(z\theta - \sqrt{2\ln\theta}) - \Phi(-z\theta - \sqrt{2\ln\theta})$ ,  $\forall \theta >$

1. Observe that

$$g'(\theta) = \left(z + \frac{1}{\theta\sqrt{2\ln\theta}}\right) \cdot \phi(z\theta - \sqrt{2\ln\theta}) \left[\frac{z\theta\sqrt{2\ln\theta} - 1}{z\theta\sqrt{2\ln\theta} + 1} + \exp(-2z\theta\sqrt{2\ln\theta})\right] > 0,$$


for all  $\theta > 1$ , because  $u := z\theta\sqrt{2\ln\theta} > 0$  and it is easy to see that

$$\frac{u - 1}{u + 1} + e^{-2u} > 0, \forall u > 0.$$

□

*Proof of Lemma 2.3.* If  $\sigma > 1$ , we can write

$$\begin{aligned} \Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right) &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-z}^z e^{-\frac{x^2}{2\sigma^2}} dx \\ &> \frac{1}{\sigma} \left[ \Phi(z) - \Phi(-z) + \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{2} \left(1 - \frac{1}{\sigma^2}\right) \int_{-z}^z x^2 e^{-\frac{x^2}{2}} dx \right], \end{aligned}$$



A Note on an Individual  
Biequivalence Setting

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 16 of 21



since  $e^{\frac{x^2}{2}(1-\frac{1}{\sigma^2})} \geq 1 + \frac{x^2}{2} \left(1 - \frac{1}{\sigma^2}\right), \forall x \in \mathbb{R} \setminus \{0\}$ . Further, from the decreasing monotonicity of  $x \mapsto e^{-\frac{x^2}{2}}$  on  $[0, z]$  deduce that

$$\Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right) \geq \frac{1}{\sigma} \left[ \Phi(z) - \Phi(-z) + \frac{1}{\sqrt{2\pi}} \left(1 - \frac{1}{\sigma^2}\right) \cdot \frac{z^3}{3} e^{-\frac{z^2}{2}} \right]$$

and, since

$$\Phi(z) - \Phi(-z) < z\sqrt{2/\pi}, \quad \forall z > 0$$

(2.3) is proved.


For the case where  $\sigma \in (0, 1)$  we have:

$$\begin{aligned} \frac{\Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right)}{\Phi(z) - \Phi(-z)} &= 1 + 2 \frac{\Phi\left(\frac{z}{\sigma}\right) - \Phi(z)}{\Phi(z) - \Phi(-z)} \\ &= 1 + 2 \frac{\int_z^{\frac{z}{\sigma}} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx}{\Phi(z) - \Phi(-z)} \\ &> 1 + 2 \frac{\frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2\sigma^2}} \cdot z \left(\frac{1}{\sigma} - 1\right)}{\Phi(z) - \Phi(-z)} \\ &> 1 + \left(\frac{1}{\sigma} - 1\right) e^{-\frac{z^2}{2\sigma^2}}, \quad \forall z > 0, \end{aligned}$$

upon using the same method as in the previous case. □

*Proof of Lemma 2.4.* Let

$$\begin{aligned} Q(z) &:= \Phi(z - \mu_1) - \Phi(-z - \mu_1) \\ &\quad - e^{-\frac{1}{2}[(\mu_1)^2 - (\mu_2)^2]} [\Phi(z - \mu_2) - \Phi(-z - \mu_2)], \quad z > 0. \end{aligned}$$



**A Note on an Individual  
Bionequivalence Setting**

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 17 of 21

Then

$$\begin{aligned} Q'(z) &= \frac{2}{\sqrt{2\pi}} e^{-\frac{1}{2}[(\mu_1)^2+z^2]} \left[ \frac{e^{z\mu_1} + e^{-z\mu_1}}{2} - \frac{e^{z\mu_2} + e^{-z\mu_2}}{2} \right] \\ &= \frac{2}{\sqrt{2\pi}} e^{-\frac{1}{2}[(\mu_1)^2+z^2]} [\cosh(z|\mu_1|) - \cosh(z|\mu_2|)], \quad \forall z > 0. \end{aligned}$$

Now using the fact that the hyperbolic cosine is an increasing function on  $(0, \infty)$ , we have that  $Q'(z) > 0$  if  $|\mu_1| > |\mu_2|$  and  $Q'(z) < 0$  if  $|\mu_1| < |\mu_2|$ .

Thus, if  $|\mu_1| > |\mu_2|$ , then  $Q(z) > 0, \forall z > 0$ , i.e. inequality (2.4) is satisfied, since the minimum therein is  $e^{-\frac{1}{2}[(\mu_1)^2-(\mu_2)^2]}$ .

If  $|\mu_1| < |\mu_2|$ , then consider the function

$$G(\mu) := \Phi(z - \mu) - \Phi(-z - \mu)$$


and observe that it is decreasing on  $(0, \infty)$ , since its derivative is

$$G'(\mu) = -\frac{1}{\sqrt{2\pi}} e^{-(z^2+\mu^2)/2} [e^{z\mu} - e^{-z\mu}] < 0, \quad \forall z > 0, \mu > 0.$$

Observing that  $G(\mu) = G(-\mu), \forall \mu \in \mathbb{R}$ , we have that  $G(\mu_2) = G(|\mu_2|) < G(|\mu_1|) = G(\mu_1)$ , which proves the inequality in this case.  $\square$

*Proof of Proposition 2.1.* In the case (i) observe that we can write

$$\frac{\Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right)}{\Phi(z) - \Phi(-z)} = \frac{\Phi\left(\frac{z-\mu}{\sigma}\right) - \Phi\left(\frac{-z-\mu}{\sigma}\right)}{\Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right)} \cdot \frac{\Phi\left(\frac{z}{\sigma}\right) - \Phi\left(\frac{-z}{\sigma}\right)}{\Phi(z) - \Phi(-z)}.$$



**A Note on an Individual  
Biequivalence Setting**

C. Surulescu and N. Surulescu

Title Page

Contents



Go Back

Close

Quit

Page 18 of 21

Now apply Lemma 2.4 for the first term in the right hand side and Lemma 2.3 for the other one.


In the case (ii), if  $\mu \in (0, \mu(\sigma))$ , then we have that  $B_2(\mu) \geq B_2(\mu(\sigma))$  (see Lemma 2.4 and Remark 1). Further, use Lemma 2.2 to obtain the last inequality in (2.2).

If  $\mu > \mu(\sigma)$ , then  $\sigma B_1(\mu) \geq \sigma B_1(\mu(\sigma))$  (see again Lemma 2.4 and Remark 1). Then Lemma 2.2 implies that

$$\frac{\sigma B_1(\mu(\sigma))}{\Phi(z) - \Phi(-z)} = \frac{F(\sigma)}{\Phi(z) - \Phi(-z)} > 1,$$

which completes the proof. □

*Proof of Proposition 3.1.* The proof follows observing that for each of the kernels above  $\hat{R}(z)$  becomes a rapport of polynomials on some finite intervals dictated by the points in  $A_m^{X,+} \cup A_n^{T,+}$ . Thus, the problem reduces to characterising the minimum of these fractional expressions on such corresponding finite intervals, which in this particular case means to find the roots of the polynomials at the numerator of the derivatives. It only remains to observe that  $\lim_{z \rightarrow \infty} \hat{R}(z) = \hat{R}(z_{max})$ . □




---

**A Note on an Individual  
Bioequivalence Setting**

C. Surulescu and N. Surulescu

---

Title Page

Contents



Go Back

Close

Quit

Page 19 of 21

## References

- [1] S. ANDERSON AND W.W. HAUCK, Consideration of individual bioequivalence, *J. of Pharmacokinetics and Biopharmaceutics*, **18** (1990), 259–274.
- [2] I.G. GRAMA, On moderate deviations for martingales, *The Annals of Probability*, **25**(1) (1997), 152–183.
- [3] B. JONES AND M.G. KENWARD, *Design and Analysis of Cross-Over Trials* (2nd edition), Chapman & Hall, 2003.
- [4] B.F.J. MANLY AND R.I.C.C. FRANCIS, Testing for mean and variance differences with samples from distributions that may be non-normal with unequal variances, *Journal of Statistical Computation and Simulation*, **72**(8) (2002), 633–646.
- [5] B.F.J. MANLY, One-sided tests of bioequivalence with nonnormal distributions and unequal variances, *Journal of Agricultural, Biological and Environmental Statistics*, **9**(3) (2004), 270–283.
- [6] L.L.MCDONALD, S. HOWLIN, J. POLYAKOVA AND C.J. BILBROUGH, *Evaluation and Comparison of Hypothesis Testing Techniques for Bond Release Applications*, a project for the Abandoned Coal Mine Lands Research Program at the University of Wyoming, 2003.
- [7] R. SCHALL, Assessment of individual and population bioequivalence using the probability that bioavailabilities are similar, *Biometrics*, **51** (1995), 615–626.



---

A Note on an Individual  
Bioequivalence Setting

C. Surulescu and N. Surulescu

---

Title Page

Contents




Go Back

Close

Quit

Page 20 of 21

- [8] R. SCHALL AND H.G. LUUS, On population and individual bioequivalence, *Statistics in Medicine*, **12** (1993), 1109–1124.
- [9] D.W. SCOTT, *Multivariate Density Estimation. Theory, Practice, and Visualization*, John Wiley & Sons, 1992.
- [10] YI-CHING YAO AND H. IYER, On an inequality for the normal distribution arising in bioequivalence studies, *J. Appl. Prob.*, **36** (1999), 279–286.



---

**A Note on an Individual  
Bioequivalence Setting**

C. Surulescu and N. Surulescu

---

Title Page

Contents



Go Back

Close

Quit

Page 21 of 21