

Journal of Inequalities in Pure and Applied Mathematics

HERMITIAN OPERATORS AND CONVEX FUNCTIONS

JEAN-CHRISTOPHE BOURIN

Université de Cergy-Pontoise
Dépt. de Mathématiques
2 rue Adolphe Chauvin
95302 Pontoise, France.

*E*Mail: bourinjc@club-internet.fr

©2000 Victoria University
ISSN (electronic): 1443-5756
105-05



volume 6, issue 5, article 139,
2005.

Received 06 April, 2005;
accepted 10 November, 2005.

Communicated by: F. Hansen

Abstract

Contents




Home Page

Go Back

Close

Quit



Abstract

We establish several convexity results for Hermitian matrices. For instance: Let A, B be Hermitian and let f be a convex function. If X and Y stand for $f(\{A+B\}/2)$ and $\{f(A) + f(B)\}/2$ respectively, then there exist unitaries U, V such that

$$X \leq \frac{UYU^* + VYV^*}{2}.$$

Consequently, $\lambda_{2j-1}(X) \leq \lambda_j(Y)$, where $\lambda_j(\cdot)$ are the eigenvalues arranged in decreasing order.

2000 Mathematics Subject Classification: 47A30 47A63.

Key words: Hermitian operators, eigenvalues, operator inequalities, Jensen's inequality.

This paper is based on the talk given by the author within the "International Conference of Mathematical Inequalities and their Applications, I", December 06-08, 2004, Victoria University, Melbourne, Australia [<http://rgmia.vu.edu.au/conference>]

Contents

1	Introduction	3
2	Compressions	5
3	Convex Combinations	10
	References	

Hermitian Operators and Convex Functions

Jean-Christophe Bourin

Title Page

Contents



Go Back

Close

Quit

Page 2 of 13

1. Introduction

The main aim of this paper is to give a matrix version of the scalar inequality

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{f(a)+f(b)}{2}$$

for convex functions f on the real line.

Capital letters A, B, \dots, Z mean n -by- n complex matrices, or operators on a finite dimensional Hilbert space \mathcal{H} ; I stands for the identity. When A is positive semidefinite, resp. positive definite, we write $A \geq 0$, resp. $A > 0$.


A classical matrix version of (1.1) is von Neuman's Trace Inequality: For Hermitians A, B ,

$$(1.2) \quad \text{Tr} f\left(\frac{A+B}{2}\right) \leq \text{Tr} \frac{f(A)+f(B)}{2}.$$

When f is convex and monotone, we showed [2] that (1.2) can be extended to an operator inequality: There exists a unitary U such that

$$(1.3) \quad f\left(\frac{A+B}{2}\right) \leq U \cdot \frac{f(A)+f(B)}{2} \cdot U^*.$$

We also established similar inequalities involving more general convex combinations. These inequalities are equivalent to an inequality for compressions. Recall that given an operator Z and a subspace \mathcal{E} with corresponding orthoprojection E , the compression of Z onto \mathcal{E} , denoted by $Z_{\mathcal{E}}$, is the restriction of EZ to \mathcal{E} . Inequality (1.3) can be derived from: For every Hermitian A , subspace \mathcal{E}



Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

and monotone convex function f , there exists a unitary operator U on \mathcal{E} such that

$$(1.4) \quad f(A_{\mathcal{E}}) \leq Uf(A)_{\mathcal{E}}U^*.$$

Inequalities (1.3) and (1.4) are equivalent to inequalities for eigenvalues. For instance (1.4) can be rephrased as

$$\lambda_j(f(A_{\mathcal{E}})) \leq \lambda_j(f(A)_{\mathcal{E}}), \quad j = 1, 2, \dots$$

where $\lambda_j(\cdot)$, $j = 1, 2, \dots$ are the eigenvalues arranged in decreasing order and counted with their multiplicities. Having proved an inequality such as (1.3) for monotone convex functions, it remains to search counterparts for general convex functions. We derived from (1.3) the following result for even convex functions f : Given Hermitians A, B , there exist unitaries U, V such that


$$(1.5) \quad f\left(\frac{A+B}{2}\right) \leq \frac{Uf(A)U^* + Vf(B)V^*}{2}.$$

This generalizes a wellknown inequality for the absolute value,

$$|A+B| \leq U|A|U^* + V|B|V^*.$$

We do not know whether (1.5) is valid for all convex functions.

In Section 2 we present a counterpart of (1.4) for all convex functions. This will enable us to give, in Section 3, a quite natural counterpart of (1.3) for all convex functions. Although (1.3) can be proven independently of (1.4) (and the same for the counterparts), we have a feeling that in the case of general convex functions, the approach via compressions is more illuminating.



Title Page

Contents



Go Back

Close

Quit

Page 4 of 13

2. Compressions

Our substitute to (1.4) for general convex functions (on the real line) is:

Theorem 2.1. *Let A be Hermitian, let \mathcal{E} be a subspace and let f be a convex function. Then, there exist unitaries U, V on \mathcal{E} such that*

$$f(A_{\mathcal{E}}) \leq \frac{Uf(A)_{\mathcal{E}}U^* + Vf(A)_{\mathcal{E}}V^*}{2}.$$

Consequently, for $j = 1, 2, \dots$,


$$\lambda_{2j-1}(f(A_{\mathcal{E}})) \leq \lambda_j(f(A)_{\mathcal{E}}).$$

Proof. We may find spectral subspaces \mathcal{E}' and \mathcal{E}'' for $A_{\mathcal{E}}$ and a real r such that

- (i) $\mathcal{E} = \mathcal{E}' \oplus \mathcal{E}''$,
- (ii) the spectrum of $A_{\mathcal{E}'}$ lies on $(-\infty, r]$ and the spectrum of $A_{\mathcal{E}''}$ lies on $[r, \infty)$,
- (iii) f is monotone both on $(-\infty, r]$ and $[r, \infty)$.

Let k be an integer, $1 \leq k \leq \dim \mathcal{E}'$. There exists a spectral subspace $\mathcal{F} \subset \mathcal{E}'$ for $A_{\mathcal{E}'}$ (hence for $f(A_{\mathcal{E}'})$), $\dim \mathcal{F} = k$, such that

$$\begin{aligned} \lambda_k[f(A_{\mathcal{E}'})] &= \min_{h \in \mathcal{F}; \|h\|=1} \langle h, f(A_{\mathcal{F}})h \rangle \\ &= \min\{f(\lambda_1(A_{\mathcal{F}})); f(\lambda_k(A_{\mathcal{F}}))\} \\ &= \min_{h \in \mathcal{F}; \|h\|=1} f(\langle h, A_{\mathcal{F}}h \rangle) \\ &= \min_{h \in \mathcal{F}; \|h\|=1} f(\langle h, Ah \rangle), \end{aligned}$$



where at the second and third steps we use the monotonicity of f on $(-\infty, r]$ and the fact that $A_{\mathcal{F}}$'s spectrum lies on $(-\infty, r]$. The convexity of f implies

$$f(\langle h, Ah \rangle) \leq \langle h, f(A)h \rangle$$

for all normalized vectors h . Therefore, by the minmax principle,

$$\begin{aligned} \lambda_k[f(A_{\mathcal{E}'})] &\leq \min_{h \in \mathcal{F}; \|h\|=1} \langle h, f(A)h \rangle \\ &\leq \lambda_k[f(A)_{\mathcal{E}'}]. \end{aligned}$$

This statement is equivalent to the existence of a unitary operator U_0 on \mathcal{E}' such that

$$f(A_{\mathcal{E}'}) \leq U_0 f(A)_{\mathcal{E}'} U_0^*.$$

Similarly we get a unitary V_0 on \mathcal{E}'' such that


$$f(A_{\mathcal{E}''}) \leq V_0 f(A)_{\mathcal{E}''} V_0^*.$$

Thus we have

$$f(A_{\mathcal{E}}) \leq \begin{pmatrix} U_0 & 0 \\ 0 & V_0 \end{pmatrix} \begin{pmatrix} f(A)_{\mathcal{E}'} & 0 \\ 0 & f(A)_{\mathcal{E}''} \end{pmatrix} \begin{pmatrix} U_0^* & 0 \\ 0 & V_0^* \end{pmatrix}.$$

Also, we note that, still in respect with the decomposition $\mathcal{E} = \mathcal{E}' \oplus \mathcal{E}''$,

$$\begin{aligned} &\begin{pmatrix} f(A)_{\mathcal{E}'} & 0 \\ 0 & f(A)_{\mathcal{E}''} \end{pmatrix} \\ &= \frac{1}{2} \left\{ \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} f(A)_{\mathcal{E}} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} + \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} f(A)_{\mathcal{E}} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \right\}. \end{aligned}$$



Hermitian Operators and Convex Functions

Jean-Christophe Bourin

Title Page

Contents



Go Back

Close

Quit

Page 6 of 13

So, letting

$$U = \begin{pmatrix} U_0 & 0 \\ 0 & V_0 \end{pmatrix} \quad \text{and} \quad V = \begin{pmatrix} U_0 & 0 \\ 0 & -V_0 \end{pmatrix}$$

we get

$$(2.1) \quad f(A_\varepsilon) \leq \frac{Uf(A)_\varepsilon U^* + Vf(A)_\varepsilon V^*}{2}.$$

It remains to check that (2.1) entails

$$\lambda_{2j-1}(f(A_\varepsilon)) \leq \lambda_j(f(A)_\varepsilon).$$

This follows from the forthcoming elementary observation. □

Proposition 2.2. *Let X, Y be Hermitians such that*

$$(2.2) \quad X \leq \frac{UYU^* + VYV^*}{2}$$

for some unitaries U, V . Then, for $j = 1, 2, \dots$,


$$\lambda_{2j-1}(X) \leq \lambda_j(Y).$$

Proof. By adding a rI term, for a suitable scalar r , both to X and Y , it suffices to show that

$$(2.3) \quad \lambda_{2j-1}(X) > 0 \implies \lambda_j(Y) > 0.$$

We need the following obvious fact: Given Hermitians A, B ,

$$\text{rank}(A + B)_+ \leq \text{rank}A_+ + \text{rank}B_+$$



Hermitian Operators and Convex Functions

Jean-Christophe Bourin

Title Page

Contents



Go Back

Close

Quit

Page 7 of 13

where the subscript + stands for positive parts. Applying this to $A = UYU^*$ and $B = VYV^*$ we infer that the negation of (2.3), that is $\lambda_{2j-1}(A+B) > 0$ and $\lambda_j(A) (= \lambda_j(B)) \leq 0$, cannot hold. Indeed, the relation

$$2j - 1 > (j - 1) + (j - 1)$$

would contradict the previous rank inequality. \square

Remark 1. From inequality (2.2) one also derives, as a straightforward consequence of Fan's Maximum Principle [1, Chapter 4],

$$\sum_{j=1}^k \lambda_j(X) \leq \sum_{j=1}^k \lambda_j(Y)$$

for $k = 1, 2, \dots$


Inequality (2.2) also implies

$$\lambda_{i+j+1}(X) \leq \frac{1}{2} \{ \lambda_{i+1}(Y) + \lambda_{j+1}(Y) \}$$

for $i, j = 0, 1, \dots$. It is a special case of Weyl's inequalities [1, Chapter 3].


Remark 2. For operators acting on an infinite dimensional (separable) space, the main inequality of Theorem 2.1 is still valid at the cost of an additional rI term in the RHS, with $r > 0$ arbitrarily small. See [3, Chapter 1] for the analogous result for (1.4).

Obviously, for a concave function f , the main inequality of Theorem 2.1 is reversed. But the following is open:



Problem 1. Let g be a concave function, let A be Hermitian and let \mathcal{E} be a subspace. Can we find unitaries U, V on \mathcal{E} such that

$$g(A)_\mathcal{E} \leq \frac{Ug(A_\mathcal{E})U^* + Vg(A_\mathcal{E})V^*}{2} ?$$



Hermitian Operators and Convex Functions

Jean-Christophe Bourin

Title Page

Contents



Go Back

Close

Quit

Page 9 of 13

3. Convex Combinations

The next two theorems can be regarded as matrix versions of Jensen's inequality. The first one is also a matrix version of the elementary scalar inequality

$$f(za) \leq zf(a)$$

for convex functions f with $f(0) \leq 0$ and scalars a and z with $0 < z < 1$.

Theorem 3.1. *Let f be a convex function, let A be Hermitian, let Z be a contraction and set $X = f(Z^*AZ)$ and $Y = Z^*f(A)Z$. Then, there exist unitaries U, V such that*

$$X \leq \frac{UYU^* + VYV^*}{2}.$$


A family $\{Z_i\}_{i=1}^m$ is an isometric column if $\sum_{i=1}^m Z_i^*Z_i = I$.

Theorem 3.2. *Let f be a convex function, let $\{A_i\}_{i=1}^m$ be Hermitians, let $\{Z_i\}_{i=1}^m$ be an isometric column and set $X = f(\sum Z_i^*A_iZ_i)$ and $Y = \sum Z_i^*f(A_i)Z_i$. Then, there exist unitaries U, V such that*

$$X \leq \frac{UYU^* + VYV^*}{2}.$$

Corollary 3.3. *Let f be a convex function, let A, B be Hermitians and set $X = f(\{A+B\}/2)$ and $Y = \{f(A) + f(B)\}/2$. Then, there exist unitaries U, V such that*

$$X \leq \frac{UYU^* + VYV^*}{2}.$$



Title Page

Contents



Go Back

Close

Quit

Page 10 of 13

Recall that the above inequality entails that for $j = 1, 2, \dots$,

$$\lambda_{2j-1}(X) \leq \lambda_j(Y).$$

We turn to the proof of Theorems 3.1 and 3.2.

Proof. Theorem 3.1 and Theorem 2.1 are equivalent. Indeed, to prove Theorem 2.1, we may assume that $f(0) = 0$. Then, Theorem 2.1 follows from Theorem 3.1 by taking Z as the projection onto \mathcal{E} .

Theorem 2.1 entails Theorem 3.1: to see that, we introduce the partial isometry J and the operator \tilde{A} on $\mathcal{H} \oplus \mathcal{H}$ defined by

$$J = \begin{pmatrix} Z & 0 \\ (I - |Z|^2)^{1/2} & 0 \end{pmatrix}, \quad \tilde{A} = \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix}.$$

Denoting by \mathcal{H} the first summand of the direct sum $\mathcal{H} \oplus \mathcal{H}$, we observe that


$$f(Z^*AZ) = f(J^*\tilde{A}J):\mathcal{H} = J^*f(\tilde{A}_{J(\mathcal{H})})J:\mathcal{H},$$

where $X : \mathcal{H}$ means the restriction of an operator X to the first summand of $\mathcal{H} \oplus \mathcal{H}$. Applying Theorem 2.1 with $\mathcal{E} = J(\mathcal{H})$, we get unitaries U_0, V_0 on $J(\mathcal{H})$ such that

$$f(Z^*AZ) \leq J^* \frac{U_0 f(\tilde{A})_{J(\mathcal{H})} U_0^* + V_0 f(\tilde{A})_{J(\mathcal{H})} V_0^*}{2} J:\mathcal{H}.$$

Equivalently, there exist unitaries U, V on \mathcal{H} such that

$$\begin{aligned} & f(Z^*AZ) \\ & \leq \frac{U J^* f(\tilde{A})_{J(\mathcal{H})} (J:\mathcal{H}) U^* + V J^* f(\tilde{A})_{J(\mathcal{H})} (J:\mathcal{H}) V^*}{2} \end{aligned}$$



Title Page

Contents



Go Back

Close

Quit

Page 11 of 13

$$\begin{aligned}
&= \frac{1}{2} \left\{ UJ^* \begin{pmatrix} f(A) & 0 \\ 0 & f(0) \end{pmatrix} (J:\mathcal{H})U^* + VJ^* \begin{pmatrix} f(A) & 0 \\ 0 & f(0) \end{pmatrix} (J:\mathcal{H})V^* \right\} \\
&= \frac{1}{2} U \{ Z^* f(A) Z + (I - |Z|^2)^{1/2} f(0) (I - |Z|^2)^{1/2} \} U^* \\
&\quad + \frac{1}{2} V \{ Z^* f(A) Z + (I - |Z|^2)^{1/2} f(0) (I - |Z|^2)^{1/2} \} V^*.
\end{aligned}$$

Using $f(0) \leq 0$ we obtain the first claim of Theorem 3.2.

Similarly, Theorem 2.1 implies Theorem 3.2 (we may assume $f(0) = 0$) by considering the partial isometry and the operator on $\oplus^m \mathcal{H}$,

$$\begin{pmatrix} Z_1 & 0 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ Z_m & 0 & \cdots & 0 \end{pmatrix}, \quad \begin{pmatrix} A_1 & & & \\ & \ddots & & \\ & & & A_m \end{pmatrix}.$$

□


We note that our theorems contain two well-known trace inequalities [4], [5]:

3.4. Brown-Kosaki: Let f be convex with $f(0) \leq 0$ and let A be Hermitian. Then, for all contractions Z ,

$$\text{Tr } f(Z^* A Z) \leq \text{Tr } Z^* f(A) Z.$$


3.5. Hansen-Pedersen: Let f be convex and let $\{A_i\}_{i=1}^m$ be Hermitians. Then, for all isometric column $\{Z_i\}_{i=1}^m$,

$$\text{Tr } f \left(\sum_i Z_i^* A_i Z_i \right) \leq \text{Tr } \sum_i Z_i^* f(A_i) Z_i.$$



References

- [1] R. BHATIA, *Matrix Analysis*, Springer, Germany, 1996.
- [2] J.-C. BOURIN, Convexity or concavity inequalities for Hermitian operators, *Math. Ineq. Appl.*, **7**(4) (2004), 607–620.
- [3] J.-C. BOURIN, Compressions, *Dilations and Matrix Inequalities*, RGMIA monograph, Victoria university, Melbourne 2004 (<http://rgmia.vu.edu.au/monograph>)
- [4] L.G. BROWN AND H. KOSAKI, Jensen's inequality in semi-finite von Neuman algebras, *J. Operator Theory*, **23** (1990), 3–19.
- [5] F. HANSEN AND G.K. PEDERSEN, Jensen's operator inequality, *Bull. London Math. Soc.*, **35** (2003), 553–564.



Hermitian Operators and
Convex Functions

Jean-Christophe Bourin

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13