





[Title Page](#)

[Contents](#)



Page 1 of 10

[Go Back](#)

[Full Screen](#)

[Close](#)

ON QUADRATURE RULES, INEQUALITIES AND ERROR BOUNDS

SZYMON WAŚOWICZ

Department of Mathematics and Computer Science

University of Bielsko-Biała

Willowa 2, 43-309 Bielsko-Biała, Poland

E-Mail: swasowicz@ath.bielsko.pl

Received: 06 June, 2007

Accepted: 22 May, 2008

Communicated by: **P. Cerone**

2000 AMS Sub. Class.: Primary: 41A55; Secondary: 26A51, 26D15.

Key words: Approximate integration, Quadrature rules, Convex functions of higher order.

Abstract: The order structure of the set of six operators connected with quadrature rules is established in the class of 5-convex functions. An error bound of the Lobatto quadrature rule with five knots is given for less regular functions as in the classical result.

Contents

1 Introduction	3
2 Results	5



**Quadrature Rules, Inequalities
and Error Bounds**

Szymon Wąsowicz

vol. 9, iss. 2, art. 36, 2008

[Title Page](#)

[Contents](#)



Page **2** of 10


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 3 of 10

Go Back

Full Screen

Close

1. Introduction

For $f : [-1, 1] \rightarrow \mathbb{R}$ we consider six operators approximating the integral mean value $\frac{1}{2} \int_{-1}^1 f(x) dx$. They are

$$\mathcal{C}(f) := \frac{1}{3} \left(f \left(-\frac{\sqrt{2}}{2} \right) + f(0) + f \left(\frac{\sqrt{2}}{2} \right) \right),$$

$$\mathcal{G}_2(f) := \frac{1}{2} \left(f \left(-\frac{\sqrt{3}}{3} \right) + f \left(\frac{\sqrt{3}}{3} \right) \right),$$

$$\mathcal{G}_3(f) := \frac{4}{9} f(0) + \frac{5}{18} \left(f \left(-\frac{\sqrt{15}}{5} \right) + f \left(\frac{\sqrt{15}}{5} \right) \right),$$

$$\mathcal{L}_4(f) := \frac{1}{12} (f(-1) + f(1)) + \frac{5}{12} \left(f \left(-\frac{\sqrt{5}}{5} \right) + f \left(\frac{\sqrt{5}}{5} \right) \right),$$

$$\mathcal{L}_5(f) := \frac{16}{45} f(0) + \frac{1}{20} (f(-1) + f(1)) + \frac{49}{180} \left(f \left(-\frac{\sqrt{21}}{7} \right) + f \left(\frac{\sqrt{21}}{7} \right) \right),$$

$$\mathcal{S}(f) := \frac{1}{6} (f(-1) + f(1)) + \frac{2}{3} f(0).$$

All of them are connected with the very well known rules of approximate integration: Chebyshev quadrature, Gauss–Legendre quadrature with two and three knots, Lobatto quadrature with four and five knots and Simpson’s Rule, respectively (see e.g. [4, 8, 9, 10, 11]).

In the paper [6] the order structure of the set of above operators was investigated in the class of 3–convex functions. In this note we establish all possible inequalities between these operators in the class of 5–convex functions. As an application we give an error bound of the operator \mathcal{L}_5 for six times differentiable functions instead of eight times differentiable ones as in the classical result.

In this paper only 5–convex functions on $[-1, 1]$ are considered. Recall that the

function $f : [-1, 1] \rightarrow \mathbb{R}$ is called *5-convex* if

$$(1.1) \quad D(x_1, \dots, x_7; f) := \begin{vmatrix} 1 & \dots & 1 \\ x_1 & \dots & x_7 \\ x_1^2 & \dots & x_7^2 \\ x_1^3 & \dots & x_7^3 \\ x_1^4 & \dots & x_7^4 \\ x_1^5 & \dots & x_7^5 \\ f(x_1) & \dots & f(x_7) \end{vmatrix} \geq 0$$

for any x_1, \dots, x_7 such that $-1 \leq x_1 < \dots < x_7 \leq 1$. More detailed introductory notes concerning higher-order convexity were given in [6]. For a wide treatment of this topic we refer the reader to Popoviciu's thesis [3], the very well known books [2] and [5] and to Hopf's thesis [1], where it appeared (without the name) for the first time.



Title Page

Contents




Page 4 of 10

Go Back

Full Screen

Close



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 5 of 10

Go Back

Full Screen

Close

2. Results

Let us start with four technical results.

Lemma 2.1. *If $f : [-1, 1] \rightarrow \mathbb{R}$ is an even 5-convex function then*

$$(w^2 - u^2)(v^2 - u^2)(w^2 - v^2)f(0) + u^2w^2(w^2 - u^2)f(v) \\ \leq w^2v^2(w^2 - v^2)f(u) + v^2u^2(v^2 - u^2)f(w)$$

for any $0 < u < v < w \leq 1$.

Proof. Fix $0 < u < v < w \leq 1$. By 5-convexity, $D(-w, -v, -u, 0, u, v, w; f) \geq 0$. Expand this determinant by the last row and perform elementary computations on Vandermonde determinants. \square

Lemma 2.2. *If $f : [-1, 1] \rightarrow \mathbb{R}$ is 5-convex then so is the function $[-1, 1] \ni x \mapsto f(-x)$.*


Proof. This result is well known from the theory of convex functions of higher order and it holds in fact for convex functions of any odd order (cf. e.g. [3]). However, the proof is easy if we use the condition (1.1) and elementary properties of determinants. \square

By (1.1) it is obvious that a sum of two 5-convex functions is also 5-convex. Then we have the following.

Lemma 2.3. *If $f : [-1, 1] \rightarrow \mathbb{R}$ is 5-convex then so is its even part, i.e. the function*

$$f_e(x) = \frac{f(x) + f(-x)}{2}, \quad x \in [-1, 1].$$

Record also the trivial



Title Page

Contents



Page 6 of 10

Go Back

Full Screen

Close

Lemma 2.4. If $T \in \{\mathcal{C}, \mathcal{G}_2, \mathcal{G}_3, \mathcal{L}_4, \mathcal{L}_5, \mathcal{S}\}$ then $T(f) = T(f_e)$ for any $f : [-1, 1] \rightarrow \mathbb{R}$.

Now we establish all possible inequalities between the considered operators in the class of 5-convex functions.

Theorem 2.5. If $f : [-1, 1] \rightarrow \mathbb{R}$ is 5-convex then $\mathcal{G}_3(f) \leq \mathcal{L}_5(f) \leq \mathcal{L}_4(f)$. In the class of 5-convex functions the operators $\mathcal{G}_2, \mathcal{C}, \mathcal{S}$ are not comparable both with each other and with $\mathcal{G}_3, \mathcal{L}_4, \mathcal{L}_5$.

Proof. By Lemmas 2.3 and 2.4, it is enough to prove the desired inequalities for even 5-convex functions. Using Lemma 2.1 for $u = \frac{\sqrt{21}}{7}, v = \frac{\sqrt{15}}{5}, w = 1$ we obtain $\mathcal{G}_3(f) \leq \mathcal{L}_5(f)$. The inequality $\mathcal{L}_5(f) \leq \mathcal{L}_4(f)$ we get for $u = \frac{\sqrt{5}}{5}, v = \frac{\sqrt{21}}{7}, w = 1$.

Now let $f = \exp, g = 1 - \cos$. Both functions are 5-convex on $[-1, 1]$ since their derivatives of the sixth order are nonnegative on this interval (cf. [2, 3, 5], for a quick reference cf. also [7]). See the table below.

Operator	\mathcal{G}_2	\mathcal{C}	\mathcal{S}	\mathcal{G}_3	\mathcal{L}_5	\mathcal{L}_4
f	1.17135	1.17373	1.18103	1.17517	1.17520	1.17524
g	0.16209	0.15984	0.15323	0.15850	0.15853	0.15857

Then


$$\mathcal{G}_2(f) < \mathcal{C}(f) < \mathcal{G}_3(f) < \mathcal{L}_5(f) < \mathcal{L}_4(f) < \mathcal{S}(f),$$

$$\mathcal{S}(g) < \mathcal{G}_3(g) < \mathcal{L}_5(g) < \mathcal{L}_4(g) < \mathcal{C}(g) < \mathcal{G}_2(g),$$

which proves the second part of the statement. \square

Remark 1. By the example given in the above proof one could expect that the inequality

$$\min\{\mathcal{G}_2, \mathcal{C}, \mathcal{S}\} \leq \mathcal{G}_3 \leq \mathcal{L}_5 \leq \mathcal{L}_4 \leq \max\{\mathcal{G}_2, \mathcal{C}, \mathcal{S}\}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 7 of 10

Go Back

Full Screen

Close

holds in the class of 5-convex functions. However this is not the case since for a 5-convex function $h(x) = x^6 - \frac{3}{2}x^4 + \frac{1}{6}$ we have

$$\mathcal{G}_2(h) = \frac{1}{27}, \quad \mathcal{C}(h) = \mathcal{S}(h) = 0, \quad \mathcal{G}_3(h) = -\frac{1}{75}, \quad \mathcal{L}_5(h) = \frac{1}{105}, \quad \mathcal{L}_4(h) = \frac{1}{25},$$

so

$$\mathcal{G}_3(h) < \mathcal{C}(h) = \mathcal{S}(h) < \mathcal{L}_5(h) < \mathcal{G}_2(h) < \mathcal{L}_4(h).$$

Let us comment on the results of Theorem 2.5. The set $\{\mathcal{C}, \mathcal{G}_2, \mathcal{G}_3, \mathcal{L}_4, \mathcal{L}_5, \mathcal{S}\}$ has 15 two-element subsets. That is why maximally 15 inequalities may be established between the operators considered. For 3-convex functions we have proved in [6] that 12 inequalities hold true and only 3 fail. We can see that for 5-convex functions the situation is quite different: only 3 inequalities are true, the rest are false. Moreover, the operators $\mathcal{G}_2, \mathcal{C}, \mathcal{S}$ comparable for 3-convex functions are not comparable for 5-convex ones, while the operators $\mathcal{G}_3, \mathcal{L}_4, \mathcal{L}_5$ comparable for 5-convex functions are not comparable for 3-convex ones.


The classical error bound of the quadrature \mathcal{L}_5 depends on the derivative of eighth order (cf. [4, 10]). Similarly to the results of the papers [6, 7] we give an error bound of this quadrature for less regular functions: in this paper for six-times differentiable functions. Let $\mathcal{I}(f) := \frac{1}{2} \int_{-1}^1 f(x) dx$. For $f \in C^6([-1, 1])$ denote

$$M(f) := \sup \{|f^{(6)}(x)| : x \in [-1, 1]\}.$$

Corollary 2.6. *If $f \in C^6([-1, 1])$ then $|\mathcal{L}_5(f) - \mathcal{I}(f)| \leq \frac{M(f)}{15750}$.*

Proof. It is well known (cf. [4, 9]) that if $f \in C^6([-1, 1])$, then $\mathcal{I}(f) = \mathcal{G}_3(f) + \frac{f^{(6)}(\xi)}{31500}$ for some $\xi \in (-1, 1)$. Assume for a while that f is 5-convex. Hence by Theorem 2.5

$$\mathcal{I}(f) \leq \mathcal{L}_5(f) + \frac{f^{(6)}(\xi)}{31500}.$$



Title Page

Contents



Page 8 of 10

Go Back

Full Screen

Close

Thus we arrive at

$$(2.1) \quad \mathcal{I}(f) - \mathcal{L}_5(f) \leq \frac{M(f)}{31500}.$$

Now let $f \in \mathcal{C}^6([-1, 1])$ be an arbitrary function and let $g(x) = \frac{M(f)x^6}{720}$. Then $|f^{(6)}| \leq g^{(6)}$ on $[-1, 1]$, whence $(g - f)^{(6)} \geq 0$ and $(g + f)^{(6)} \geq 0$ on $[-1, 1]$. This implies that $g - f$ and $g + f$ are 5-convex on $[-1, 1]$. It is easy to see that $M(g - f) \leq 2M(f)$ and $M(g + f) \leq 2M(f)$. Then we infer by 5-convexity and (2.1),

$$\mathcal{I}(g - f) - \mathcal{L}_5(g - f) \leq \frac{M(g - f)}{31500} \leq \frac{M(f)}{15750} \quad \text{and}$$

$$\mathcal{I}(g + f) - \mathcal{L}_5(g + f) \leq \frac{M(g + f)}{31500} \leq \frac{M(f)}{15750}.$$


It is easy to see that $\mathcal{I}(g) = \mathcal{L}_5(g)$. Since the operators $\mathcal{I}, \mathcal{L}_5$ are linear, then

$$-\mathcal{I}(f) + \mathcal{L}_5(f) \leq \frac{M(f)}{15750} \quad \text{and} \quad \mathcal{I}(f) - \mathcal{L}_5(f) \leq \frac{M(f)}{15750},$$

which concludes the proof. \square

References

- [1] E. HOPF, Über die Zusammenhänge zwischen gewissen höheren Differenzenquotienten reeller Funktionen einer reellen Variablen und deren Differenzierbarkeitseigenschaften, Dissertation, Friedrich–Wilhelms–Universität Berlin, 1926.
- [2] M. KUCZMA, *An Introduction to the Theory of Functional Equations and Inequalities. Cauchy's Equation and Jensen's Inequality*, Państwowe Wydawnictwo Naukowe (Polish Scientific Publishers) and Uniwersytet Śląski, Warszawa–Kraków–Katowice 1985.
- [3] T. POPOVICIU, Sur quelques propriétés des fonctions d'une ou de deux variables réelles, *Mathematica (Cluj)*, **8** (1934), 1–85.
- [4] A. RALSTON, *A First Course in Numerical Analysis*, McGraw–Hill Book Company, New York, St. Louis, San Francisco, Toronto, London, Sydney, 1965.
- [5] A.W. ROBERTS AND D.E. VARBERG, *Convex Functions*, Academic Press, New York 1973.
- [6] S. WAŚOWICZ, Inequalities between the quadrature operators and error bounds of quadrature rules, *J. Ineq. Pure & Appl. Math.*, **8**(2) (2007), Art. 42. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=848>].
- [7] S. WAŚOWICZ, On error bounds for Gauss–Legendre and Lobatto quadrature rules, *J. Ineq. Pure & Appl. Math.*, **7**(3) (2006), Art. 84. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=704>].





Quadrature Rules, Inequalities
and Error Bounds

Szymon Waśowicz

vol. 9, iss. 2, art. 36, 2008

Title Page

Contents



Page 9 of 10


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [8] E.W. WEISSTEIN, Chebyshev Quadrature, From MathWorld – A Wolfram Web Resource. [ONLINE: <http://mathworld.wolfram.com/ChebyshevQuadrature.html>].
- [9] E.W. WEISSTEIN, Legendre–Gauss Quadrature, From MathWorld – A Wolfram Web Resource. [ONLINE: <http://mathworld.wolfram.com/Legendre-GaussQuadrature.html>].
- [10] E.W. WEISSTEIN, Lobatto Quadrature, From MathWorld – A Wolfram Web Resource. [ONLINE: <http://mathworld.wolfram.com/LobattoQuadrature.html>].
- [11] E.W. WEISSTEIN, Simpson’s Rule, From MathWorld – A Wolfram Web Resource. [ONLINE: <http://mathworld.wolfram.com/SimpsonsRule.html>].

**Quadrature Rules, Inequalities
and Error Bounds**

Szymon Wąsowicz

vol. 9, iss. 2, art. 36, 2008

Title Page

Contents



Page 10 of 10

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756