

# Journal of Inequalities in Pure and Applied Mathematics

## A NOTE ON THE ACCURACY OF RAMANUJAN'S APPROXIMATIVE FORMULA FOR THE PERIMETER OF AN ELLIPSE

MARK B. VILLARINO

Escuela de Matemática,  
Universidad de Costa Rica,  
2060 San José, Costa Rica.

*EMail:* [mvallari@cariari.ucr.ac.cr](mailto:mvallari@cariari.ucr.ac.cr)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
192-05



---

volume 7, issue 1, article 21,  
2006.

*Received 27 June, 2005;  
accepted 19 November, 2005.*  
*Communicated by: S.S. Dragomir*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

We present a detailed error analysis, with best possible constants, of Ramanujan's most accurate approximation to the perimeter of an ellipse.


*2000 Mathematics Subject Classification:* 26D15, 33E05, 41A44.

*Key words:* Ramanujan's approximative formula, Best constants, Elliptic functions and integrals.

Support from the Vicerrectoría de Investigación of the University of Costa Rica is acknowledged. We thank the referee for his valuable comments and additional references.

## Contents

1	Introduction .....	3
2	Later History .....	6
3	Fundamental Lemma .....	7
4	Ivory's Identity .....	14
5	The Accuracy Lemma .....	16
6	The Accuracy of Ramanujan's Approximation .....	19
	References	



---

**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 22

# 1. Introduction

Let  $a$  and  $b$  be the semi-major and semi-minor axes of an ellipse with perimeter  $p$  and whose eccentricity is  $k$ . The final sentence of Ramanujan's famous paper *Modular Equations and Approximations to  $\pi$* , [6], says:

“The following approximation for  $p$  [was] obtained empirically:

$$(1.1) \quad p = \pi \left\{ (a + b) + \frac{3(a - b)^2}{10(a + b) + \sqrt{a^2 + 14ab + b^2}} + \varepsilon \right\}$$

where  $\varepsilon$  is about  $\frac{3ak^{20}}{68719476736}$ .”

Ramanujan never explained his “empirical” method of obtaining this approximation, nor ever subsequently returned to this approximation, neither in his published work, nor in his Notebooks [4]. Indeed, although the notebooks do contain the above approximation (see Entry 3 of Chapter XVIII) the statement there does not even mention his asymptotic error estimate stated above.

Twenty years later Watson [7] claimed to have proven that Ramanujan's approximation is *in defect*, but he never published his proof.

In 1978, we established the following optimal version of Ramanujan's approximation:

**Theorem 1.1 (Ramanujan's Approximation Theorem).** *Ramanujan's approximative perimeter*

$$(1.2) \quad p_R := \pi \left\{ (a + b) + \frac{3(a - b)^2}{10(a + b) + \sqrt{a^2 + 14ab + b^2}} \right\}$$



---

A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 22

underestimates the true perimeter,  $p$ , by

$$(1.3) \quad \epsilon := \pi(a + b) \cdot \theta(\lambda) \cdot \lambda^{10},$$

where

$$(1.4) \quad \lambda := \frac{a - b}{a + b},$$

and where the function  $\theta(\lambda)$  grows monotonically in  $0 \leq \lambda \leq 1$  while at the same time it satisfies the optimal inequalities

$$(1.5) \quad \frac{3}{2^{17}} < \theta(\lambda) \leq \frac{14}{11} \left( \frac{22}{7} - \pi \right).$$

Please take note of the striking form of the sharp upper bound since it involves the number  $\left(\frac{22}{7} - \pi\right)$  which measures the accuracy of Archimedes' famous approximation,  $\frac{22}{7}$ , to the transcendental number  $\pi$ !


**Corollary 1.2.** *The error in defect,  $\epsilon$ , as a function of  $\lambda$ , grows monotonically for  $0 \leq \lambda \leq 1$ .*

**Corollary 1.3.** *The error in defect,  $\epsilon$ , as a function of the eccentricity,  $e$ , is given by*

$$(1.6) \quad \epsilon(e) := a \left\{ \delta(e) \left( \frac{2}{1 + \sqrt{1 - e^2}} \right)^{19} \right\} e^{20}.$$

Moreover,  $\epsilon(e)$  grows monotonically with  $e$ ,  $0 \leq e \leq 1$ , while  $\delta(e)$  satisfies the optimal inequalities

$$(1.7) \quad \frac{3\pi}{68719476736} < \delta(e) \leq \frac{7}{2^{18}} \left( \frac{22}{7} - \pi \right).$$



A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse

Mark B. Villarino

Title Page

Contents




Go Back

Close

Quit

Page 4 of 22

This Corollary 1.3 explains the significance of Ramanujan's own error estimate in (1.1). The latter is an asymptotic *lower bound* for  $\epsilon(e)$  but it is not the optimal one. That is given in (1.7).



---

**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit


Page 5 of 22

## 2. Later History

We sent an (updated) copy of our 1978 preprint to Bruce Berndt in 1988 and he subsequently quoted its conclusions in his edition of Volume 3 of the Notebooks (see p. 150 [4]). However the details of our proofs have never been published and so we have decided to present them in this paper.

Berndt's discussion of Ramanujan's approximation includes Almkvist's very plausible suggestion that Ramanujan's "empirical process" was to develop a *continued fraction expansion* of Ivory's infinite series for the perimeter ([1]) as well as a proof, due independently to Almkvist and Askey, of our fundamental lemma (see §3). However, their proof is different from ours.

The most recent work on the subject includes that by R. Barnard, K. Pearce, and K. Richards in [3], published in the year 2000, and the paper by H. Alzer and Qui, S.-L. (see [2]), which was published in the year 2004. The former also prove the major conclusion in our fundamental lemma, but their methods too are quite different from ours. The latter includes a sharp lower bound for elliptical arc length in terms of a power-mean type function. But their methods are also quite different from ours.



---

**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 22

### 3. Fundamental Lemma

**Lemma 3.1 (Fundamental Lemma).** Define the functions  $A(x)$  and  $B(x)$  and the coefficients  $A_n$  and  $B_n$  by:

$$(3.1) \quad A(x) := 1 + \frac{3x}{10 + \sqrt{4 - 3x}} := 1 + A_1x + A_2x^2 + \dots,$$

$$(3.2) \quad B(x) := \sum_{n=0}^{\infty} \left\{ \frac{1}{2n-1} \frac{1}{4^n} \binom{2n}{n} \right\}^2 x^n := 1 + B_1x + B_2x^2 + \dots.$$

Then:


$$(3.3) \quad A_1 = B_1, \quad A_2 = B_2, \quad A_3 = B_3, \quad A_4 = B_4$$

$$(3.4) \quad A_5 < B_5, \quad A_6 < B_6, \dots, \quad A_n < B_n, \dots,$$

where the strict inequalities in (3.4) are valid for all  $n \geq 5$ .

*Proof.* First we prove (3.3). We read this off directly from the numerical values of the expansion:

$$\begin{aligned} A_1 = B_1 &= \frac{1}{4} \\ A_2 = B_2 &= \frac{1}{16} \\ A_3 = B_3 &= \frac{1}{64} \\ A_4 = B_4 &= \frac{25}{4096}. \end{aligned}$$



A Note On the Accuracy of Ramanujan's Approximative Formula for the Perimeter of an Ellipse

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 7 of 22

Now we prove (3.4). For  $A_5$ ,  $B_5$ ,  $A_6$ , and  $B_6$  we verify (3.4) directly from their explicit numerical values. Namely,

$$A_5 = \frac{47\frac{1}{2}}{2^{14}}, B_5 = \frac{49}{2^{14}} \Rightarrow A_5 - B_5 = \frac{-\frac{3}{2}}{2^{14}} < 0$$

$$A_6 = \frac{803}{2^{21}}, B_6 = \frac{882}{2^{21}} \Rightarrow A_6 - B_6 = \frac{-79}{2^{21}} < 0.$$

Therefore it is sufficient to prove

$$(3.5) \quad A_n < B_n$$

for all


$$(3.6) \quad n \geq 7.$$

Now the *explicit* formula for  $A_n$  is

$$(3.7) \quad A_n = a_{n-1} + a_{n-2} + a_{n-3} + \cdots + a_1 + a_0$$

where

$$(3.8) \quad \begin{aligned} a_{n-1} &:= \frac{1}{2n-3} \cdot \frac{1}{16^n} \binom{2n-2}{n-1} 3^{n-1} \\ a_{n-2} &:= \frac{1}{2n-5} \cdot \frac{1}{16^{n-1}} \binom{2n-4}{n-2} 3^{n-2} \left(\frac{-1}{2^5}\right) \\ &\vdots \\ a_1 &:= \frac{1}{2 \cdot 1 - 1} \frac{1}{16^2} \binom{2}{1} 3^1 \left(\frac{-1}{2^5}\right)^{n-2} \\ a_0 &:= \frac{4}{16} \left(\frac{-1}{2^5}\right)^{n-1}. \end{aligned}$$



**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 8 of 22



Next we write

$$(3.9) \quad A_n = a_{n-1} \left( 1 + \frac{a_{n-2}}{a_{n-1}} + \frac{a_{n-3}}{a_{n-1}} + \frac{a_{n-4}}{a_{n-1}} + \cdots + \frac{a_1}{a_{n-1}} + \frac{a_0}{a_{n-1}} \right)$$

and assert:

**Claim 1.** *The ratios  $\frac{a_{n-k-1}}{a_{n-k}}$  increase monotonically in absolute value as  $k$  increases from  $k = 1$  to  $k = n - 1$ .*

*Proof.* For  $k = 1, \dots, n - 2$ ,

$$\begin{aligned} \left| \frac{a_{n-k-1}}{a_{n-k}} \right| &= \left( 1 + \frac{2}{2n - 2k - 3} \right) \left( \frac{1}{2} + \frac{1}{4n - 4k - 2} \right) \frac{1}{12} \\ &\leq \frac{1}{6} \quad (\text{which is the worst case and occurs when } k = n - 2) \\ &< 1 \end{aligned}$$


For  $k = n - 1$ ,

$$\left| \frac{a_0}{a_1} \right| = \frac{1}{3} < 1.$$

This completes the proof. □

**Claim 2.** *The ratios  $\frac{a_{n-k-1}}{a_{n-k}}$  alternate in sign.*

*Proof.* This is a consequence of the definition of the  $a_k$ . □




---

**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 9 of 22

By Claim 1 and Claim 2 we can write (3.9) in the form

$$A_n = a_{n-1} (1 - \text{something positive and smaller than } 1) < a_{n-1}.$$

Therefore, to prove (3.8) for  $n \geq 7$ , it suffices to prove that

$$(3.10) \quad a_{n-1} < B_n$$

for all  $n \geq 7$ .

By (3.8) and the definition of  $B_n$ , this last affirmation is equivalent to proving

$$\frac{1}{2n-3} \cdot \frac{1}{16^n} \binom{2n-2}{n-1} 3^{n-1} < \left\{ \frac{1}{2n-1} \cdot \frac{1}{4^n} \binom{2n}{n} \right\}^2,$$

which, after some algebra, reduces to proving the implication

$$n \geq 7 \Rightarrow \frac{\frac{n}{2} \cdot \frac{2n-1}{2n-3}}{\binom{2n}{n}} \cdot 3^{n-1} < 1.$$


If we define for all integers  $n \geq 7$

$$(3.11) \quad f(n) := \frac{\frac{n}{2} \cdot \frac{2n-1}{2n-3}}{\binom{2n}{n}} \cdot 3^{n-1}$$

then the affirmation (3.10) turns out to be equivalent to

$$(3.12) \quad n \geq 7 \Rightarrow f(n) < 1$$

This latter affirmation is a consequence of the following two conditions:



**A Note On the Accuracy of Ramanujan's Approximative Formula for the Perimeter of an Ellipse**

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 10 of 22

**Condition 1.**  $f(7) < 1$ .

**Condition 2.**  $f(7) > f(8) > f(9) > \dots > f(k) > f(k+1) > \dots$

□

*Proof of Condition 1.* By direct numerical computation,

$$f(7) = \frac{1701}{1936} < 1.$$

□

*Proof of Condition 2.* We must show that

$$k \geq 7 \Rightarrow f(k) > f(k+1).$$

If we define


$$(3.13) \quad g(k) := \frac{f(k)}{f(k+1)},$$

then we must show that

$$(3.14) \quad k \geq 7 \Rightarrow g(k) > 1.$$

Using the definition (3.11) of  $f(n)$  and the definition (3.14) of  $g(n)$ , and reducing algebraically we find

$$g(k) = \frac{2k}{6k-9} \left( \frac{2k-1}{k+1} \right)^2,$$



---

A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 11 of 22

and we must show that

$$(3.15) \quad k \geq 7 \Rightarrow \frac{2k}{6k-9} \left( \frac{2k-1}{k+1} \right)^2 > 1.$$

Define the rational function of the real variable  $x$ :

$$(3.16) \quad g(x) := \frac{2x}{6x-9} \left( \frac{2x-1}{x+1} \right)^2.$$

Then the graph of  $y = g(x)$  has a vertical asymptote at  $x = \frac{3}{2}$  and

$$(3.17) \quad \lim_{x \rightarrow \frac{3}{2}^+} g(x) = +\infty.$$

Moreover, the derivative of  $g(x)$  is given by:


$$g'(x) = \frac{2(2x^2 - 7x + 1)}{x(x+1)(2x-1)(2x+3)},$$

which implies that

$$g'(x) \begin{cases} < 0 & \text{if } \frac{3}{2} < x < \frac{7+\sqrt{41}}{4}, \\ = 0 & \text{if } x = \frac{7+\sqrt{41}}{4}, \\ > 0 & \text{if } x > \frac{7+\sqrt{41}}{4}. \end{cases}$$

Therefore, for  $x \geq \frac{3}{2}$ ,  $g(x)$  decreases from “ $+\infty$ ” at  $x = \frac{3}{2}$  (see (3.17)) to an *absolute minimum value* (in  $\frac{3}{2} \leq x < \infty$ )

$$g \left( \frac{7 + \sqrt{41}}{4} \right) = 1 + \frac{37 - \sqrt{41}}{399 + 69\sqrt{41}} = 1.0363895208 \dots$$



**A Note On the Accuracy of Ramanujan's Approximative Formula for the Perimeter of an Ellipse**

Mark B. Villarino

Title Page

Contents




Go Back

Close

Quit

Page 12 of 22

and then *increases monotonically* as  $x \rightarrow \infty$  to its asymptotic limit  $y = \frac{4}{3}$  and this is enough to complete the proof of the Fundamental Lemma.  $\square$



---

**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 13 of 22

## 4. Ivory's Identity


In 1796, J. Ivory [5] published the following identity (in somewhat different notation):

**Theorem 4.1 (Ivory's Identity).** *If  $0 \leq x \leq 1$  then the following formula for  $B(x)$  is valid:*

$$(4.1) \quad \frac{1}{\pi} \int_0^\pi \sqrt{1 + 2\sqrt{x} \cos(2\phi) + x} d\phi = \sum_{n=0}^{\infty} \left\{ \frac{1}{2n-1} \frac{1}{4^n} \binom{2n}{n} \right\}^2 x^n \equiv B(x).$$

*Proof.* We sketch his elegant proof.

$$\begin{aligned} & \frac{1}{\pi} \int_0^\pi \sqrt{1 + 2\sqrt{x} \cos(2\phi) + x} d\phi \\ &= \frac{1}{\pi} \int_0^\pi \sqrt{1 + \sqrt{x}e^{2i\phi}} \sqrt{1 + \sqrt{x}e^{-2i\phi}} d\phi \\ &= \frac{1}{\pi} \int_0^\pi \sum_{m=0}^{\infty} \left\{ \frac{1}{2m-1} \cdot \frac{1}{4^m} \binom{2m}{m} (\sqrt{x})^m e^{2\pi im\phi} \right\} \\ & \quad \times \sum_{n=0}^{\infty} \left\{ \frac{1}{2n-1} \cdot \frac{1}{4^n} \binom{2n}{n} (\sqrt{x})^n e^{-2\pi in\phi} \right\} d\phi \\ &= \frac{1}{\pi} \sum_{m=0}^{\infty} \left\{ \frac{1}{2m-1} \cdot \frac{1}{4^m} \binom{2m}{m} (\sqrt{x})^m \right\} \end{aligned}$$



A Note On the Accuracy of Ramanujan's Approximative Formula for the Perimeter of an Ellipse

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 14 of 22

$$\begin{aligned} & \times \sum_{n=0}^{\infty} \left\{ \frac{1}{2n-1} \cdot \frac{1}{4^n} \binom{2n}{n} (\sqrt{x})^n \right\} \int_0^{\pi} e^{2\pi i(m-n)\phi} d\phi \\ & = \sum_{n=0}^{\infty} \left\{ \frac{1}{2n-1} \cdot \frac{1}{4^n} \binom{2n}{n} \right\}^2 x^n \end{aligned}$$

□

We will need the following evaluation in our investigation of the accuracy of Ramanujan's approximation.


**Corollary 4.2.**

$$(4.2) \quad \mathbf{B}(1) = \frac{4}{\pi}.$$

*Proof.* By Ivory's identity,

$$\begin{aligned} \mathbf{B}(1) &= \frac{1}{\pi} \int_0^{\pi} \sqrt{1 + 2\sqrt{1} \cos(2\phi) + 1} d\phi \\ &= \frac{1}{\pi} \int_0^{\pi} \sqrt{2 + 2 \cos(2\phi)} d\phi \\ &= \frac{1}{\pi} \int_0^{\pi} \sqrt{4 \cos^2(\phi)} d\phi \\ &= \frac{4}{\pi}. \end{aligned}$$

□



**A Note On the Accuracy of Ramanujan's Approximative Formula for the Perimeter of an Ellipse**

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 15 of 22

## 5. The Accuracy Lemma

**Theorem 5.1 (Accuracy Lemma).** For  $0 \leq x \leq 1$ , the function

$$(5.1) \quad \mathbf{A}(x) := 1 + \frac{3x}{10 + \sqrt{4 - 3x}}$$

underestimates the function

$$(5.2) \quad \mathbf{B}(x) := \sum_{n=0}^{\infty} \left\{ \frac{1}{2n-1} \frac{1}{4^n} \binom{2n}{n} \right\}^2 x^n$$

by a discrepancy,  $\Delta(x)$  which is never more than  $\left(\frac{4}{\pi} - \frac{14}{11}\right) x^5$  and which is always more than  $\frac{3}{2^{17}} x^5$ :

$$(5.3) \quad \frac{3}{2^{17}} x^5 < \Delta(x) \leq \left(\frac{4}{\pi} - \frac{14}{11}\right) x^5.$$


Moreover, the constants  $\left(\frac{4}{\pi} - \frac{14}{11}\right)$  and  $\frac{3}{2^{17}} x^5$  are the best possible.

*Proof.* By the definition of  $\mathbf{A}(x)$  and  $\mathbf{B}(x)$  given in Theorem 1.1, the discrepancy  $\Delta(x)$  is given by the series

$$\begin{aligned} \Delta(x) &:= \mathbf{B}(x) - \mathbf{A}(x) \\ &= (B_5 - A_5)x^5 + (B_6 - A_6)x^6 + \dots \\ &:= \delta_5 x^5 + \delta_6 x^6 + \dots, \end{aligned}$$

where, again by Theorem 1.1,

$$\delta_k > 0 \quad \text{for } k = 5, 6, \dots$$



---

**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 16 of 22



On the one hand

$$\begin{aligned}
 \Delta(x) &= x^5(\delta_5 + \delta_6x + \dots) \\
 &\leq x^5(\delta_5 + \delta_6 + \delta_7 + \dots) \\
 &= x^5\Delta(1) \\
 &= x^5\{\mathbf{B}(1) - \mathbf{A}(1)\} \\
 &= x^5\left(\frac{4}{\pi} - \frac{14}{11}\right)
 \end{aligned}$$

where we used Corollary 1.2 of Ivory's identity in the last equality. Therefore

$$\Delta(x) \leq \left(\frac{4}{\pi} - \frac{14}{11}\right)x^5.$$

This is half of the accuracy lemma. Moreover, the constant  $\left(\frac{4}{\pi} - \frac{14}{11}\right)$  is assumed for  $x = 1$  and thus cannot be replaced by anything smaller, i.e., it is the best possible constant.

On the other hand, we can write


$$\Delta(x) = x^5\{\delta_5 + G(x)\},$$

where

$$G(x) := \delta_6x + \delta_7x^2 + \dots \Rightarrow \begin{cases} G(x) \geq 0 & \text{for all } 0 \leq x \leq 1, \\ G(x) \rightarrow 0 & \text{as } x \rightarrow 0. \end{cases}$$

This shows that

$$\Delta(x) > \delta_5x^5 = \frac{3}{2^{17}}x^5$$



**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 17 of 22

and that

$$\lim_{x \rightarrow 0} \frac{\Delta(x)}{x^5} = \frac{3}{2^{17}}.$$

This proves both the other inequality in the theorem and the optimality of the constant  $\delta_5 = \frac{3}{2^{17}}$ , i.e., that it cannot be replaced by any larger constant.

This completes the proof of the Accuracy Lemma.  $\square$



---

**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 22

## 6. The Accuracy of Ramanujan's Approximation

Now we can achieve the main goal of this paper, namely to prove *Ramanujan's Approximation Theorem*.

First we express the perimeter of an ellipse and Ramanujan's approximative perimeter in terms of the functions  $\mathbf{A}(x)$  and  $\mathbf{B}(x)$ .

**Theorem 6.1.** *If  $p$  is the perimeter of an ellipse with semimajor axes  $a$  and  $b$ , and if  $p_R$  is Ramanujan's approximative perimeter, then:*

$$(6.1) \quad p = \pi(a+b) \cdot \mathbf{B} \left\{ \left( \frac{a-b}{a+b} \right)^2 \right\}$$

$$p_R = \pi(a+b) \cdot \mathbf{A} \left\{ \left( \frac{a-b}{a+b} \right)^2 \right\}.$$

*Proof.* We begin with *Ivory's Identity* (§4) and in it we substitute  $x := \left( \frac{a-b}{a+b} \right)^2$ . Then the integral becomes

$$\begin{aligned} \frac{1}{\pi} \int_0^\pi \sqrt{1 + 2\sqrt{\left( \frac{a-b}{a+b} \right)^2} \cos(2\phi) + \left( \frac{a-b}{a+b} \right)^2} d\phi \\ = \frac{4}{\pi(a+b)} \int_0^{\frac{\pi}{2}} (a^2 \sin^2 \phi + b^2 \cos^2 \phi) d\phi \end{aligned}$$



A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 19 of 22

and therefore

$$\mathbf{B} \left\{ \left( \frac{a-b}{a+b} \right)^2 \right\} = \frac{4}{\pi(a+b)} \int_0^{\frac{\pi}{2}} (a^2 \sin^2 \phi + b^2 \cos^2 \phi) d\phi.$$

But, it is well known (Berndt [4]) that the perimeter,  $p$ , of an ellipse with semi-axes  $a$  and  $b$  is given by

$$p = 4 \int_0^{\frac{\pi}{2}} (a^2 \sin^2 \phi + b^2 \cos^2 \phi) d\phi,$$

and thus

$$(6.2) \quad p = \pi(a+b) \cdot \mathbf{B} \left\{ \left( \frac{a-b}{a+b} \right)^2 \right\}.$$


Moreover, some algebra shows us that

$$\begin{aligned} \mathbf{A} \left\{ \left( \frac{a-b}{a+b} \right)^2 \right\} &= 1 + \frac{3 \left( \frac{a-b}{a+b} \right)^2}{10 + \sqrt{4 - 3 \left( \frac{a-b}{a+b} \right)^2}} \\ &= \frac{1}{a+b} \left\{ (a+b) + \frac{3(a-b)^2}{10(a+b) + \sqrt{a^2 + 14ab + b^2}} \right\} \end{aligned}$$

and we conclude that Ramanujan's approximative formula,  $p_R$  is given by

$$(6.3) \quad p_R = \pi(a+b) \mathbf{A} \left\{ \left( \frac{a-b}{a+b} \right)^2 \right\}.$$

□



**A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse**

Mark B. Villarino

Title Page

Contents



Go Back

Close

Quit

Page 20 of 22

The formula for  $p$  above was the object of Ivory's original paper [5].  
 Now we complete the proof of Theorem 1.1.

*Proof.* Writing

$$\lambda := \frac{a - b}{a + b},$$


and using the notation of the statement of Theorem 1.1. we conclude that

$$\begin{aligned} \epsilon &:= \pi(a + b) \cdot \theta(\lambda) \cdot \lambda^{10} \\ &= \pi(a + b) \cdot \frac{\Delta(\lambda^2)}{\lambda^{10}} \cdot \lambda^{10}, \end{aligned}$$

where

$$(6.4) \quad \theta(\lambda) \equiv \frac{\Delta(\lambda^2)}{\lambda^{10}} = \delta_5 + \delta_6 \lambda^2 + \dots$$

Now we apply the Accuracy Lemma and the proof is complete. □




---

**A Note On the Accuracy of  
 Ramanujan's Approximative  
 Formula for the Perimeter of an  
 Ellipse**

Mark B. Villarino

---

Title Page

Contents



Go Back


Close

Quit

Page 21 of 22

## References

- [1] G. ALMKVIST AND B. BERNDT, Gauss, Landen, Ramanujan, the Arithmetic-geometric Mean, Ellipses,  $\pi$ , and the Ladies Diary, *Amer. Math. Monthly*, **95** (1988), 585–608.
- [2] H. ALZER AND S.-L. QIU, Monotonicity theorems and inequalities for the complete elliptic integrals, *J. Comp. Appl. Math.*, **172** (2004), 289–312.
- [3] R.W. BARNARD, K. PEARCE AND K.C. RICHARDS, A monotonicity property involving  ${}_3F_2$  and comparisons of the classical approximations of elliptical arc length, *SIAM J. Math. Anal.*, **32** (2000), 403–419.
- [4] B. BERNDT, *Ramanujan's Notebooks, Volume 3*, Springer, New York, 1998.
- [5] J. IVORY, A new series for the rectification of the ellipsis; together with some observations on the evolution of the formula  $(a^2 + b^2 - 2ab \cos \phi)^n$ , *Trans. R. Soc. Edinburgh*, **4** (1796), 177–190.
- [6] S. RAMANUJAN, *Ramanujan's Collected Works*, Chelsea, New York, 1962.
- [7] G.N. WATSON, The Marquis and the land agent, *Mathematical Gazette*, **17** (1933), 5–17.



---

A Note On the Accuracy of  
Ramanujan's Approximative  
Formula for the Perimeter of an  
Ellipse

Mark B. Villarino

---

Title Page

Contents



Go Back

Close

Quit

Page 22 of 22