

Journal of Inequalities in Pure and Applied Mathematics

INEQUALITIES FOR CONVEX SETS

PAUL R. SCOTT AND POH WAH AWYONG

Department of Pure Mathematics
University of Adelaide
S.A. 5005 AUSTRALIA.
EMail: pscott@maths.adelaide.edu.au

Division of Mathematics
National Institute of Education
469 Bukit Timah Road, SINGAPORE 259756
EMail: awyongpw@nievax.nie.ac.sg

©2000 [School of Communications and Informatics](#), Victoria University of Technology
ISSN (electronic): 1443-5756
016-99



volume 1, issue 1, article 6,
2000.

*Received 29 November, 1999;
accepted 4 February, 2000.*

Communicated by: C.E.M. Pearce

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

This paper collects together known inequalities relating the area, perimeter, width, diameter, inradius and circumradius of planar convex sets. Also, a technique for finding new inequalities is stated and illustrated.


2000 Mathematics Subject Classification: 52A10, 52A40

Key words: planar convex set, inequality, area, perimeter, diameter, width, inradius, circumradius

Contents

1	Introduction	3
2	Inequalities Involving Two Functionals	4
3	Inequalities Involving Three Functionals	6
4	Obtaining New Inequalities	10

References



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back


Close

Quit

Page 2 of 13

1. Introduction

Let K be a convex set in the plane. Associated with K are a number of well-known functionals: the area $A = A(K)$, the perimeter $p = p(K)$, the diameter $d = d(K)$, the width $w = w(K)$, the inradius $r = r(K)$ and the circumradius $R = R(K)$. For many years we have been interested in inequalities involving these functionals. As the literature is extensive and spans more than 80 years, it will be helpful to summarize the known inequalities. This is done in sections 2 and 3. Then in section 4 we explore a technique for suggesting new inequalities, and give a number of conjectured new inequalities, mostly obtained by this method.



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

2. Inequalities Involving Two Functionals

Table 2.1 lists the known inequalities involving two functionals. The **extremal sets** referred to in the table are described below the table. These demonstrate that the inequalities are best possible. All the proofs of the results in this table can be found in the indicated sections of Yaglom and Boltyanskiĭ's book [17]. The (d, R) and (w, r) results are respectively known as Jung's Theorem and Blaschke's Theorem. Where a dagger (\dagger) appears in the reference column, the result is trivial.



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents




Go Back

Close

Quit

Page 4 of 13



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit

Page 5 of 13

Parameters	Inequality	Extremal Set	Reference
A, d	$4A \leq \pi d^2$	\bigcirc	p. 239, ex. 610a
A, p	$4\pi A \leq p^2$	\bigcirc	p. 207, ex. 5.8
A, r	$\pi r^2 \leq A$	\bigcirc	†
A, R	$A \leq \pi R^2$	\bigcirc	†
A, w	$w^2 \leq \sqrt{3}A$	\triangle_E	p. 221, ex. 6.4
d, p	$2d < p \leq \pi d$	$, W$	†; p. 257, ex. 7.17a
d, r	$2r \leq d$	\bigcirc	†
d, R	$\sqrt{3}R \leq d \leq 2R$	\triangle_E, \bigcirc	p. 213, ex. 6.1; †
d, w	$w \leq d$	W	†
p, r	$2\pi r \leq p$	\bigcirc	†
p, R	$4R < p \leq 2\pi R$	$, W$	†
p, w	$\pi w \leq p$	W	p. 258, ex. 7.18a
r, R	$r \leq R$	\bigcirc	†
r, w	$2r \leq w \leq 3r$	\bigcirc, \triangle_E	†; p. 215, ex. 6.2
R, w	$w \leq 2R$	\bigcirc	†

Table 2.1: Inequalities involving two functionals.


The extremal sets

- $|$ line segment
- \bigcirc circle
- \triangle_E equilateral triangle
- W sets of constant width

3. Inequalities Involving Three Functionals

Table 3.1 lists the known inequalities involving three functionals. Extra information, signified in the **Notes** column, appears after the table. The **extremal sets** referred to in the table are also listed after the table. These demonstrate that the inequalities are best possible. Some of the results in the table are established in [17]. These are indicated by a number in the References column, and details appear after the table. No inequalities appear to be known relating (p, r, R) or (p, R, w) .

Param.	Condition	Inequality	Note	Ext. Set	Ref.
1. A, d, p	$2d \leq p \leq 3d$ $3d \leq p \leq \pi d$	$8\phi A \leq p(p - 2d \cos \phi)$ $d(p - 2d) \leq 4A \leq pd$	1a	\emptyset $ \cdot, \circ$	[8] ¹ [5]
		$4A \geq (p - 2d)\sqrt{4pd - p^2}$ $4A \geq \sqrt{3}d(p - 2d)$	1b	\triangle_I \triangle_E	[8, 9] ² [9]
2. A, d, r		$A < 2dr$		\parallel	[6]
3. A, d, R		$(2R - d)A$ $\leq \pi(3\sqrt{3} - 5)R^3$	3a		[15]
4. A, d, w	$2w \leq \sqrt{3}d$ $\sqrt{3} < 2w < 2d$	$A < wd$		\parallel	[8]
		$2A \leq w\sqrt{d^2 - w^2}$ $+ 2d^2 \arcsin(w/d)$		\oplus	[8] ³
		$2A \geq wd$		\triangle	[8] ⁴
		$2A > \pi w^2 - \sqrt{3}d^2$ $+ 6w^2(\tan \delta - \delta)$	4a	\triangle_Y	[16]
		$2A \geq 3dw - \sqrt{3}d^2$ $2A \geq \pi w^2 - \sqrt{3}d^2$		\triangle_E \triangle_R	[16] [16]
5. A, p, r		$A \leq r(p - \pi r)$		\circ	[2]
		$2A \geq pr$		\triangle	[3]
6. A, p, R		$A \leq R(p - \pi R)$		\circ	[2]
		$A < 2R(p - 2R)$	6a		[6]
		$A > R(p - 4R)$		$ $	[4]



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents




Go Back

Close

Quit

Page 6 of 13



7. A, p, w	$2\sqrt{3}w \leq p$ $\pi w < p < 2\sqrt{3}w$ $\pi w = p$	$2A \leq w \left(p - \frac{1}{2}\pi w \right)$ $A \geq A_1^*$ $2A \geq w \left(p - \sqrt{3}w \sec^2 \gamma \right)$ $2A \geq \left(\pi - \sqrt{3} \right) w^2$ $6A \geq 4\sqrt{3}w^2 - pw$ $4A \geq pw - \frac{2}{\sqrt{3}}w^2$ $6A \geq pw$	7a 7b	$\circ \equiv \circ$ \triangle_I \triangle_Y \triangle_R \triangle_E \triangle_E \triangle_E	$[8]^5$ $[18]^6$ $[18]$ $[10, 11]^7$ $[16]$ $[7]$ $[16]$
8. A, r, R		$A < 4rR$ $A > 2rR$		\parallel $ $	$[6]$ $[6]$
9. A, r, w		$4(w-2r)A < w^3$ $\sqrt{3}(w-2r)A \leq w^2r$ $(w-2r)A \leq \sqrt{3}wr^2$ $\leq 3\sqrt{3}r^3$		\sqcup \triangle_E \triangle_E	$[14]$ $[14]$ $[14]$
10. A, R, w		$A < 4Rw$ $A > \sqrt{3}Rw$		\parallel \triangle_E	$[6]$ $[6]$
11. d, p, r		$p < 2d + 4r$		\parallel	$[6]$
12. d, p, R		$(2R-d)p$ $\leq (2\sqrt{3}-3)\pi R^2$		W	$[15]$
13. d, p, w		$p \leq 2\sqrt{d^2 - w^2}$ $+ 2d \arcsin(w/d)$ $p \geq 2\sqrt{d^2 - w^2}$ $+ 2w \arcsin(w/d)$		\oplus \diamond	$[8]^8$ $[8]^9$
14. d, r, R		$(2R-d)r \leq (3\sqrt{3}-5)R^2$		\triangle_R	$[15]$
15. d, r, w		$\sqrt{3}(w-2r)d \leq 2wr \leq 6r^2$ $2(w-2r)d \leq w^2$		\triangle_E \sqcup	$[13, 14]$ $[14]$
16. d, R, w		$(2R-d)w$ $\leq \sqrt{3}(2-\sqrt{3})R^2$ $3(2R-d)$ $\leq 2(2-\sqrt{3})w$		\triangle_R \triangle_E	$[15]$ $[1]$
17. p, r, R					

Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit


Page 7 of 13

18. p, r, W	$\sqrt{3}(w-2r)p \leq 2w^2$	\triangle_E	[14]
	$(w-2r)p \leq 2\sqrt{3}wr$ $\leq 6\sqrt{3}r^2$	\triangle_E	[14]
19. p, R, w			
20. r, R, w	$4(w-2r)R \leq w^2$	\sqcup	[14]
	$3(w-2r)R \leq 2wr$ $\leq 6r^2$	\triangle_E	[14]

Table 3.1: Inequalities involving three functionals

The extremal sets

	line segment	○	circle
()	lens: the intersection of two congruent circular disks	◊	convex hull of a disk and two symmetrically placed points
○≡○	convex hull of two congruent circles	⊖	intersection of a disk and a symmetrically placed strip
△	triangle	△ _E	equilateral triangle
△ _I	isoceses triangle	△ _R	Reuleaux triangle
△ _Y	equilateral Yamanouti triarc, bounded by three circular arcs of radius w whose centres lie at the vertices of an equilateral triangle of side length d , and by the six tangents drawn from the vertices of this triangle to these arcs	⊔	half strip, occuring as the limit of an isoceses triangle with increasing altitude on a given base
W	sets of constant width		infinite strip bounded by two parallel lines



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit

Page 8 of 13

Notes

1. A, p, d Note 1a: $2\phi d = p \sin \phi$.
Note 1b: This is not best possible unless $p = 3d$.
3. A, d, R Note 3a: This bound is not best possible. See Conjecture 4.7 below.
4. A, d, w Note 4a: Here $\delta = \arccos(w/d)$.
6. A, p, r Note 6a: This bound is not best possible.
7. A, d, w Note 7a: Here A_1^* is the middle root of the equation


$$128px^3 - 16w(5p^2 + w^2)x^2 + 16w^2p^3x - w^3p^4 = 0.$$

Note 7b: $6w(\tan \gamma - \gamma) = p - \pi w$.

References

The precise references to the proofs in Yaglom and Boltyanskiĭ [17] are:

- | | | |
|----------------------------------|----------------------------------|----------------------------------|
| ¹ p. 240, ex. 6.11(a) | ² p. 229, ex. 6.8(a) | ³ p. 240, ex. 6.10(b) |
| ⁴ p. 227, ex. 6.7 | ⁵ p. 241, ex. 6.11(b) | ⁶ p. 231, ex. 6.8(b) |
| ⁷ p. 260, ex. 7.20 | ⁸ p. 257, ex. 7.17(b) | ⁹ p. 258, ex. 7.18(b) |



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit

Page 9 of 13

4. Obtaining New Inequalities

Osserman [12] takes the classical isoperimetric inequality

$$(4.1) \quad p^2 \geq 4\pi A$$


and from it derives the new inequality

$$(4.2) \quad p^2 - 4\pi A \geq \pi^2(R - r)^2.$$

We can think of this in the following way. The function $f(K) = \pi^2(R - r)^2$ takes the value 0 for the extremal set (the circle) of (4.1). Hence for the circle, (4.2) and (4.1) are identical. Osserman shows that in fact (4.2) is satisfied for all other sets K .

Let us see if we can adapt this method to find other new inequalities. The method will become clear by an example. From Table 2.1 we have Jung's Theorem: $d \geq \sqrt{3}R$, with equality when and only when the set K is an equilateral triangle. For an equilateral triangle, we know that $2A = dw$ or $d = \frac{2A}{w}$. Substituting into Jung's inequality suggests the inequality $2A \geq \sqrt{3}Rw$, which is in fact true and was discovered by Henk (see [6]). So the technique is as follows:

- Consider a known inequality and its extremal set.
- Take an equality relating functionals of the extremal set which includes a functional in the chosen inequality.
- Substitute for this functional in the chosen inequality to obtain a new inequality.



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit

Page 10 of 13

Of course the new inequality may be incorrect; it is then quickly discarded. Again, the new inequality may be trivial, in the sense that it is a combination of known simpler inequalities. For example, the technique gives

$$(w - 2r)p \leq 2A, \text{ with extremal set } \triangle_E;$$

this occurs as a combination of $(w-2r)p \leq 2w^2/\sqrt{3}$ (Table 3.1) and $w^2 \leq \sqrt{3}A$ (Table 2.1). Similarly,

$$4\pi r^2 \leq pd, \text{ with extremal set } \bigcirc,$$

occurs as a combination of $2\pi r \leq p$ and $2r \leq d$ (Table 2.1). However, the following more interesting conjectures have been obtained by this method.

Conjecture 4.1. $2w^2 \leq \sqrt{3}pr$ with extremal set \triangle_E .

Conjecture 4.2. $\sqrt{3}wR \leq pr$, \triangle_E .

Conjecture 4.3. $(p - 2d)w \leq 2A$, \triangle_E .


Conjecture 4.4. $wp \leq 9dr$, \triangle .

Conjecture 4.5. $3(w - 2r)(p - 2d) \leq 2A$, \triangle_E .

Conjecture 4.6. $dw \leq pr$, \triangle .

Finally we have the older [15]

Conjecture 4.7. $2(2R - d)A \leq 3(2 - \sqrt{3})(\pi - \sqrt{3})R^3$, \triangle_R .



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit

Page 11 of 13

References

- [1] P.W. AWYONG, An inequality relating the circumradius and diameter of two-dimensional lattice-point-free convex bodies, *Amer. Math. Monthly*, to appear.
- [2] T. BONNESSEN, *Les problèmes des isopérimètres et des isépiphanes*, Gauthier-Villars, Paris, 1929.
- [3] T. BONNESSEN AND W. FENCHEL, *Théorie der konvexen Körper*, Springer, Berlin, 1929.
- [4] J. FAVARD, Problèmes d'extremums relatifs aux courbes convexes? *Ann. École Norm. Sup.*, **46** (1929), 345–369.
- [5] T. HAYASHI, The extremal chords of an oval, *Tôhoku Math. J.* **22** (1923), 387–393.
- [6] M. HENK AND G.A. TSINTSIFAS, Some inequalities for planar convex figures, *Elem. Math.*, **49** (1994), 120–124.
- [7] S. KAWAI, An inequality for the closed convex curve, *Tôhoku Math. J.*, **36** (1932), 50–57.
- [8] T. KUBOTA, Einige Ungleichheitsbeziehungen über Eilinen und Eiflächen, *Sci. Rep. of the Tôhoku Univ. Ser. (1)*, **12** (1923), 45–65.
- [9] T. KUBOTA, Eine Ungleichheit für Eilinen, *Math. Z.*, **20** (1924), 264–266.



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents




Go Back

Close

Quit

Page 12 of 13

- [10] H. LEBESGUE, Sur le problème des isopérimètres et sur les domaines de largeur constante, *Bull. Soc. Math. France, C. R.*, **42** (1914), 72–76.
- [11] H. LEBESGUE, Sur quelques questions de minimum, relatives aux courbes orbiformes et sur leur rapports avec le calcul des variations, *J. Math. Pures App.*, **4**(8) (1921), 67–96.
- [12] R. OSSERMAN, The isoperimetric inequality, *Bull. Amer. Math. Soc.*, **84**(6) (1978), 1182–1238.
- [13] P.R. SCOTT, Two inequalities for convex sets in the plane, *Bull. Austral. Math. Soc.*, **19** (1978), 131–133.
- [14] P.R. SCOTT, A family of inequalities for convex sets, *Bull. Austral. Math. Soc.*, **20** (1979), 237–245.
- [15] P.R. SCOTT, Sets of constant width and inequalities, *Quart. J. Math. Oxford Ser. (2)*, **32** (1981), 345–348.
- [16] M. SHOLANDER, On certain minimum problems in the theory of convex curves, *Trans. Amer. Math. Soc.*, **73** (1952), 139–173.
- [17] I.M. YAGLOM AND V.G. BOLTYANSKIÏ, *Convex figures*, (Translated by P.J. Kelly and L.F. Walton), Holt, Rinehart and Winston, New York, 1961.
- [18] M. YAMANOUTI, Notes on closed convex figures, *Proceedings of the Physico-Mathematical Society of Japan Ser. (3)*, **14** (1932), 605–609.



Inequalities for Convex Sets

Paul R. Scott and
Poh Wah Awyong

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13