

Journal of Inequalities in Pure and Applied Mathematics

REVERSE WEIGHTED L_p -NORM INEQUALITIES IN CONVOLUTIONS


SABUROU SAITOH, VŨ KIM TUẤN AND MASAHIRO YAMAMOTO

Department of Mathematics
Faculty of Engineering
Gunma University
Kiryu 376-8515, JAPAN.
EMail: ssaitoh@eg.gunma-u.ac.jp

Department of Mathematics and Computer Science
Faculty of Science
Kuwait University
P.O. Box 5969, Safat 13060, KUWAIT.
EMail: vu@sci.kuniv.edu.kw

Department of Mathematical Sciences
The University of Tokyo
3-8-1 Komaba
Tokyo 153-8914, JAPAN.
EMail: myama@ms.u-tokyo.ac.jp

©2000 [School of Communications and Informatics](#), Victoria University of Technology
ISSN (electronic): 1443-5756
018-99



volume 1, issue 1, article 7,
2000.

*Received and accepted
14 December, 1999.*

Communicated by: J.E. Pečarić

Abstract

Contents




Home Page

Go Back

Close

Quit



Abstract

Various weighted L_p ($p > 1$)-norm inequalities in convolutions were derived by using Hölder's inequality. Therefore, by using reverse Hölder inequalities one can obtain reverse weighted L_p -norm inequalities. These inequalities are important in studying stability of some inverse problems.

2000 Mathematics Subject Classification: 44A35, 26D20

Key words: Convolution, weighted L_p inequality, reverse Hölder inequality, inverse problems, Green's function, integral transform, stability.

The authors wish to express their sincere thanks to Professor Josip Pečarić for his valuable information on the reverse Hölder inequality. The work of the second named author was supported by Kuwait University Research Administration under project SM 187.

Contents

1	Introduction	3
2	A general reverse weighted L_p convolution inequality	6
3	Examples	9
	3.1 The first order differential equation	9
	3.2 Picard transform	10
	3.3 Poisson integrals	12
	3.4 Heat equation	14

References

Reverse Weighted L_p -Norm Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 2 of 16

1. Introduction

For the Fourier convolution

$$(f * g)(x) = \int_{-\infty}^{\infty} f(x - \xi)g(\xi) d\xi,$$

the Young's inequality

$$(1.1) \quad \|f * g\|_r \leq \|f\|_p \|g\|_q, \quad f \in L_p(\mathbb{R}), g \in L_q(\mathbb{R}), r^{-1} = p^{-1} + q^{-1} - 1 \quad (p, q, r > 0),$$

is fundamental. Note, however, that for the typical case of $f, g \in L_2(\mathbb{R}^n)$, the inequality (1.1) does not hold. In a series of papers [4, 5, 6, 7] (see also [1]) the first author obtained the following weighted L_p ($p > 1$) inequality for convolution.

Proposition 1.1. ([7]). *For two nonvanishing functions $\rho_j \in L_1(\mathbb{R})$ ($j = 1, 2$) the following L_p ($p > 1$) weighted convolution inequality*

$$(1.2) \quad \left\| \left((F_1 \rho_1) * (F_2 \rho_2) \right) (\rho_1 * \rho_2)^{\frac{1}{p}-1} \right\|_p \leq \|F_1\|_{L_p(\mathbb{R}, |\rho_1|)} \|F_2\|_{L_p(\mathbb{R}, |\rho_2|)}$$


holds for $F_j \in L_p(\mathbb{R}, |\rho_j|)$ ($j = 1, 2$). Equality holds if and only if

$$(1.3) \quad F_j(x) = C_j e^{\alpha x},$$

where α is a constant such that $e^{\alpha x} \in L_p(\mathbb{R}, |\rho_j|)$ ($j = 1, 2$).

Here

$$\|F\|_{L_p(\mathbb{R}; \rho)} = \left\{ \int_{-\infty}^{\infty} |F(x)|^p \rho(x) dx \right\}^{\frac{1}{p}}.$$



Reverse Weighted L_p -Norm
Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and
Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

Unlike the Young's inequality, the inequality (1.2) holds also in case $p = 2$.

In many cases of interest, the convolution is given in the form

$$(1.4) \quad \rho_2(x) \equiv 1, \quad F_2(x) = G(x),$$

where $G(x - \xi)$ is some Green's function. Then the inequality (1.2) takes the form

$$(1.5) \quad \|(F\rho) * G\|_p \leq \|\rho\|_p^{1-\frac{1}{p}} \|G\|_p \|F\|_{L_p(\mathbb{R}, |\rho|)},$$

where ρ , F , and G are such that the right hand side of (1.5) is finite.


The inequality (1.5) enables us to estimate the output function

$$(1.6) \quad \int_{-\infty}^{\infty} F(\xi)\rho(\xi)G(x - \xi) d\xi$$

in terms of the input function F . In this paper we are interested in the reverse type inequality for (1.5), namely, we wish to estimate the input function F by means of the output (1.6). This kind of estimate is important in inverse problems. Our estimate is based on the following version of the reverse Hölder inequality

Proposition 1.2. ([2], see also [3], pages 125-126). *For two positive functions f and g satisfying*

$$(1.7) \quad 0 < m \leq \frac{f}{g} \leq M < \infty$$



Reverse Weighted L_p -Norm Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit


Page 4 of 16

on the set X , and for $p, q > 0$, $p^{-1} + q^{-1} = 1$,

$$(1.8) \quad \left(\int_X f d\mu \right)^{\frac{1}{p}} \left(\int_X g d\mu \right)^{\frac{1}{q}} \leq A_{p,q} \left(\frac{m}{M} \right) \int_X f^{\frac{1}{p}} g^{\frac{1}{q}} d\mu,$$

if the right hand side integral converges. Here

$$A_{p,q}(t) = p^{-\frac{1}{p}} q^{-\frac{1}{q}} t^{-\frac{1}{pq}} (1-t) \left(1 - t^{\frac{1}{p}}\right)^{-\frac{1}{p}} \left(1 - t^{\frac{1}{q}}\right)^{-\frac{1}{q}}.$$



**Reverse Weighted L_p -Norm
Inequalities in Convolutions**

Saburoou Saitoh, Vũ Kim Tuấn and
Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 5 of 16

2. A general reverse weighted L_p convolution inequality

Our main result is the following

Theorem 2.1. *Let F_1 and F_2 be positive functions satisfying*

$$(2.1) \quad 0 < m_1^{\frac{1}{p}} \leq F_1(x) \leq M_1^{\frac{1}{p}} < \infty, \quad 0 < m_2^{\frac{1}{p}} \leq F_2(x) \leq M_2^{\frac{1}{p}} < \infty, \quad p > 1, \quad x \in \mathbb{R}.$$

Then for any positive functions ρ_1 and ρ_2 we have the reverse L_p -weighted convolution inequality

$$(2.2) \quad \left\| ((F_1\rho_1) * (F_2\rho_2)) (\rho_1 * \rho_2)^{\frac{1}{p}-1} \right\|_p \geq \left\{ A_{p,q} \left(\frac{m_1 m_2}{M_1 M_2} \right) \right\}^{-1} \|F_1\|_{L_p(\mathbb{R}, \rho_1)} \|F_2\|_{L_p(\mathbb{R}, \rho_2)}.$$


Inequality (2.2) and others should be understood in the sense that if the left hand side is finite, then so is the right hand side, and in this case the inequality holds.

Proof. Let

$$f(\xi) = F_1^p(\xi) F_2^p(x - \xi) \rho_1(\xi) \rho_2(x - \xi), \quad g(\xi) = \rho_1(\xi) \rho_2(x - \xi).$$

Then condition (2.1) implies

$$m_1 m_2 \leq \frac{f(\xi)}{g(\xi)} \leq M_1 M_2, \quad \xi \in \mathbb{R}.$$



Reverse Weighted L_p -Norm
Inequalities in Convolutions

Sabrou Saitoh, Vũ Kim Tuấn and
Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

Hence, one can apply the reverse Hölder inequality (1.8) for f and g to get

$$A_{p,q} \left(\frac{m_1 m_2}{M_1 M_2} \right) \int_{-\infty}^{\infty} F_1(\xi) \rho_1(\xi) F_2(x - \xi) \rho_2(x - \xi) d\xi \\ \geq \left\{ \int_{-\infty}^{\infty} F_1^p(\xi) F_2^p(x - \xi) \rho_1(\xi) \rho_2(x - \xi) d\xi \right\}^{\frac{1}{p}} \left\{ \int_{-\infty}^{\infty} \rho_1(\xi) \rho_2(x - \xi) d\xi \right\}^{1 - \frac{1}{p}}.$$


Hence,

$$(2.3) \quad \left\{ \int_{-\infty}^{\infty} F_1(\xi) \rho_1(\xi) F_2(x - \xi) \rho_2(x - \xi) d\xi \right\}^p \left\{ \int_{-\infty}^{\infty} \rho_1(\xi) \rho_2(x - \xi) d\xi \right\}^{1-p} \\ \geq \left\{ A_{p,q} \left(\frac{m_1 m_2}{M_1 M_2} \right) \right\}^{-p} \int_{-\infty}^{\infty} F_1^p(\xi) F_2^p(x - \xi) \rho_1(\xi) \rho_2(x - \xi) d\xi.$$

Taking integration of both sides of (2.3) with respect to x from $-\infty$ to ∞ we obtain the inequality

$$(2.4) \quad \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} F_1(\xi) \rho_1(\xi) F_2(x - \xi) \rho_2(x - \xi) d\xi \right\}^p \left\{ \int_{-\infty}^{\infty} \rho_1(\xi) \rho_2(x - \xi) d\xi \right\}^{1-p} dx \\ \geq \left\{ A_{p,q} \left(\frac{m_1 m_2}{M_1 M_2} \right) \right\}^{-p} \int_{-\infty}^{\infty} F_1^p(\xi) \rho_1(\xi) d\xi \int_{-\infty}^{\infty} F_2^p(x) \rho_2(x) dx.$$

Raising both sides of the inequality (2.4) to power $\frac{1}{p}$ yields the inequality (2.2). \square



Reverse Weighted L_p -Norm Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

Inequality (1.8) reverses the sign if $0 < p < 1$. Hence, inequality (2.2) reverses the sign if $0 < p < 1$.


In formula (2.3) replacing ρ_2 by 1, and $F_2(x - \xi)$ by $G(x - \xi)$, and taking integration with respect to x from c to d we arrive at the following inequality

$$(2.5) \quad \int_c^d \left(\int_{-\infty}^{\infty} F(\xi) \rho(\xi) G(x - \xi) d\xi \right)^p dx \\ \geq \left\{ A_{p,q} \left(\frac{m}{M} \right) \right\}^{-p} \left(\int_{-\infty}^{\infty} \rho(\xi) d\xi \right)^{p-1} \int_{-\infty}^{\infty} F^p(\xi) \rho(\xi) d\xi \int_{c-\xi}^{d-\xi} G^p(x) dx,$$

valid if positive continuous functions ρ , F , and G satisfy

$$(2.6) \quad 0 < m^{\frac{1}{p}} \leq F(\xi)G(x - \xi) \leq M^{\frac{1}{p}}, \quad x \in [c, d], \quad \xi \in \mathbb{R}.$$

Inequality (2.5) is especially important when $G(x - \xi)$ is a Green's function. See examples in the next section.



Reverse Weighted L_p -Norm Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16

3. Examples

3.1. The first order differential equation

The solution $y(x)$ of the first order differential equation

$$y'(x) + \lambda y(x) = F(x), \quad y(0) = 0,$$

is represented in the form

$$y(x) = \int_0^x F(t) e^{-\lambda(x-t)} dt.$$

So we shall consider the integral transform

$$f(x) = \int_0^x F(t) \rho(t) e^{-\lambda(x-t)} dt, \quad \lambda > 0.$$

Take


$$G(x) = \begin{cases} e^{-\lambda x}, & x > 0 \\ 0, & x < 0 \end{cases}.$$

The condition (2.6) reads

$$(3.1) \quad 0 < m^{\frac{1}{p}} \leq F(t) e^{-\lambda(x-t)} \leq M^{\frac{1}{p}}.$$

It will be satisfied for $0 \leq t \leq x \leq d < \infty$, if we have

$$(3.2) \quad 0 < m^{\frac{1}{p}} e^{\lambda d - \lambda t} \leq F(t) \leq M^{\frac{1}{p}}, \quad 0 < d < \frac{1}{p\lambda} \log \frac{M}{m}.$$



Reverse Weighted L_p -Norm
Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and
Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 9 of 16

Notice that

$$\int_{c-\xi}^{d-\xi} G^p(x) dx = \begin{cases} \frac{e^{-\lambda pc} - e^{-\lambda pd}}{\lambda p} e^{\lambda p \xi}, & \xi < c, \\ \frac{1 - e^{\lambda p \xi - \lambda pd}}{\lambda p}, & c < \xi < d. \end{cases}$$

Thus the inequality (2.5) yields

$$(3.3) \quad \int_c^d f^p(x) \left(\int_0^x \rho(t) dt \right)^{1-p} dx \\ \geq \left\{ A_{p,q} \left(\frac{m}{M} \right) \right\}^{-p} \frac{1}{\lambda p} \left[(e^{-\lambda pc} - e^{-\lambda pd}) \int_0^c F^p(\xi) \rho(\xi) e^{\lambda p \xi} d\xi \right. \\ \left. + \int_c^d F^p(\xi) \rho(\xi) (1 - e^{-\lambda pd} e^{\lambda p \xi}) d\xi \right].$$

Here we assume that ρ is a positive continuous function on $[0, d]$, and F satisfies (3.2).


3.2. Picard transform

Note that $\frac{1}{2}e^{-|x-t|}$ is the Green's function for the boundary value problem

$$y'' - y = 0, \quad \lim_{x \rightarrow \pm\infty} y(x) = 0.$$

So, we shall consider the Picard transform

$$f(x) = \frac{1}{2} \int_{-\infty}^{\infty} F(t) \rho(t) e^{-|x-t|} dt.$$



Reverse Weighted L_p -Norm
Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and
Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

Take $G(x) = e^{-|x|}$. Since

$$e^{-a}e^{|t|} \leq e^{|x-t|} \leq e^ae^{|t|}, \quad |x| \leq a,$$

we see that the condition (2.6)

$$(3.4) \quad 0 < m^{\frac{1}{p}} \leq F(t)e^{-|x-t|} \leq M^{\frac{1}{p}},$$

holds if

$$(3.5) \quad 0 < m^{\frac{1}{p}}e^ae^{|t|} \leq F(t) \leq M^{\frac{1}{p}}e^{-a}e^{|t|}, \quad t \in \mathbb{R}, \quad 0 < a < \frac{1}{2p} \log \frac{M}{m}.$$


We have

$$\int_{c-t}^{d-t} G^p(x) dx = \int_{c-t}^{d-t} e^{-p|x|} dx = \begin{cases} \frac{e^{pt}}{p} [e^{-pc} - e^{-pd}], & t < c, \\ \frac{e^{-pt}}{p} [e^{pd} - e^{pc}], & t > d, \\ \frac{1}{p} (2 - e^{pc-pt} - e^{pt-pd}), & c < t < d. \end{cases}$$

Thus, for $-a \leq c, d \leq a$ the inequality (2.5) yields

$$(3.6) \quad \int_c^d f^p(x) dx \geq \frac{1}{2^p p} \left\{ A_{p,q} \left(\frac{m}{M} \right) \right\}^{-p} \left(\int_{-\infty}^{\infty} \rho(t) dt \right)^{p-1} \\ \left[(e^{-pc} - e^{-pd}) \int_{-\infty}^c F^p(t) \rho(t) e^{pt} dt + (e^{pd} - e^{pc}) \int_d^{\infty} F^p(t) \rho(t) e^{-pt} dt \right. \\ \left. + \int_c^d F^p(t) \rho(t) (2 - e^{pc-pt} - e^{pt-pd}) dt \right],$$

if ρ is positive continuous, and F satisfies (3.5).



Reverse Weighted L_p -Norm Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 11 of 16

3.3. Poisson integrals

Consider the Poisson integral

$$(3.7) \quad u(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} F(\xi) \rho(\xi) \frac{y}{(x - \xi)^2 + y^2} d\xi.$$

Take

$$G(x) = \frac{y}{x^2 + y^2}.$$

Let

$$\xi \in [a, b], \quad x \in [c, d].$$

Denote

$$\alpha = \max\{|a - c|, |a - d|, |b - c|, |b - d|\}.$$

We have


$$\frac{y}{\alpha^2 + y^2} \leq \frac{y}{(x - \xi)^2 + y^2} \leq \frac{1}{y}.$$

Thus,

$$\int_{c-\xi}^{d-\xi} G^p(x) dx = \int_{c-\xi}^{d-\xi} \left(\frac{y}{x^2 + y^2} \right)^p dx \geq (d - c) \left(\frac{y}{\alpha^2 + y^2} \right)^p.$$

Hence, for a function F satisfying

$$\frac{\alpha^2 + y^2}{y} m^{\frac{1}{p}} \leq F(\xi) \leq y M^{\frac{1}{p}},$$



**Reverse Weighted L_p -Norm
Inequalities in Convolutions**

Sabouro Saitoh, Vũ Kim Tuấn and
Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 12 of 16

and for a positive continuous function ρ on $[a, b]$ we obtain

$$(3.8) \quad \int_c^d u^p(x, y) dx \geq \frac{(d-c)}{\pi^p} \left(\frac{y}{\alpha^2 + y^2} \right)^p \left\{ A_{p,q} \left(\frac{m}{M} \right) \right\}^{-p} \\ \left(\int_a^b \rho(\xi) d\xi \right)^{p-1} \int_a^b F^p(\xi) \rho(\xi) d\xi.$$

Consider now the conjugate Poisson integral

$$(3.9) \quad v(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} F(\xi) \rho(\xi) \frac{x - \xi}{(x - \xi)^2 + y^2} d\xi.$$

Take

$$G(x) = \frac{x}{x^2 + y^2}.$$

For

$$\xi \in [a, b], \quad x \in [c, d], \quad (b < c),$$

we have


$$\frac{c-b}{(d-a)^2 + y^2} \leq \frac{x-\xi}{(x-\xi)^2 + y^2} \leq \frac{d-a}{(c-b)^2 + y^2}.$$

Thus,

$$\int_{c-\xi}^{d-\xi} G^p(x) dx = \int_{c-\xi}^{d-\xi} \left(\frac{x}{x^2 + y^2} \right)^p dx \geq (d-c) \left(\frac{c-b}{(d-a)^2 + y^2} \right)^p.$$

Hence, for a function F satisfying

$$\frac{(d-a)^2 + y^2}{c-b} m^{\frac{1}{p}} \leq F(\xi) \leq \frac{(c-b)^2 + y^2}{d-a} M^{\frac{1}{p}},$$



Reverse Weighted L_p -Norm Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 13 of 16

and for a positive continuous function ρ on $[a, b]$ we obtain

$$(3.10) \quad \int_c^d v^p(x, y) dx \geq \frac{(d-c)}{\pi^p} \left(\frac{c-b}{(d-a)^2 + y^2} \right)^p \left\{ A_{p,q} \left(\frac{m}{M} \right) \right\}^{-p} \\ \left(\int_a^b \rho(\xi) d\xi \right)^{p-1} \int_a^b F^p(\xi) \rho(\xi) d\xi.$$

3.4. Heat equation

We consider the Weierstrass transform

$$(3.11) \quad u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} F(\xi) \rho(\xi) \exp\left(-\frac{(x-\xi)^2}{4t}\right) d\xi,$$

which gives the formal solution $u(x, t)$ of the heat equation

$$u_t = \Delta u \quad \text{on} \quad \mathbb{R}_+ \times \mathbb{R},$$

subject to the initial condition


$$u(x, 0) = F(x) \rho(x) \quad \text{on} \quad \mathbb{R}.$$

Take

$$G(x) = e^{-\frac{x^2}{4t}}.$$

Let

$$x \in [-a, a], \quad \xi \in [-b, b], \quad a + b \leq \sqrt{\frac{4t}{p} \log \frac{M}{m}}.$$



**Reverse Weighted L_p -Norm
Inequalities in Convolutions**

Sabouro Saitoh, Vũ Kim Tuấn and
Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 14 of 16

From

$$1 \leq \exp\left(\frac{(x - \xi)^2}{4t}\right) \leq \exp\left(\frac{(a + b)^2}{4t}\right),$$

we have

$$0 < m^{\frac{1}{p}} \leq F(\xi) \exp\left(-\frac{(x - \xi)^2}{4t}\right) \leq M^{\frac{1}{p}},$$

if

$$(3.12) \quad m^{\frac{1}{p}} \exp\left(\frac{(a + b)^2}{4t}\right) \leq F(\xi) \leq M^{\frac{1}{p}}, \quad \xi \in [-b, b].$$

It is easy to see that

$$\int_{c-\xi}^{d-\xi} e^{-\frac{px^2}{4t}} dx = \sqrt{\frac{\pi t}{p}} \left[\operatorname{erf}\left(\frac{\sqrt{p}(d - \xi)}{2\sqrt{t}}\right) - \operatorname{erf}\left(\frac{\sqrt{p}(c - \xi)}{2\sqrt{t}}\right) \right],$$


where

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

is the error function. Therefore, for $-a \leq c < d \leq a$, the inequality (2.5) yields

$$(3.13) \quad \int_c^d u(x, t)^p dx \geq \frac{1}{2^p (\pi t)^{(p-1)/2} \sqrt{p}} \left\{ A_{p,q} \left(\frac{m}{M} \right) \right\}^{-p} \left(\int_{-b}^b \rho(\xi) d\xi \right)^{p-1} \\ \int_{-b}^b F^p(\xi) \rho(\xi) \left[\operatorname{erf}\left(\frac{\sqrt{p}(d - \xi)}{2\sqrt{t}}\right) - \operatorname{erf}\left(\frac{\sqrt{p}(c - \xi)}{2\sqrt{t}}\right) \right] d\xi,$$

where ρ is a positive continuous function on $[-b, b]$, and F satisfies (3.12).



Reverse Weighted L_p -Norm Inequalities in Convolutions

Saburou Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back


Close

Quit

Page 15 of 16

References

- [1] M. CWICKEL AND R. KERMAN, On a convolution inequality of Saitoh, *Proc. Amer. Math. Soc.*, **124** (1996), 773-777.
- [2] L. XIAO-HUA, On the inverse of Hölder inequality, *Math. Practice and Theory*, **1** (1990), 84-88.
- [3] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [4] S. SAITOH, A fundamental inequality in the convolution of L_2 functions on the half line, *Proc. Amer. Math. Soc.*, **91** (1984), 285-286.
- [5] S. SAITOH, Inequalities in the most simple Sobolev space and convolution of L_2 functions with weights, *Proc. Amer. Math. Soc.*, **118** (1993), 515-520.
- [6] S. SAITOH, Various operators in Hilbert spaces introduced by transforms, *Intern. J. Appl. Math.*, **1** (1999), 111-126.
- [7] S. SAITOH, Weighted L_p -norm inequalities in convolutions, *Handbook on Classical Inequalities*, Kluwer Academic Publishers, Dordrecht (to appear).
- [8] H.M. SRIVASTAVA AND R.G. BUSCHMAN, *Theory and Applications of Convolution Integral Equations*, Kluwer Academic Publishers, Dordrecht, 1992.



Reverse Weighted L_p -Norm Inequalities in Convolutions

Sabouro Saitoh, Vũ Kim Tuấn and Masahiro Yamamoto

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16