


# Journal of Inequalities in Pure and Applied Mathematics

## INEQUALITIES ON POLYNOMIAL HEIGHTS

LAURENȚIU PANAITOPOL AND DORU ȘTEFĂNESCU

University of Bucharest  
70109 Bucharest 1,  
ROMANIA.  
EMail: [pan@al.math.unibuc.ro](mailto:pan@al.math.unibuc.ro)

University of Bucharest,  
P.O.Box 39–D5, Bucharest 39,  
ROMANIA  
EMail: [stef@irma.u-strasbg.fr](mailto:stef@irma.u-strasbg.fr)



---

volume 2, issue 1, article 7,  
2001.

*Received 21 June, 2000;  
accepted 26 October 2000.*

*Communicated by: H. Gauchman*

---

Abstract

Contents




Home Page

Go Back

Close

Quit



## Abstract

We give explicit bounds for the absolute values of the coefficients of the divisors of a complex polynomial. They are expressed in function of the coefficients and of upper and lower bounds for the roots. These bounds are compared with other estimates, in particular with the inequality of Beauzamy [B. Beauzamy, Products of polynomials and a priori estimates for coefficients in polynomial decompositions: A sharp result, *J. Symbolic Comput.*, **13** (1992), 463–472]. Through examples it is proved that for some cases our evaluations give better upper limits.

*2000 Mathematics Subject Classification:* 12D05, 12D10, 12E05, 26C05.

*Key words:* Inequalities, Polynomials.

We are grateful to the referee for the pertinent suggestions concerning Propositions 2.1 and 2.3.

## Contents

1	Introduction .....	3
2	A Height Estimation .....	4
3	A Smallest Divisor .....	11
	References	

---

### Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 13

# 1. Introduction

If  $P(X) = \sum_{i=0}^d a_i X^i \in \mathbb{C}[X]$ , the height of the polynomial  $P$  is defined by  $H(P) = \max(|a_0|, |a_1|, \dots, |a_d|)$ . Other polynomial sizes are the norm  $\|P\| = \sqrt{\sum_{j=0}^d |a_j|^2}$ , the measure  $M(P) = \exp \left\{ \int_0^1 \frac{1}{2\pi} \log |P(e^{2i\pi\theta})| d\theta \right\}$  and Bombieri's norm  $[P]_2 = \sqrt{\sum_{j=0}^d |a_j|^2 / \binom{d}{j}}$ .


There exist many estimates for the height of an arbitrary polynomial divisor  $Q$  of  $P$ . They can be expressed in function of polynomial sizes as the norm, the measure or Bombieri's norm. We mention, for example, the estimate

$$H(Q) \leq \binom{d}{\lfloor d/2 \rfloor} M(P),$$

where the measure can be easily computed by the method of M. Mignotte [5].

For integer polynomials one of the best height estimates uses the norm of Bombieri [2] and was obtained by Beauzamy [1]

$$H(Q) \leq l_n [P]_2, \quad \text{with} \quad l_n = \frac{3^{3/4} \cdot 3^{n/2}}{2(\pi n)^{1/2}}.$$



Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

## 2. A Height Estimation

We present another estimate for the heights of proper divisors of a complex polynomial. It makes use of an inequality of M. Mignotte [6]. A key step is the consideration of complex polynomials with roots of moduli greater than 2.

**Proposition 2.1.** *Let  $P(X) = X^n + a_{n-1}X^{n-1} + \dots + a_1X + a_0 \in \mathbb{C}[X] \setminus \mathbb{C}$ , such that  $P(0) \neq 0$  and let  $\mu$  be a lower bound of the absolute values of the roots of  $P$ . If  $Q$  is a monic proper divisor of  $P$  in  $[X]$ , then*

$$H(Q) < H(P) \quad \text{if } \mu \geq 2,$$

$$H(Q) < \max_{0 \leq i \leq n} \left| a_i \left( \frac{2}{\mu} \right)^{n-i} \right| \quad \text{if } \mu < 2.$$

*Proof.* We suppose

$$Q(X) = X^k + b_{k-1}X^{k-1} + \dots + b_1X + b_0.$$


Let  $\xi_1, \dots, \xi_n \in \mathbb{C}$  be the roots of  $P$ . Without loss of generality we may suppose that  $\xi_1, \dots, \xi_k$  are the roots of  $Q$ . Note that

$$(2.1) \quad P = (X - \xi_{k+1}) \cdots (X - \xi_n)Q.$$

By an inequality of M. Mignotte [6], we have

$$(2.2) \quad \left| |\xi_{k+1}| - 1 \right| \cdots \left| |\xi_n| - 1 \right| H(Q) < H(P).$$

We look to  $\mu > 0$ , which is a lower bound for the absolute values of the roots of  $P$ .



Inequalities on Polynomial  
Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 4 of 13

If  $\mu \geq 2$  then  $|\xi_{k+1}| - 1, \dots, |\xi_n| - 1 \geq 1$  and by (2.2) we obtain

$$(2.3) \quad H(Q) < H(P).$$

If  $\mu < 2$  we associate the polynomials

$$P_\mu(X) = (2/\mu)^n P(\mu X/2) = X^n + a_{n-1} 2/\mu X^{n-1} + \dots + a_0 (2/\mu)^n,$$

$$Q_\mu(X) = (2/\mu)^k Q(\mu X/2) = X^k + b_{k-1} 2/\mu X^{k-1} + \dots + b_0 (2/\mu)^k.$$

Let  $\eta_1, \dots, \eta_n \in \mathbb{C}$  be the roots of  $P_\mu$ . Then

$$|\eta_i| = 2|\xi_i|/\mu \geq 2$$

and from (2.2) it follows that

$$H(Q) = \max_j |b_j| \leq \max_j |b_j (2/\mu)^{k-j}| = H(Q_\mu) < H(P_\mu) = \max_i |a_i (2/\mu)^{n-i}|.$$


Therefore

$$(2.4) \quad H(Q) < \max_i \left| a_i \left( \frac{2}{\mu} \right)^{n-i} \right|.$$

□

**Corollary 2.2.** *If  $\mu < 2$  is a lower bound of the moduli of the roots of the complex polynomial  $P$  and  $Q$  is proper divisor of  $P$  in  $[X]$ , then*

$$H(Q) < (2/\mu)^n H(P).$$



## Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 5 of 13

*Proof.* Note that  $2/\mu > 1$ . Therefore, by Proposition 2.1,

$$\begin{aligned} H(Q) &< \max_{0 \leq i \leq n} \left| a_i \left( \frac{2}{\mu} \right)^{n-i} \right| \\ &< \max_{0 \leq i \leq n} \left( \frac{2}{\mu} \right)^{n-i} \cdot H(P) \\ &= \left( \frac{2}{\mu} \right)^n H(P). \end{aligned}$$

□

*Example 2.1.* Consider  $P(X) = X^6 + 2X^5 + 5X^4 + 10X^3 + 21X^2 + 42X + 83$ . By the criterion of Eneström–Kakeya (see M. Marden [4], p. 137), the zeros of a polynomial  $a_0 + a_1X + \dots + a_nX^n$  with real positive coefficients lie in the ring


$$\min\{a_i/a_{i+1}\} \leq |z| \leq \max\{a_i/a_{i+1}\}.$$

Thus, the roots of the polynomial  $P$  are in the ring  $83/42 \leq |z| \leq 5/2$ . If  $Q$  is a divisor of  $P$  we obtain

$$H(Q) < 89.183 \quad \text{by Proposition 2.1,}$$

$$H(Q) < 602.455 \quad \text{by B. Beauzamy [1].}$$

**Proposition 2.3.** Let  $P(X) = a_nX^n + a_{n-1}X^{n-1} + \dots + a_1X + a_0 \in \mathbb{C}[X] \setminus \mathbb{C}$  such that  $P(0) \neq 0$  and let  $\nu > 0$  be an upper bound for the absolute values of



## Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 6 of 13

the roots of  $P$ . If  $Q$  is a proper divisor of  $P$  in  $[X]$ ,  $Q(0) = c_0$ , then

$$H(Q) < \left| \frac{c_0}{a_0} \right| \cdot H(P), \quad \text{if } \nu \leq \frac{1}{2},$$

$$H(Q) < \left| \frac{c_0}{a_0} \right| \cdot \max_{0 \leq i \leq n} |a_i (2\nu)^i|, \quad \text{if } \nu > \frac{1}{2}.$$

*Proof.* Let  $P^*$  and  $Q^*$  be the reciprocal polynomials of  $P$  and  $Q$  respectively. Considering

$$P_1(X) = \frac{1}{a_0} P^*(X) = \frac{1}{a_0} X^n P\left(\frac{1}{X}\right)$$

it is possible to obtain information about the heights with respect to the upper bounds of the moduli of the roots of  $P$ . The polynomial  $P_1$  is monic and

$$\frac{1}{\nu} \text{ is a lower bound for the roots of } P_1.$$

On the other hand, if  $P_1(X) = \sum_{i=0}^n b_i X^i$ , then  $b_i = a_{n-i}/a_0$  and


$$\max_{0 \leq i \leq n} \left| b_i \left( \frac{2}{\nu^{-1}} \right)^{n-i} \right| = \frac{1}{|a_0|} \max_{0 \leq i \leq n} |a_i (2\nu)^i|.$$

Let  $Q \in \mathbb{C}[X]$  be a proper divisor of  $P$ . Therefore

$$Q_1(X) = \frac{1}{c_0} Q^*(X) = \frac{1}{c_0} X^{\deg(Q)} Q(X^{-1})$$

is a proper divisor of  $P_1$ . However,

$$H(Q_1) = \frac{1}{|c_0|} H(Q).$$



## Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 7 of 13

If  $\nu \leq 1/2$ , by Proposition 2.1 we have  $H(Q_1) < H(P_1)$ , hence

$$\frac{1}{|c_0|} \cdot H(Q) < \frac{1}{|a_0|} \cdot H(P),$$

which gives the first inequality. The second relation in Proposition 2.1 gives the other inequality.  $\square$

*Remark 2.1.* If  $|P(0)| = 1$ , the inequalities in Proposition 2.3 become

$$H(Q) < H(P), \quad \text{if } \nu \leq \frac{1}{2},$$

$$H(Q) < \max_{0 \leq i \leq n} |a_i (2\nu)^i|, \quad \text{if } \nu > \frac{1}{2}.$$

Indeed, if  $|P(0)| = 1$  we have  $|a_0| = |c_0| = 1$ .

The same inequalities are valid if  $P \in [X]$  and  $Q$  is a proper divisor of  $P$  over  $\mathbb{Z}$ . In this case  $a_0$  and  $c_0$  are integers and  $c_0$  divides  $a_0$ , therefore  $|\frac{c_0}{a_0}| \leq 1$ .


As in Corollary 2.2 we deduce

**Corollary 2.4.** *If  $\nu > \frac{1}{2}$  is an upper bound of the moduli of the roots of the complex polynomial  $P$  and  $Q$  is proper divisor of  $P$  in  $\mathbb{C}[X]$ , then*

$$H(Q) < \left| \frac{c_0}{a_0} \right| (2\nu)^n H(P).$$

*Example 2.2.* Let  $P = 2381X^5 - 597X^4 - 150X^3 - 37X^2 + 9X + 2$ .

If  $z \in \mathbb{C}$  is a root of  $P$ , then  $|z| \leq 2 \max \left| \frac{a_{i-1}}{a_i} \right| = \frac{100}{199}$ , by the criterion of T.



### Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 8 of 13



Kojima [3]. If  $Q$  is a possible divisor of  $P$ , then

$$H(Q) < 2440.403 \quad \text{By Proposition 2.3,}$$

$$H(Q) < 10745.533 \quad \text{by B. Beuzamy [1].}$$

*Remark 2.2.* The estimates from Propositions 2.1 and 2.3 apply to any complex polynomial, while the estimate of Beuzamy refers only to integer polynomials.

*Example 2.3.* Let  $P = 381X^5 - 95iX^4 + (45 - 9i)X^3 + 17iX^2 + (2 + 7i)X + 3$ .

If  $z \in \mathbb{C}$  is a root of  $P$ , then  $|z| \leq 2 \max_i \frac{|a_{i-1}|}{|a_i|} = \frac{\sqrt{2106}}{95}$ , again by the criterion of T. Kojima [3]. If  $Q$  is a possible divisor of  $P$ , then

$$H(Q) < 320.703 \quad \text{By Proposition 2.3,}$$

$$H(Q) < 2374.689 \quad \text{by M. Mignotte [5].}$$

*Example 2.4.* Let  $P = 127X^7 + 64X^6 + 32X^5 + 16X^4 + 8X^3 + 4X^2 + 2X + 1$ .


The roots of  $P$  are in the ring  $1/2 \leq |z| \leq 64/127$ , by the criterion of Eneström–Kakeya (M. Marden, loc. cit.). If  $Q$  is a possible divisor of  $P$ , then

$$H(Q) < 134.167 \quad \text{By Proposition 2.3,}$$

$$H(Q) < 1472.464 \quad \text{by Beuzamy.}$$

**Corollary 2.5.** Let  $P[X] = \sum_{i=0}^n a_i X^i \in \mathbb{R}[X] \setminus \mathbb{R}$  be monic and let  $Q \in [X]$  be a proper monic divisor of  $P$ . If  $a_i > 0$  for all  $i$ , there exists  $\mu \in \mathbb{R}$  such that  $2^{1-\frac{1}{n}} < \mu < 2$  and  $a_{i-1} \geq \mu a_i$  for  $i = 1, 2, \dots, n$ , then

$$H(Q) < 2 H(P).$$




---

### Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

---

Title Page

Contents



Go Back

Close

Quit

Page 9 of 13

*Proof.* From the condition  $a_{i-1} \geq \mu a_i$  for all  $i$  and the criterion of Eneström–Kakeya (M. Marden, loc. cit.) it follows that  $\mu$  is a lower bound for the absolute values of the roots of  $P$ . Because  $\mu < 2$ , the inequality (2.4) from Proposition 2.1 is verified. Noting that

$$\frac{1}{2} < \frac{1}{\mu} < 2^{\frac{1}{n}-1}$$


we obtain

$$\begin{aligned} H(Q) &< \max_{1 \leq i \leq n} |a_i| \cdot \left| \left( \frac{2}{\mu} \right)^{n-i} \right| \\ &= H(P) \cdot \max_{1 \leq i \leq n} \left( \frac{2}{\mu} \right)^{n-i} \\ &< H(P) \cdot \max_{1 \leq i \leq n} 2^{\frac{n-i}{n}} < 2H(P). \end{aligned}$$

□

*Example 2.5.* Let  $P(X) = X^7 + 2X^6 + 4X^5 + 8X^4 + 15X^3 + 28X^2 + 51X + 92$ . Then, by the theorem of Eneström–Kakeya (M. Marden loc. cit.),  $\mu = 51/28$  is a lower bound for the absolute values of roots of  $P$ . We have  $2^{1-\frac{1}{7}} < 51/28 < 2$ , so the hypotheses of Corollary 2.5 are fulfilled.

*Remark 2.3.* The same conclusion as in Corollary 2.5 holds if  $a_i > 0$  and there exists  $\nu > 0$  such that  $a_{i-1} \leq \nu a_i$  and  $\frac{1}{2} < \nu \leq 2^{\frac{1}{n}-1}$ .



## Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 10 of 13

### 3. A Smallest Divisor

In this section we give a limit for the smallest height of a proper divisor of a complex polynomial.

**Theorem 3.1.** *Let  $P$  be a nonconstant complex polynomial and suppose that its roots  $\xi_1, \dots, \xi_n \in \mathbb{C}$  are such that*

$$|\xi_1| \geq \dots \geq |\xi_k| > 1 \geq |\xi_{k+1}| \geq \dots \geq |\xi_n|.$$

If  $P = P_1 P_2$  is a factorization of  $P$  over  $\mathbb{Z}$ , we have

$$H(P) > \min\{H(P_1), H(P_2)\} \cdot (|\xi_k| - 1)^{k/2} \cdot (1 - |\xi_{k+1}|)^{(n-k)/2}.$$

*Proof.* We observe that


$$H(P) > H(P_1) \prod_{s \in J_1} (|\xi_s| - 1)$$

and

$$H(P) > H(P_2) \prod_{t \in J_2} (|\xi_t| - 1),$$

where  $\{J_1, J_2\}$  is a partition of  $\{1, 2, \dots, n\}$ . It follows that

$$H(P)^2 > H(P_1) H(P_2) \prod_{j=1}^n (|\xi_j| - 1),$$



Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back

Close

Quit

Page 11 of 13

therefore

$$H(P) > \min(H(P_1), H(P_2)) \cdot \sqrt{\prod_{j=1}^n ||\xi_j| - 1|}.$$

We finally observe that

$$\prod_{j=1}^n ||\xi_j| - 1| \geq (|\xi_k| - 1)^{k/2} \cdot (1 - |\xi_{k+1}|)^{(n-k)/2},$$

which ends the proof.  $\square$

*Example 3.1.* Let  $P(X) = X^5 - 49/6 X^4 + 59/3 X^3 - 6X^2 - 18X - 9/2$ . We have  $\xi_1 = \xi_2 = \xi_3 = 3$ ,  $\xi_4 = -1/2$  and  $\xi_5 = -1/3$ . Therefore

$$H(P) > \min\{H(P_1), H(P_2)\} \cdot (3 - 1)^{3/2} \cdot (1 - 1/2)^{2/2}.$$


But  $(3 - 1)^{3/2} \cdot (1 - 1/2)^{2/2} > 1.415$ , hence

$$\min\{H(P_1), H(P_2)\} < \frac{1}{1.415} H(P) < \frac{8}{11} H(P).$$

*Remark 3.1.* The index  $k$  from Theorem 3.1 can be computed by the Schur–Cohn criterion (see M. Marden [4], p. 198). Since it is usually not possible to find the roots  $\xi_k$  and  $\xi_{k+1}$ , we need to know lower bounds for roots outside the unit circle, respectively upper bounds for roots outside the unit circle.

**Corollary 3.2.** *If  $P$  has no roots on the unit circle, we have*

$$\min\{H(P_1), H(P_2)\} < H(P) / (|\xi_k| - 1)^{k/2} \cdot (1 - |\xi_{k+1}|)^{(n-k)/2}.$$



## Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

Title Page

Contents



Go Back


Close

Quit

Page 12 of 13

## References

- [1] B. BEAUZAMY, Products of polynomials and a priori estimates for coefficients in polynomial decompositions: A sharp result, *J. Symbolic Comput.*, **13** (1992), 463–472.
- [2] B. BEAUZAMY, E. BOMBIERI, P. ENFLO AND H. MONTGOMERY, Products of polynomials in many variables, *J. Number Theory*, **36** (1990), 219–245.
- [3] T. KOJIMA, On the limits of the roots of an algebraic equation, *Tôhoku Math. J.*, **11** (1917), 119-127.
- [4] M. MARDEN, *Geometry of Polynomials*, AMS Surveys **3**, 4th edition, Providence, Rhode Island (1989).
- [5] M. MIGNOTTE, An inequality about factors of polynomials, *Math. Comp.*, **28** (1974), 1153 – 1157.
- [6] M. MIGNOTTE, An inequality on the greatest roots of a polynomial, *Elem. d. Math.*, (1991), 86-87.
- [7] L. PANAITOPOL AND D. ȘTEFĂNESCU, New bounds for factors of integer polynomials, *J. UCS*, **1**(8) (1995), 599–609.



---

### Inequalities on Polynomial Heights

Laurențiu Panaitopol and  
Doru Ștefănescu

---

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13