

Journal of Inequalities in Pure and Applied Mathematics

THE INEQUALITIES $G \leq L \leq I \leq A$ IN n VARIABLES

ZHEN-GANG XIAO AND ZHI-HUA ZHANG

Department of Mathematics,
Hunan Institute of Science and Technology,
Yueyang City, Hunan 414006,
CHINA.

E-Mail: xgzgzh@163.com

Zixing Educational Research Section,
Chenzhou City,
Hunan 423400, CHINA.

E-Mail: zxzh1234@163.com

©2000 Victoria University
ISSN (electronic): 1443-5756
110-02



volume 4, issue 2, article 39,
2003.

*Received 21 October, 2002;
accepted 28 March, 2003.*

Communicated by: F. Qi

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

In the short note, the inequalities $G \leq L \leq I \leq A$ for the geometric, logarithmic, identric, and arithmetic means in n variables are proved.


2000 Mathematics Subject Classification: 26D15.

Key words: Inequality, Arithmetic mean, Geometric mean, Logarithmic mean, Identric mean, n variables, Van der Monde determinant.

The authors would like to thank Professor Feng Qi and the anonymous referee for some valuable suggestions which have improved the final version of this paper.

Contents

1	Introduction	3
2	Definitions and the Main Result	4
3	Proof of Theorem 2.1	6



**The Inequalities $G \leq L \leq I \leq A$
in n Variables**

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 2 of 13

1. Introduction

For positive numbers a_i , $1 \leq i \leq 2$, let

$$(1.1) \quad A = A(a_1, a_2) = \frac{a_1 + a_2}{2};$$

$$(1.2) \quad G = G(a_1, a_2) = \sqrt{a_1 a_2};$$


$$(1.3) \quad I = I(a_1, a_2) = \begin{cases} \exp \left[\frac{a_2 \ln a_2 - a_1 \ln a_1}{a_2 - a_1} - 1 \right], & a_1 < a_2, \\ a_1, & a_1 = a_2; \end{cases}$$

$$(1.4) \quad L = L(a_1, a_2) = \begin{cases} \frac{a_2 - a_1}{\ln a_2 - \ln a_1}, & a_1 < a_2, \\ a_1, & a_1 = a_2. \end{cases}$$

These are respectively called the arithmetic, geometric, identric, and logarithmic means.

The logarithmic mean [1, 3], somewhat surprisingly, has applications to the economical index analysis. K.B. Stolarsky first introduced the identric mean I and proved $G \leq L \leq I \leq A$ in two variables in [4]. See [2] also.

The purpose of this short note is to prove the inequalities $G \leq L \leq I \leq A$ of the geometric, logarithmic, identric, and arithmetic means in n variables.



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

2. Definitions and the Main Result

Let $a = (a_1, a_2, \dots, a_n)$ and $a_i > 0$ for $1 \leq i \leq n$, then the arithmetic, geometric, identric, and logarithmic means in n variables are defined respectively as follows

$$(2.1) \quad A = A(a) = A(a_1, \dots, a_n) = \frac{a_1 + a_2 + \dots + a_n}{n},$$


$$(2.2) \quad G = G(a) = G(a_1, \dots, a_n) = \sqrt[n]{a_1 a_2 \dots a_n},$$

$$(2.3) \quad I = I(a) = I(a_1, \dots, a_n) \\ = \exp \left[\frac{1}{V(a)} \sum_{i=1}^n (-1)^{n+i} a_i^{n-1} V_i(a) \ln a_i - m \right],$$

$$(2.4) \quad L = L(a) = L(a_1, \dots, a_n) = \frac{(n-1)!}{V(\ln a)} \sum_{i=1}^n (-1)^{n+i} a_i V_i(\ln a),$$

where $\ln a = (\ln a_1, \dots, \ln a_n)$, $a_i \neq a_j$ for $i \neq j$,

$$(2.5) \quad V(a) = \begin{vmatrix} 1 & 1 & \dots & 1 \\ a_1 & a_2 & \dots & a_n \\ a_1^2 & a_2^2 & \dots & a_n^2 \\ \dots & \dots & \dots & \dots \\ a_1^{n-1} & a_2^{n-1} & \dots & a_n^{n-1} \end{vmatrix} = \prod_{1 \leq j < i \leq n} (a_i - a_j)$$



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 4 of 13

is the determinant of Van der Monde's matrix of the n -th order,

$$(2.6) \quad V_i(a) = \begin{vmatrix} 1 & 1 & \dots & 1 & 1 & \dots & 1 \\ a_1 & a_2 & \dots & a_{i-1} & a_{i+1} & \dots & a_n \\ a_1^2 & a_2^2 & \dots & a_{i-1}^2 & a_{i+1}^2 & \dots & a_n^2 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ a_1^{n-2} & a_2^{n-2} & \dots & a_{i-1}^{n-2} & a_{i+1}^{n-2} & \dots & a_n^{n-2} \end{vmatrix},$$


$m = \sum_{k=1}^{n-1} \frac{1}{k}$, and $1 \leq i \leq n$.

The main result of this short note can be stated as

Theorem 2.1. *Let $a = (a_1, a_2, \dots, a_n)$ and $a_i > 0$ for $1 \leq i \leq n$, then the inequalities*

$$(2.7) \quad G(a) \leq L(a) \leq I(a) \leq A(a)$$

of the geometric, logarithmic, identric, and arithmetic means in n variables hold. The equalities in (2.7) are valid if and only if $a_1 = a_2 = \dots = a_n$.



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 5 of 13

3. Proof of Theorem 2.1

To prove inequalities in (2.7), we introduce the following means

$$(3.1) \quad I_r(a) = \prod_{\substack{i_1+i_2+\dots+i_n=n+r-1 \\ i_1, i_2, \dots, i_n \geq 1}} \left[\sum_{k=1}^n \frac{i_k}{n+r-1} a_k \right]^{\frac{1}{\binom{n+r-2}{r-1}}},$$

$$(3.2) \quad I'_r(a) = \prod_{\substack{i_1+i_2+\dots+i_n=r \\ i_1, i_2, \dots, i_n \geq 0}} \left[\sum_{k=1}^n \frac{i_k}{r} a_k \right]^{\frac{1}{\binom{n+r-1}{r}}},$$


$$(3.3) \quad L_r(a) = \frac{1}{\binom{n+r-1}{r}} \sum_{\substack{i_1+i_2+\dots+i_n=r \\ i_1, i_2, \dots, i_n \geq 0}} \prod_{k=1}^n a_k^{i_k/r},$$

$$(3.4) \quad L'_r(a) = \frac{1}{\binom{n+r-2}{r-1}} \sum_{\substack{i_1+i_2+\dots+i_n=n+r-1 \\ i_1, i_2, \dots, i_n \geq 1}} \prod_{k=1}^n a_k^{i_k/(n+r-1)},$$

where $a = (a_1, a_2, \dots, a_n)$ and $a_i > 0$ for $1 \leq i \leq n$.

Lemma 3.1. *Let $a = (a_1, a_2, \dots, a_n)$ and $a_i > 0$ for $1 \leq i \leq n$, then we have*

1. $I_1(a) = L_1(a) = A(a)$;
2. $I'_1(a) = L'_1(a) = G(a)$;



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 6 of 13

3. For $1 \leq j \leq n - 1$ and

$$(3.5) \quad \pi_j = \prod_{\substack{i_1+i_2+\dots+i_n=n+r-1 \\ i_{k_1}=i_{k_2}=\dots=i_{k_j}=0, \text{ for the rest } i_k \geq 1}} \sum_{k=1}^n \frac{i_k}{n+r-1} a_k,$$

we have

$$(3.6) \quad I'_{n+r-1}(a) = \prod_{j=1}^{n-1} \pi_j^{1/\binom{2n+r-2}{n+r-1}} [I_r(a)]^{\binom{n+r-2}{r-1}/\binom{2n+r-2}{n+r-1}};$$

4. For $1 \leq j \leq n - 1$ and

$$(3.7) \quad \delta_j = \sum_{\substack{i_1+i_2+\dots+i_n=n+r-1 \\ i_{k_1}=i_{k_2}=\dots=i_{k_j}=0, \text{ for the rest } i_k \geq 1}} \prod_{k=1}^n a_k^{i_k/(n+r-1)},$$


we have

$$(3.8) \quad L_{n+r-1}(a) = \frac{1}{\binom{2n+r-2}{n+r-1}} \left[\sum_{j=1}^{n-1} \delta_j + \binom{n+r-2}{r-1} L'_r(a) \right];$$

5. If $r \in \mathbb{N}$, then

(a) $I_r(a) \geq I_{r+1}(a)$,

(b) $I'_r(a) \leq I'_{r+1}(a)$,



**The Inequalities $G \leq L \leq I \leq A$
in n Variables**

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 7 of 13


- (c) $L_r(a) \geq L_{r+1}(a)$,
- (d) $L'_r(a) \leq L'_{r+1}(a)$,
- (e) $I_r(a) \geq L_r(a)$,
- (f) $I'_r(a) \geq L'_r(a)$,

where equalities above hold if and only if $a_1 = a_2 = \dots = a_n$.

Proof. The formula (3.6) follows from standard arguments and formulas (3.1) and (3.2).

If $r \in \mathbb{N}$ and $i_1 + i_2 + \dots + i_n = n + r - 1$, then

$$\begin{aligned}
 \sum_{k=1}^n \frac{i_k}{n+r-1} a_k &= \sum_{k=1}^n \frac{i_k}{n+r} \cdot \frac{n+r}{n+r-1} a_k \\
 &= \frac{n+r}{n+r-1} \sum_{k=1}^n \frac{i_k}{n+r} a_k \\
 &= \frac{1}{n+r-1} \left[\sum_{j=1}^n i_j + 1 \right] \sum_{k=1}^n \frac{i_k}{n+r} a_k \\
 &= \frac{1}{n+r-1} \left[\sum_{j=1}^n i_j \sum_{k=1}^n \frac{i_k}{n+r} a_k + \sum_{k=1}^n \frac{i_k}{n+r} a_k \right] \\
 &= \frac{1}{n+r-1} \left[\sum_{j=1}^n i_j \sum_{k=1}^n \frac{i_k}{n+r} a_k + \sum_{j=1}^n \frac{i_j}{n+r} a_j \right]
 \end{aligned}$$



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 8 of 13


$$= \sum_{j=1}^n \frac{i_j}{n+r-1} \left[\sum_{k=1}^n \frac{i_k a_k}{n+r} + \frac{a_j}{n+r} \right].$$

By using the weighted arithmetic-geometric mean inequality, we have

$$\sum_{k=1}^n \frac{i_k}{n+r-1} a_k \geq \prod_{j=1}^n \left[\sum_{k=1}^n \frac{i_k a_k}{n+r} + \frac{a_j}{n+r} \right]^{i_j/(n+r-1)},$$

and then

$$\begin{aligned} (3.9) \quad & \prod_{\substack{i_1+i_2+\dots+i_n=n+r-1 \\ i_1, i_2, \dots, i_n \geq 1}} \sum_{k=1}^n \frac{i_k}{n+r-1} a_k \\ & \geq \prod_{\substack{i_1+i_2+\dots+i_n=n+r-1 \\ i_1, i_2, \dots, i_n \geq 1}} \prod_{j=1}^n \left[\sum_{k=1}^n \frac{i_k a_k}{n+r} + \frac{a_j}{n+r} \right]^{i_j/(n+r-1)} \\ & = \prod_{j=1}^n \prod_{\substack{i_1+i_2+\dots+i_n=n+r-1 \\ i_1, i_2, \dots, i_n \geq 1}} \left[\sum_{k=1}^n \frac{i_k a_k}{n+r} + \frac{a_j}{n+r} \right]^{i_j/(n+r-1)} \\ & = \prod_{j=1}^n \prod_{\substack{\nu_1+\nu_2+\dots+\nu_n=n+r \\ \nu_1, \nu_2, \dots, \nu_{j-1}, \nu_{j+1}, \dots, \nu_n \geq 1; \nu_j \geq 2}} \left[\sum_{k=1}^n \frac{\nu_k}{n+r} a_k \right]^{(\nu_j-1)/(n+r-1)} \\ & = \prod_{\substack{\nu_1+\nu_2+\dots+\nu_n=n+r \\ \nu_1, \nu_2, \dots, \nu_n \geq 1}} \left[\sum_{k=1}^n \frac{\nu_k}{n+r} a_k \right]^{\sum_{j=1}^n (\nu_j-1)/(n+r-1)} \end{aligned}$$



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents




Go Back

Close

Quit

Page 9 of 13



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 10 of 13

$$\begin{aligned}
 &= \prod_{\substack{\nu_1 + \nu_2 + \dots + \nu_n = n+r \\ \nu_1, \nu_2, \dots, \nu_n \geq 1}} \left[\sum_{k=1}^n \frac{\nu_k}{n+r} a_k \right]^{(\sum_{j=1}^n \nu_j - n) / (n+r-1)} \\
 &= \prod_{\substack{\nu_1 + \nu_2 + \dots + \nu_n = n+r \\ \nu_1, \nu_2, \dots, \nu_n \geq 1}} \left[\sum_{k=1}^n \frac{\nu_k}{n+r} a_k \right]^{r / (n+r-1)} \\
 &= \prod_{\substack{\nu_1 + \nu_2 + \dots + \nu_n = n+r \\ \nu_1, \nu_2, \dots, \nu_n \geq 1}} \left[\sum_{k=1}^n \frac{\nu_k}{n+r} a_k \right]^{(\binom{n+r-2}{r-1}) / (\binom{n+r-1}{r})},
 \end{aligned}$$

notice that the result from line 4 to line 5 in (3.9) follows from a simple fact that

$$\left[\sum_{k=1}^n \frac{\nu_k}{n+r} a_k \right]^{(\nu_j - 1) / (n+r-1)} = 1 \text{ for } \nu_j = 1.$$

The equalities above are valid if and only if

$$\sum_{k=1}^n \frac{i_k a_k}{n+r} + \frac{a_1}{n+r} = \sum_{k=1}^n \frac{i_k a_k}{n+r} + \frac{a_2}{n+r} = \dots = \sum_{k=1}^n \frac{i_k a_k}{n+r} + \frac{a_n}{n+r},$$

which is equivalent to $a_1 = a_2 = \dots = a_n$. This implies that $I_r(a) \geq I_{r+1}(a)$.

The inequality $I_r(a) \geq L_r(a)$ follows easily from the generalized Hölder inequality

$$(3.10) \quad \prod_{\substack{i_1 + i_2 + \dots + i_n = r+1 \\ i_1 \geq 1, i_2 \geq 1, \dots, i_n \geq 1}} \left[\sum_{j=1}^n i_j a_j \right]^{\frac{1}{r}} \geq \sum_{\substack{i_1 + i_2 + \dots + i_n = r \\ i_1 \geq 0, i_2 \geq 0, \dots, i_n \geq 0}} \left[\prod_{j=1}^n a_j^{i_j} \right]^{\frac{1}{r}}.$$

The proofs of other formulas and inequalities will be left to the readers. \square

Lemma 3.2. Let $a = (a_1, a_2, \dots, a_n)$ and $a_i > 0$ for $1 \leq i \leq n$, then we have


1. $\lim_{r \rightarrow \infty} I_r(a) = \lim_{r \rightarrow \infty} I'_r(a) = I(a)$,
2. $\lim_{r \rightarrow \infty} L_r(a) = \lim_{r \rightarrow \infty} L'_r(a) = L(a)$.

Proof. It is easy to see that $\lim_{r \rightarrow \infty} I_r(a) = \lim_{r \rightarrow \infty} I'_r(a)$ and $\lim_{r \rightarrow \infty} L_r(a) = \lim_{r \rightarrow \infty} L'_r(a)$, since

$$\lim_{r \rightarrow \infty} \pi_j^{1/(2n+r-2)} = 1, \quad \lim_{r \rightarrow \infty} \frac{1}{\binom{2n+r-2}{n+r-1}} \sum_{j=1}^{n-1} \delta_j = 0, \quad \lim_{r \rightarrow \infty} \frac{\binom{n+r-2}{r-1}}{\binom{2n+r-2}{n+r-1}} = 1.$$

Straightforward computation yields

$$\begin{aligned} & \ln \lim_{r \rightarrow \infty} I'_r(a) \\ &= \lim_{r \rightarrow \infty} \ln I'_r(a) \\ &= \lim_{r \rightarrow \infty} \frac{1}{\binom{n+r-1}{r}} \sum_{\substack{i_1+i_2+\dots+i_n=r \\ i_1, i_2, \dots, i_n \geq 0}} \ln \sum_{k=1}^n \frac{i_k}{r} a_k \\ &= (n-1)! \int \cdots \int_{\substack{x_1+x_2+\dots+x_{n-1} \leq 1 \\ x_\ell \geq 0, 1 \leq \ell \leq n-1}} \ln \left[\left(1 - \sum_{i=1}^{n-1} x_i \right) a_1 + \sum_{j=2}^n x_{j-1} a_j \right] dx_1 dx_2 \cdots dx_{n-1} \\ &= \frac{(n-1)!}{V(a)} \sum_{i=1}^n (-1)^{n+i} a_i^{n-1} V_i(a) (\ln a_i - m) = \ln I(a) \end{aligned}$$



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit


Page 11 of 13

and

$$\begin{aligned}
 (3.11) \quad & \lim_{r \rightarrow \infty} L_r(a) \\
 &= \lim_{r \rightarrow \infty} \frac{1}{\binom{n+r-1}{r}} \sum_{\substack{i_1+i_2+\dots+i_n=r \\ i_1, i_2, \dots, i_n \geq 0}} \prod_{k=1}^n a_k^{i_k/r} \\
 &= (n-1)! \int \cdots \int_{\substack{x_1+x_2+\dots+x_{n-1} \leq 1 \\ x_\ell \geq 0, 1 \leq \ell \leq n-1}} a_1^{1-\sum_{i=1}^{n-1} x_i} a_2^{x_1} \cdots a_n^{x_{n-1}} dx_1 dx_2 \cdots dx_{n-1} \\
 &= \frac{(n-1)!}{V(\ln a)} \sum_{i=1}^n (-1)^{n+i} a_i V_i(\ln a) = L(a).
 \end{aligned}$$

The proof is complete. □

Proof of Theorem 2.1. This follows from combination of Lemma 3.1 and Lemma 3.2. □



**The Inequalities $G \leq L \leq I \leq A$
in n Variables**

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back


Close

Quit

Page 12 of 13

References

- [1] E. LEACH AND M. SHOLANDER, Extended mean values, *Amer. Math. Monthly*, **85** (1978), 84–90.
- [2] A.O. PITTENGER, Two logarithmic mean in n variables, *Amer. Math. Monthly*, **92** (1985), 99–104.
- [3] G. PÓLYA AND G. SZEGÖ, *Isoperimetric Inequalities in Mathematical Physics*, Princeton University Press, Princeton, 1951.
- [4] K.B. STOLARSKY, Generalizations of the logarithmic mean, *Mag. Math.*, **48** (1975), 87–92.



The Inequalities $G \leq L \leq I \leq A$
in n Variables

Zhen-Gang Xiao and
Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13