



Sums of Squares and Products of Jacobsthal Numbers

Zvonko Čerin

Department of Mathematics

University of Zagreb

Bijenička 30

Zagreb 10 000

Croatia

cerin@math.hr

Abstract

We consider sums of squares of odd and even terms of the Jacobsthal sequence and sums of their products. We also study the analogous alternating sums. These sums are related to products of appropriate Jacobsthal numbers and several integer sequences. The formulas that we discover show that a certain translation property for these sums holds, so that in practice, only sums of initial values and the information where the summation begins are necessary.

1 Introduction

The Jacobsthal and Jacobsthal-Lucas sequences J_n and j_n are defined by the recurrence relations

$$J_0 = 0, \quad J_1 = 1, \quad J_n = J_{n-1} + 2J_{n-2} \quad \text{for } n \geq 2,$$

and

$$j_0 = 2, \quad j_1 = 1, \quad j_n = j_{n-1} + 2j_{n-2} \quad \text{for } n \geq 2.$$

In Sections 2–4 we consider sums of squares of odd and even terms of the Jacobsthal sequence and sums of their products. These sums have nice representations as products of appropriate Jacobsthal and Jacobsthal-Lucas numbers.

The numbers J_k appear as the integer sequence A001045 from [9] while the numbers j_k is A014551. The properties of these numbers are summarized in [7]. For the convenience of the reader we shall now explicitly define these sequences.

The first eleven terms of the sequence J_k are 0, 1, 1, 3, 5, 11, 21, 43, 85, 171 and 341. It is given by the formula $J_n = \frac{2^n - (-1)^n}{3}$.

The first eleven terms of the sequence j_k are 2, 1, 5, 7, 17, 31, 65, 127, 257, 511 and 1025. It is given by the formula $j_n = 2^n + (-1)^n$.

In the last three sections we look into the alternating sums of squares of odd and even terms of the Jacobsthal sequence and the alternating sums of products of two consecutive Jacobsthal numbers. These sums also have nice representations as products of appropriate Jacobsthal and Jacobsthal-Lucas numbers.

These formulas for ordinary sums and for alternating sums have been discovered with the help of a PC computer and all algebraic identities needed for the verification of our theorems can be easily checked in either Derive, Mathematica or Maple V. Running times of all these calculations are in the range of a few seconds.

Similar results for Fibonacci, Lucas, Pell, and Pell-Lucas numbers have recently been discovered by G. M. Gianella and the author in papers [1, 2, 3, 4, 5, 6]. They improved some results in [8].

2 Jacobsthal even squares

The following lemma is needed to accomplish the inductive step in the proof of the first part of our first theorem.

For any $n = 0, 1, 2, \dots$ let $\beta_n = \frac{2^{4n+2}-1}{3} = J_{4n+2}$, $\gamma_n = \frac{2^{4n+2}+1}{5} = \frac{j_{4n+2}}{5}$ and $\tau_n = \frac{2^{4n}-1}{15} = \frac{J_{4n}}{5}$.

Lemma 1. *For every $m \geq 0$ and $k \geq 0$ the following equality holds*

$$\begin{aligned} (\beta_{n+1} \gamma_{n+1} - \beta_n \gamma_n) J_{2k}^2 + 8(\beta_{n+1} \tau_{n+1} - \beta_n \tau_n) J_{2k} \\ = J_{2k+4n+4}^2 + J_{2k+4n+2}^2 - J_{4n+4}^2 - J_{4n+2}^2. \end{aligned} \quad (2.1)$$

Proof. (P1). Let $A = 2^{2k}$, $B = 2^{8n}$ and $C = 2^{4n}$. The difference of the left hand side and the right hand side of the relation (2.1) is equal to

$$\begin{aligned} \frac{544}{9} \left((-1)^{1+2k} + 1 \right) AB + \frac{40}{9} \left((-1)^{4n+2k} + (-1)^{1+4n} \right) (A+1)C \\ + \frac{272}{9} \left((-1)^{4k} + 2(-1)^{1+2k} + 1 \right) B + \frac{2}{9} \left((-1)^{8n} + (-1)^{1+8n+4k} \right). \end{aligned}$$

It is obvious that all coefficients in the above expression are zero and the proof is complete. \square

There are essentially three types of proofs in this paper that we indicate as **P1**, **P2** and **P3**.

In **P1** and **P2** we use the substitutions $A = 2^{2k}$, $B = 2^{8n}$, $C = 2^{4n}$ and $M = 2^k$, $P = (-1)^k$, respectively. We prove that the difference of the left hand side and the right hand side of the relation from the statement is equal to zero with algebraic simplification to an expression that is obviously vanishing. This part is easily done with the help of a computer (for example in Maple V) so that we shall only indicate the final expression.

In **P3** we argue by induction on n taking care first of the initial value $n = 0$ and then showing that it holds for $n = r + 1$ under the assumption that it is true for $n = r$.

The following lemma is needed to accomplish the inductive step in the proof of the second part of our first theorem.

For any $n = 0, 1, 2, \dots$ let $\eta_n = 2^{4n+4} + 1 = j_{4n+4}$.

Lemma 2. *For every $m \geq 0$ and $k \geq 0$ the following equality holds*

$$\begin{aligned} (\tau_{n+2} \eta_{n+1} - \tau_{n+1} \eta_n) J_{2k}^2 + 8 (\beta_{n+1} \tau_{n+2} - \beta_n \tau_{n+1}) J_{2k} \\ = J_{2k+4n+6}^2 + J_{2k+4n+4}^2 - J_{4n+6}^2 - J_{4n+4}^2. \end{aligned} \quad (2.2)$$

Proof. (**P1**).

$$\begin{aligned} \frac{8704}{9} \left((-1)^{1+2k} + 1 \right) AB + \frac{160}{9} \left((-1)^{4n+2k} + (-1)^{1+4n} \right) (A+1)C \\ + \frac{4352}{9} \left((-1)^{4k} + 2(-1)^{1+2k} + 1 \right) B + \frac{2}{9} \left((-1)^{8n} + (-1)^{1+8n+4k} \right). \end{aligned}$$

□

Theorem 1. *For every $m \geq 0$ and $k \geq 0$ the following equalities hold*

$$\sum_{i=0}^m J_{2k+2i}^2 = \sum_{i=0}^m J_{2i}^2 + \beta_n J_{2k} [\gamma_n J_{2k} + 8 \tau_n], \quad (2.3)$$

if $m = 2n$ and $n = 0, 1, 2, \dots$ and

$$\sum_{i=0}^m J_{2k+2i}^2 = \sum_{i=0}^m J_{2i}^2 + \tau_{n+1} J_{2k} [\eta_n J_{2k} + 8 \beta_n], \quad (2.4)$$

if $m = 2n + 1$ and $n = 0, 1, 2, \dots$

Proof of (2.3). (**P3**). The proof is by induction on n . When $n = 0$ we obtain

$$J_{2k}^2 = J_0^2 + \beta_0 J_{2k} [\gamma_0 J_{2k} + 8 \tau_0] = J_{2k} J_{2k} = J_{2k}^2,$$

because $J_0 = 0$, $\beta_0 = 1$, $\gamma_0 = 1$ and $\tau_0 = 0$.

Assume that the relation (2.3) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)} J_{2k+2i}^2 &= J_{2k+4r+2}^2 + J_{2k+4r+4}^2 + \sum_{i=0}^{2r} J_{2k+2i}^2 = J_{2k+4r+2}^2 + J_{2k+4r+4}^2 + \\ &\sum_{i=0}^{2r} J_{2i}^2 + \beta_r J_{2k} [\gamma_r J_{2k} + 8 \tau_r] = \sum_{i=0}^{2(r+1)} J_{2i}^2 + \beta_{r+1} J_{2k} [\gamma_{r+1} J_{2k} + 8 \tau_{r+1}], \end{aligned}$$

where the last step uses Lemma 1 for $n = r + 1$. Hence, (2.3) is true for $n = r + 1$ and the proof is completed. □

Proof of (2.4). (**P3**). When $n = 0$ we obtain

$$J_{2k}^2 + J_{2k+2}^2 = J_0^2 + J_2^2 + \tau_1 J_{2k} [\eta_0 J_{2k} + 8\beta_0] = 17 J_{2k}^2 + 8 J_{2k} + 1,$$

since $J_0 = 0$, $J_1 = 1$, $\beta_0 = 1$, $\eta_0 = 17$ and $\tau_1 = 1$. The above equality is equivalent to $J_{2k+2}^2 = 16 J_{2k}^2 + 8 J_{2k} + 1 = (4 J_{2k} + 1)^2$ which is true because it follows from the relation $J_{n+2} = 4 J_n + 1$.

Assume that the relation (2.4) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)+1} J_{2k+2i}^2 &= J_{2k+4r+4}^2 + J_{2k+4r+6}^2 + \sum_{i=0}^{2r+1} J_{2k+2i}^2 = J_{2k+4r+4}^2 + J_{2k+4r+6}^2 + \\ &\sum_{i=0}^{2r+1} J_{2i}^2 + \tau_{r+1} J_{2k} [\eta_r J_{2k} + 8\beta_r] = \sum_{i=0}^{2(r+1)+1} J_{2i}^2 + \tau_{r+2} J_{2k} [\eta_{r+1} J_{2k} + 8\beta_{r+1}], \end{aligned}$$

where the last step uses Lemma 2 for $n = r + 1$. □

3 Jacobsthal odd squares

The initial step in an inductive proof of the first part of our second theorem uses the following lemma.

Lemma 3. *For every $k \geq 0$ the following identity holds*

$$J_{2k+1}^2 = 16 J_{2k} J_{2k-2} + 8 J_{2k} + 1.$$

Proof. (**P2**). Let $M = 2^k$ and $P = (-1)^k$. The difference of the left and the right hand side is equal to $\frac{(P-1)(P+1)}{3} (8M^2 - 5P^2 + 3) = 0$. □

The initial step in an inductive proof of the second part of our second theorem uses the following lemma.

Lemma 4. *For every $k \geq 0$ the following identity holds*

$$J_{2k+1}^2 + J_{2k+3}^2 = 10 + 8 J_{2k} (34 J_{2k-2} + 15).$$

Proof. (**P2**). $10(P-1)(P+1)(4M^2 - 3P^2 + 1)$. □

The following lemma is needed to accomplish the inductive step in the proof of the first part of our second theorem.

For any $n = 0, 1, 2, \dots$ let $\pi_n = \frac{2^{4n+1}-1}{3} = J_{4n+1}$.

Lemma 5. *For every $m \geq 0$ and $k \geq 0$ the following equality holds*

$$\begin{aligned} 8 J_{2k} [2(\beta_{n+1} \gamma_{n+1} - \beta_n \gamma_n) J_{2k-2} + \beta_{n+1} \pi_{n+1} - \beta_n \pi_n] \\ = J_{2k+4n+5}^2 + J_{2k+4n+3}^2 - J_{4n+5}^2 - J_{4n+3}^2. \end{aligned} \quad (3.1)$$

Proof. (P1).

$$\begin{aligned} & \frac{5440}{9} \left((-1)^{1+2k} + 1 \right) AB + \frac{80}{9} \left((-1)^{4n+2k+1} + (-1)^{4n} \right) (A+1)C \\ & \quad + \frac{1088}{9} \left(4(-1)^{4k} + 5(-1)^{1+2k} + 1 \right) B + \frac{2}{9} \left((-1)^{8n} + (-1)^{8n+4k+1} \right). \end{aligned}$$

□

The following lemma is needed to accomplish the inductive step in the proof of the second part of our second theorem.

For any $n = 0, 1, 2, \dots$ let $\sigma_n = 5 J_{4n+3}$.

Lemma 6. *For every $m \geq 0$ and $k \geq 0$ the following equality holds*

$$\begin{aligned} & 8 J_{2k} [2(\tau_{n+2} \eta_{n+1} - \tau_n \eta_n) J_{2k-2} + \tau_{n+2} \sigma_{n+1} - \tau_{n+1} \sigma_n] \\ & \quad = J_{2k+4n+7}^2 + J_{2k+4n+5}^2 - J_{4n+7}^2 - J_{4n+5}^2. \end{aligned} \quad (3.2)$$

Proof. (P1).

$$\begin{aligned} & \frac{87040}{9} \left((-1)^{1+2k} + 1 \right) AB + \frac{320}{9} \left((-1)^{4n+2k+1} + (-1)^{4n} \right) (A+1)C \\ & \quad + \frac{17408}{9} \left(4(-1)^{4k} + 5(-1)^{1+2k} + 1 \right) B + \frac{2}{9} \left((-1)^{8n} + (-1)^{8n+4k+1} \right). \end{aligned}$$

□

Theorem 2. *For every $m \geq 0$ and $k \geq 0$ the following equalities hold*

$$\sum_{i=0}^m J_{2k+2i+1}^2 = \sum_{i=0}^m J_{2i+1}^2 + 8 \beta_n J_{2k} [2 \gamma_n J_{2k-2} + \pi_n], \quad (3.3)$$

if $m = 2n$ and $n = 0, 1, 2, \dots$ and

$$\sum_{i=0}^m J_{2k+2i+1}^2 = \sum_{i=0}^m J_{2i+1}^2 + 8 \tau_{n+1} J_{2k} [2 \eta_n J_{2k-2} + \sigma_n], \quad (3.4)$$

if $m = 2n + 1$ and $n = 0, 1, 2, \dots$

Proof of (3.3). (P3). When $n = 0$ we obtain

$$J_{2k+1}^2 = J_1^2 + 8 \beta_0 J_{2k} [2 \gamma_0 J_{2k-2} + \pi_0] = 1 + 8 J_{2k} [2 J_{2k-2} + 1],$$

because $J_1 = 1$, $\beta_0 = 1$, $\gamma_0 = 1$ and $\pi_0 = 1$. But, this equality is true by Lemma 3.

Assume that the relation (3.3) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)} J_{2k+2i+1}^2 &= J_{2k+4r+5}^2 + J_{2k+4r+3}^2 + \sum_{i=0}^{2r} J_{2k+2i+1}^2 = J_{2k+4r+5}^2 + J_{2k+4r+3}^2 + \sum_{i=0}^{2r} J_{2i+1}^2 \\ & \quad + 8 \beta_n J_{2k} [2 \gamma_n J_{2k-2} + \pi_n] = \sum_{i=0}^{2(r+1)} J_{2i+1}^2 + 8 \beta_{n+1} J_{2k} [2 \gamma_{n+1} J_{2k-2} + \pi_{n+1}], \end{aligned}$$

where the last step uses Lemma 5 for $n = r + 1$.

□

Proof of (3.4). (**P3**). The proof is by induction on n . When $n = 0$ we obtain

$$J_{2k+1}^2 + J_{2k+3}^2 = J_1^2 + J_3^2 + 8 \tau_1 J_{2k} [2 \eta_0 J_{2k-2} + \sigma_0] = 10 + 8 J_{2k} [34 J_{2k-2} + 15],$$

since $J_1 = 1$, $J_3 = 3$, $\tau_1 = 1$, $\eta_0 = 17$ and $\sigma_0 = 15$. The above equality is true by Lemma 4.

Assume that the relation (3.4) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)+1} J_{2k+2i+1}^2 &= J_{2k+4r+7}^2 + J_{2k+4r+5}^2 + \sum_{i=0}^{2r+1} J_{2k+2i+1}^2 = \\ &= J_{2k+4r+7}^2 + J_{2k+4r+5}^2 + \sum_{i=0}^{2r+1} J_{2i+1}^2 + 8 \tau_{r+1} J_{2k} [2 \eta_r J_{2k-2} + \sigma_r] \\ &= \sum_{i=0}^{2(r+1)+1} J_{2i+1}^2 + 8 \tau_{r+2} J_{2k} [2 \eta_{r+1} J_{2k-2} + \sigma_{r+1}], \end{aligned}$$

where the last step uses Lemma 6 for $n = r + 1$. □

4 Jacobsthal products

For the first two steps in a proof by induction of our next theorem we require the following lemma.

Lemma 7. *For every $k \geq 0$ the following equalities hold*

$$J_{2k+1} = 8 J_{2k-2} + 3. \tag{4.1}$$

$$J_{2k} J_{2k+1} + J_{2k+2} J_{2k+3} = 3 + J_{2k} (136 J_{2k-2} + 55). \tag{4.2}$$

Proof of (4.1). By the formula $J_k = \frac{2^k - (-1)^k}{3}$ we have

$$J_{2k+1} = \frac{2^{2k+1} - (-1)^{2k+1}}{3} = \frac{2^{2k+1} + 1}{3} = 8 \cdot \frac{2^{2k-2} - (-1)^{2k-2}}{3} + 3 = 8 J_{2k-2} + 3. \tag{4.3}$$

□

Proof of (4.2). (**P2**). $\frac{(P-1)(P+1)}{3} (55 M^2 - 46 P^2 + 9)$. □

With the following lemma we shall make the inductive step in the proof of the first part of our third theorem.

Lemma 8. *For every $m \geq 0$ and $k \geq 0$ the following equality holds*

$$\begin{aligned} J_{2k} [8 (\beta_{n+1} \gamma_{n+1} - \beta_n \gamma_n) J_{2k-2} + \beta_{n+1} \mu_{n+1} - \beta_n \mu_n] = \\ J_{2k+4n+5} J_{2k+4n+4} + J_{2k+4n+3} J_{2k+4n+2} - J_{4n+5} J_{4n+4} - J_{4n+3} J_{4n+2}. \end{aligned} \tag{4.3}$$

Proof. (P1).

$$\frac{2720}{9} \left((-1)^{1+2k} + 1 \right) AB + \frac{20}{9} \left((-1)^{4n+2k} + (-1)^{4n+1} \right) (A+1)C + \frac{544}{9} \left(4(-1)^{4k} + 5(-1)^{1+2k} + 1 \right) B + \frac{2}{9} \left((-1)^{8n+1} + (-1)^{8n+4k} \right).$$

□

With the following lemma we shall make the inductive step in the proof of the second part of our third theorem.

Lemma 9. *For every $m \geq 0$ and $k \geq 0$ the following equality holds*

$$J_{2k} [8(\tau_{n+2} \eta_{n+1} - \tau_{n+1} \eta_n) J_{2k-2} + \tau_{n+2} \nu_{n+1} - \tau_{n+1} \nu_n] = J_{2k+4n+5} J_{2k+4n+4} + J_{2k+4n+7} J_{2k+4n+6} - J_{4n+5} J_{4n+4} - J_{4n+7} J_{4n+6}. \quad (4.4)$$

Proof. (P1).

$$\frac{43520}{9} \left((-1)^{1+2k} + 1 \right) AB + \frac{80}{9} \left((-1)^{4n+2k} + (-1)^{4n+1} \right) (A+1)C + \frac{8704}{9} \left(4(-1)^{4k} + 5(-1)^{1+2k} + 1 \right) B + \frac{2}{9} \left((-1)^{8n+1} + (-1)^{8n+4k} \right).$$

□

For any $n = 0, 1, 2, \dots$ let $\mu_n = \frac{2^{4n+3}+1}{3} = J_{4n+3}$ and $\nu_n = \frac{5(2^{4n+5}+1)}{3} = 5 J_{4n+5}$.

Theorem 3. *For every $m \geq 0$ and $k \geq 0$ the following equalities hold*

$$\sum_{i=0}^m J_{2k+2i} J_{2k+2i+1} = \sum_{i=0}^m J_{2i} J_{2i+1} + \beta_n J_{2k} [8\gamma_n J_{2k-2} + \mu_n], \quad (4.5)$$

if $m = 2n$ and $n = 0, 1, 2, \dots$ and

$$\sum_{i=0}^m J_{2k+2i} J_{2k+2i+1} = \sum_{i=0}^m J_{2i} J_{2i+1} + \tau_{n+1} J_{2k} [8\eta_n J_{2k-2} + \nu_n], \quad (4.6)$$

if $m = 2n + 1$ and $n = 0, 1, 2, \dots$

Proof of (4.5). (P3). For $n = 0$ the relation (4.5) is

$$J_{2k} J_{2k+1} = J_0 J_1 + \beta_0 J_{2k} (8\gamma_0 J_{2k-2} + \mu_0) = J_{2k} (8 J_{2k-2} + 3)$$

which is true since $J_{2k+1} = 8 J_{2k-2} + 3$ by the relation (4.1) in Lemma 7.

Assume that the relation (4.5) is true for $n = r$. Then

$$\begin{aligned}
\sum_{i=0}^{2(r+1)} J_{2k+2i} J_{2k+2i+1} &= J_{2k+4r+4} J_{2k+4r+5} + J_{2k+4r+2} J_{2k+4r+3} + \sum_{i=0}^{2r} J_{2k+2i} J_{2k+2i+1} = \\
&J_{2k+4r+4} J_{2k+4r+5} + J_{2k+4r+2} J_{2k+4r+3} + \sum_{i=0}^{2r} J_{2i} J_{2i+1} + \beta_n J_{2k} [8\gamma_n J_{2k-2} + \mu_n] \\
&= \sum_{i=0}^{2(r+1)} J_{2i} J_{2i+1} + \beta_{n+1} J_{2k} [8\gamma_{n+1} J_{2k-2} + \mu_{n+1}],
\end{aligned}$$

where the last step uses Lemma 8 for $n = r + 1$. \square

Proof of (4.6). (**P3**). For $n = 0$ the relation (4.6) is

$$J_{2k} J_{2k+1} + J_{2k+2} J_{2k+3} = J_0 J_1 + J_2 J_3 + \tau_1 J_{2k} (8\eta_0 J_{2k-2} + \nu_0) = 3 + J_{2k} (136 J_{2k-2} + 55)$$

which is true by (4.2) in Lemma 7.

Assume that the relation (4.6) is true for $n = r$. Then

$$\begin{aligned}
\sum_{i=0}^{2(r+1)+1} J_{2k+2i} J_{2k+2i+1} &= J_{2k+4r+4} J_{2k+4r+5} + J_{2k+4r+6} J_{2k+4r+7} + \sum_{i=0}^{2r+1} J_{2k+2i} J_{2k+2i+1} = \\
&J_{2k+4r+4} J_{2k+4r+5} + J_{2k+4r+6} J_{2k+4r+7} + \sum_{i=0}^{2r+1} J_{2i} J_{2i+1} + \tau_{n+1} J_{2k} [8\eta_n J_{2k-2} + \nu_n] \\
&= \sum_{i=0}^{2(r+1)+1} J_{2i} J_{2i+1} + \tau_{n+2} J_{2k} [8\eta_{n+1} J_{2k-2} + \nu_{n+1}],
\end{aligned}$$

where the last step uses Lemma 9 for $n = r + 1$. \square

5 Alternating Jacobsthal even squares

In this section we look for formulas that give closed forms for alternating sums of squares of Jacobsthal numbers with even indices.

Lemma 10. *For every $k \geq 0$ we have*

$$J_{2k} = 4 J_{2k-2} + 1. \quad (5.1)$$

Proof. By the formula $J_k = \frac{2^k - (-1)^k}{3}$ we get

$$4 J_{2k-2} + 1 = 4 \cdot \frac{2^{2k-2} - (-1)^{2k-2}}{3} + 1 = \frac{2^{2k} - (-1)^{2k}}{3} = J_{2k}.$$

\square

Lemma 11. For every $k \geq 0$ we have

$$J_{2k+2}^2 - J_{2k}^2 = 1 + J_{2k} (60 J_{2k-2} + 23). \quad (5.2)$$

Proof. (P2). $\frac{(P-1)(P+1)}{3} (23 M^2 - 20 P^2 + 3)$. \square

Let $\tau_0^* = 1$ and $\tau_{n+1}^* - \tau_n^* = 2^{4n+3} (50 \cdot 2^{4n} - 1)$, for $n = 0, 1, 2, \dots$. For any $n = 0, 1, 2, \dots$ let $\gamma_n^* = \frac{2^{8n+4}+1}{17} = \frac{J_{8n+4}}{17}$.

Lemma 12. For every $k \geq 0$ and every $n \geq 0$ we have

$$J_{2k} (4 (\gamma_{n+1}^* - \gamma_n^*) J_{2k-2} + \tau_{n+1}^* - \tau_n^*) = J_{2k+4n+4}^2 - J_{2k+4n+2}^2 - J_{4n+4}^2 + J_{4n+2}^2. \quad (5.3)$$

Proof. (P1).

$$\begin{aligned} \frac{400}{3} \left((-1)^{1+2k} + 1 \right) AB + \frac{8}{3} \left((-1)^{4n+2k} - 1 \right) AC + \\ \frac{80}{3} \left(4 (-1)^{4k} + 5 (-1)^{1+2k} + 1 \right) B + \frac{8}{3} \left((-1)^{2k} + (-1)^{4n+1} \right) C. \end{aligned}$$

\square

Let $\beta_0^* = 23$ and $\beta_{n+1}^* - \beta_n^* = 2^{4n+5} (200 \cdot 2^{4n} - 1)$ for any $n = 0, 1, 2, \dots$. For the same values of n , let $\eta_n^* = \frac{2^{8n+8}-1}{17} = \frac{3}{17} J_{8n+8}$.

Lemma 13. For every $k \geq 0$ and every $n \geq 0$ we have

$$J_{2k} (4 (\eta_{n+1}^* - \eta_n^*) J_{2k-2} + \beta_{n+1}^* - \beta_n^*) = J_{2k+4n+6}^2 - J_{2k+4n+4}^2 + J_{4n+4}^2 - J_{4n+6}^2. \quad (5.4)$$

Proof. (P1).

$$\begin{aligned} \frac{6400}{3} \left((-1)^{1+2k} + 1 \right) AB + \frac{32}{3} \left((-1)^{4n+2k} - 1 \right) AC + \\ \frac{1280}{3} \left(4 (-1)^{4k} + 5 (-1)^{1+2k} + 1 \right) B + \frac{32}{3} \left((-1)^{2k} + (-1)^{4n+1} \right) C. \end{aligned}$$

\square

Theorem 4. For every $m \geq 0$ and $k \geq 0$ the following equalities hold

$$\sum_{i=0}^m (-1)^i J_{2k+2i}^2 = \sum_{i=0}^m (-1)^i J_{2i}^2 + J_{2k} [4 \gamma_n^* J_{2k-2} + \tau_n^*], \quad (5.5)$$

if $m = 2n$ and $n = 0, 1, 2, \dots$ and

$$\sum_{i=0}^m (-1)^i J_{2k+2i}^2 = \sum_{i=0}^m (-1)^i J_{2i}^2 - J_{2k} [4 \eta_n^* J_{2k-2} + \beta_n^*], \quad (5.6)$$

if $m = 2n + 1$ and $n = 0, 1, 2, \dots$

Proof of (5.5). **(P3).** For $n = 0$ the relation (5.5) is

$$J_{2k}^2 = J_0^2 + J_{2k} (4 \gamma_0^* J_{2k-2} + \tau_0^*) = J_{2k} (4 J_{2k-2} + 1)$$

(i. e., the relation (5.1) multiplied by J_{2k}) which is true by Lemma 10.
Assume that the relation (5.5) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)} (-1)^i \cdot J_{2k+2i}^2 &= Q_{2k+4r+4}^2 - Q_{2k+4r+2}^2 + \sum_{i=0}^{2r} (-1)^i \cdot J_{2k+2i}^2 \\ &= Q_{2k+4r+4}^2 - Q_{2k+4r+2}^2 + \sum_{i=0}^{2r} (-1)^i \cdot J_{2i}^2 + J_{2k} [4 \gamma_n^* J_{2k-2} + \tau_n^*] = \\ &\qquad\qquad\qquad \sum_{i=0}^{2(r+1)} (-1)^i \cdot J_{2i}^2 + J_{2k} [4 \gamma_{n+1}^* J_{2k-2} + \tau_{n+1}^*], \end{aligned}$$

where the last step uses Lemma 12. □

Proof of (5.6). **(P3).** For $n = 0$ the relation (5.6) is

$$J_{2k}^2 - J_{2k+2}^2 = (J_0^2 - J_2^2) - J_{2k} [4 \eta_0^* J_{2k-2} + \beta_0^*] = -1 - J_{2k} [60 J_{2k-2} + 23],$$

which is true by Lemma 11.

Assume that the relation (5.6) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)+1} (-1)^i \cdot J_{2k+2i}^2 &= J_{2k+4r+4}^2 - J_{2k+4r+6}^2 + \sum_{i=0}^{2r+1} (-1)^i \cdot J_{2k+2i}^2 \\ &= J_{2k+4r+4}^2 - J_{2k+4r+6}^2 + \sum_{i=0}^{2r+1} (-1)^i \cdot J_{2i}^2 - J_{2k} [4 \eta_n^* J_{2k-2} + \beta_n^*] \\ &\qquad\qquad\qquad = \sum_{i=0}^{2(r+1)+1} (-1)^i \cdot J_{2i}^2 - J_{2k} [4 \eta_{n+1}^* J_{2k-2} + \beta_{n+1}^*], \end{aligned}$$

where the last step uses Lemma 13. □

6 Alternating Jacobsthal odd squares

In this section we look for formulas that give closed forms for alternating sums of squares of Jacobsthal numbers with odd indices.

Lemma 14. *For every $k \geq 0$ we have*

$$J_{2k+1}^2 = 1 + 8 J_{2k} [2 J_{2k-2} + 1]. \tag{6.1}$$

Proof. **(P2).** $\frac{(P-1)(P+1)}{3} (8 M^2 - 5 P^2 + 3)$. □

Lemma 15. For every $k \geq 0$ we have

$$J_{2k+3}^2 - J_{2k+1}^2 = 8 + 8 J_{2k} (30 J_{2k-2} + 13). \quad (6.2)$$

Proof. (P2). $\frac{(P-1)(P+1)}{3} (13 M^2 - 10 P^2 + 3)$. □

Let $\tau_0^{**} = 1$ and $\tau_{n+1}^{**} - \tau_n^{**} = 101 \cdot 2^{4n+1} + 25 \cdot 2^{4n+3} (2^{4n} - 1)$ for $n = 0, 1, 2, \dots$

Lemma 16. For every $k \geq 0$ and every $n \geq 0$ we have

$$8 J_{2k} (2 (\gamma_{n+1}^* - \gamma_n^*) J_{2k-2} + \tau_{n+1}^{**} - \tau_n^{**}) = J_{2k+4n+5}^2 - J_{2k+4n+3}^2 - J_{4n+5}^2 + J_{4n+3}^2. \quad (6.3)$$

Proof. (P1).

$$\begin{aligned} & \frac{1600}{3} \left((-1)^{1+2k} + 1 \right) AB + \frac{16}{3} \left((-1)^{4n+2k+1} + 1 \right) AC + \\ & \quad \frac{320}{3} \left(4 (-1)^{4k} + 5 (-1)^{1+2k} + 1 \right) B + \frac{16}{3} \left((-1)^{2k+1} + (-1)^{4n} \right) C. \end{aligned}$$

□

Let $\beta_0^{**} = 13$ and $\beta_{n+1}^{**} - \beta_n^{**} = 401 \cdot 2^{4n+3} + 100 \cdot 2^{4n+5} (2^{4n} - 1)$ for any $n = 0, 1, 2, \dots$

Lemma 17. For every $k \geq 0$ and every $n \geq 0$ we have

$$8 J_{2k} (2 (\eta_{n+1}^* - \eta_n^*) J_{2k-2} + \beta_{n+1}^{**} - \beta_n^{**}) = J_{2k+4n+7}^2 - J_{2k+4n+5}^2 + J_{4n+5}^2 - J_{4n+7}^2. \quad (6.4)$$

Proof. (P1).

$$\begin{aligned} & \frac{25600}{3} \left((-1)^{1+2k} + 1 \right) AB + \frac{64}{3} \left((-1)^{4n+2k+1} + 1 \right) AC + \\ & \quad \frac{5120}{3} \left(4 (-1)^{4k} + 5 (-1)^{1+2k} + 1 \right) B + \frac{64}{3} \left((-1)^{2k+1} + (-1)^{4n} \right) C. \end{aligned}$$

□

Theorem 5. For every $m \geq 0$ and $k \geq 0$ the following equalities hold

$$\sum_{i=0}^m (-1)^i J_{2k+2i+1}^2 = \sum_{i=0}^m (-1)^i J_{2i+1}^2 + 8 J_{2k} [2 \gamma_n^* J_{2k-2} + \tau_n^{**}], \quad (6.5)$$

if $m = 2n$ and $n = 0, 1, 2, \dots$ and

$$\sum_{i=0}^m (-1)^i J_{2k+2i+1}^2 = \sum_{i=0}^m (-1)^i J_{2i+1}^2 - 8 J_{2k} [2 \eta_n^* J_{2k-2} + \beta_n^{**}], \quad (6.6)$$

if $m = 2n + 1$ and $n = 0, 1, 2, \dots$

Proof of (6.5). (P3). For $n = 0$ the relation (6.5) is

$$J_{2k+1}^2 = J_1^2 + 8 J_{2k} (2 \gamma_0^* J_{2k-2} + \tau_0^{**})$$

(i. e., the relation (6.1)) which is true by Lemma 14.

Assume that the relation (6.5) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)} (-1)^i J_{2k+2i+1}^2 &= J_{2k+4r+5}^2 - J_{2k+4r+3}^2 + \sum_{i=0}^{2r} (-1)^i J_{2k+2i+1}^2 \\ &= J_{2k+4r+5}^2 - J_{2k+4r+3}^2 + \sum_{i=0}^{2r} (-1)^i J_{2i+1}^2 + 8 J_{2k} [2 \gamma_n^* J_{2k-2} + \tau_n^{**}] \\ &= \sum_{i=0}^{2(r+1)} (-1)^i J_{2i+1}^2 + 8 J_{2k} [2 \gamma_{n+1}^* J_{2k-2} + \tau_{n+1}^{**}], \end{aligned}$$

where the last step uses Lemma 16. □

Proof of (6.6). (P3). The proof is again by induction on n . For $n = 0$ the relation (6.6) is $J_{2k+1}^2 - J_{2k+3}^2 = J_1^2 - J_3^2 - 8 J_{2k} [2 \eta_0^* J_{2k-2} + \beta_0^{**}]$ which is true by Lemma 15 since $\eta_0^* = 15$ and $\beta_0^{**} = 13$.

Assume that the relation (6.6) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)+1} (-1)^i J_{2k+2i+1}^2 &= J_{2k+4r+5}^2 - J_{2k+4r+7}^2 + \sum_{i=0}^{2r+1} (-1)^i J_{2k+2i+1}^2 \\ &= J_{2k+4r+5}^2 - J_{2k+4r+7}^2 + \sum_{i=0}^{2r+1} (-1)^i J_{2i+1}^2 - 8 J_{2k} [2 \eta_n^* J_{2k-2} + \beta_n^{**}] \\ &= \sum_{i=0}^{2(r+1)+1} (-1)^i J_{2i+1}^2 - 8 J_{2k} [2 \eta_{n+1}^* J_{2k-2} + \beta_{n+1}^{**}], \end{aligned}$$

where the last step uses Lemma 17. □

7 Alternating Jacobsthal products

Lemma 18. *For every $k \geq 0$ we have*

$$J_{2k+3} J_{2k+2} - J_{2k+1} J_{2k} = 3 + J_{2k} [120 J_{2k-2} + 49]. \quad (7.1)$$

Proof. (P2). $\frac{(P-1)(P+1)}{3} (49 M^2 - 40 P^2 + 9)$. □

Lemma 19. *For every $k \geq 0$ and every $n \geq 0$ we have*

$$\begin{aligned} J_{2k} (8 (\gamma_{n+1}^* - \gamma_n^*) J_{2k-2} + \tau_{n+1}^{***} - \tau_n^{***}) &= \\ J_{2k+4n+5} J_{2k+4n+4} - J_{2k+4n+3} J_{2k+4n+2} - J_{4n+5} J_{4n+4} + J_{4n+3} J_{4n+2}. \end{aligned} \quad (7.2)$$

Proof. (P1).

$$\frac{800}{3} \left((-1)^{1+2k} + 1 \right) AB + \frac{4}{3} \left((-1)^{4n+2k} - 1 \right) AC + \frac{160}{3} \left(4 (-1)^{4k} + 5 (-1)^{1+2k} + 1 \right) B + \frac{4}{3} \left((-1)^{2k} + (-1)^{4n+1} \right) C.$$

□

Let $\beta_0^{***} = 49$ and $\beta_{n+1}^{***} - \beta_n^{***} = 799 \cdot 2^{4n+4} + 25 \cdot 2^{4n+9} (2^{4n} - 1)$ for any $n = 0, 1, 2, \dots$

Lemma 20. *For every $k \geq 0$ and every $n \geq 0$ we have*

$$J_{2k} (8 (\eta_{n+1}^* - \eta_n^*) J_{2k-2} + \beta_{n+1}^{***} - \beta_n^{***}) = J_{2k+4n+6} J_{2k+4n+7} - J_{2k+4n+4} J_{2k+4n+5} + J_{4n+5} J_{4n+4} - J_{4n+6} J_{4n+7}. \quad (7.3)$$

Proof. (P1).

$$\frac{12800}{3} \left((-1)^{1+2k} + 1 \right) AB + \frac{16}{3} \left((-1)^{4n+2k} - 1 \right) AC + \frac{2560}{3} \left(4 (-1)^{4k} + 5 (-1)^{1+2k} + 1 \right) B + \frac{16}{3} \left((-1)^{2k} + (-1)^{4n+1} \right) C.$$

□

Theorem 6. *For every $m \geq 0$ and $k \geq 0$ the following equalities hold*

$$\sum_{i=0}^m (-1)^i J_{2k+2i} J_{2k+2i+1} = \sum_{i=0}^m (-1)^i J_{2i} J_{2i+1} + J_{2k} [8 \gamma_n^* J_{2k-2} + \tau_n^{***}], \quad (7.4)$$

if $m = 2n$ and $n = 0, 1, 2, \dots$ and

$$\sum_{i=0}^m (-1)^i J_{2k+2i} J_{2k+2i+1} = \sum_{i=0}^m (-1)^i J_{2i} J_{2i+1} - J_{2k} [8 \eta_n^* J_{2k-2} + \beta_n^{***}], \quad (7.5)$$

if $m = 2n + 1$ and $n = 0, 1, 2, \dots$

Proof of (7.4). (P3). For $n = 0$ the relation (7.4) is

$$J_{2k} J_{2k+1} = J_0 J_1 + J_{2k} [8 \gamma_0^* J_{2k-2} + \tau_0^{***}] = J_{2k} [8 J_{2k-2} + 3]$$

which is true by the relation (4.1) in Lemma 7 since $J_0 = 0$, $J_1 = 1$, $\gamma_0^* = 1$ and $\tau_0^{***} = 3$.

Assume that the relation (7.4) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)} (-1)^i J_{2k+2i} J_{2k+2i+1} &= \sum_{i=0}^{2r} (-1)^i J_{2k+2i} J_{2k+2i+1} + J_{2k+4r+4} J_{2k+4r+5} - J_{2k+4r+2} J_{2k+4r+3} \\ &= J_{2k+4r+4} J_{2k+4r+5} - J_{2k+4r+2} J_{2k+4r+3} + \sum_{i=0}^{2r} (-1)^i J_{2i} J_{2i+1} + J_{2k} [8 \gamma_n^* J_{2k-2} + \tau_n^{***}] \\ &= \sum_{i=0}^{2(r+1)} (-1)^i J_{2i} J_{2i+1} + J_{2k} [8 \gamma_{n+1}^* J_{2k-2} + \tau_{n+1}^{***}], \end{aligned}$$

where the last step uses Lemma 19.

□

Proof of (7.5). (P3). For $n = 0$ the relation (7.5) is

$$J_{2k} J_{2k+1} - J_{2k+2} J_{2k+3} = J_0 J_1 - J_2 J_3 - J_{2k} [8 \eta_0^* J_{2k-2} + \beta_0^{***}] = -3 - J_{2k} [120 J_{2k-2} + 49],$$

(i. e., the relation (7.1)) which is true by Lemma 18.

Assume that the relation (7.5) is true for $n = r$. Then

$$\begin{aligned} \sum_{i=0}^{2(r+1)+1} (-1)^i J_{2k+2i} J_{2k+2i+1} &= \sum_{i=0}^{2r+1} (-1)^i J_{2k+2i} J_{2k+2i+1} + J_{2k+4r+4} J_{2k+4r+5} - \\ J_{2k+4r+6} J_{2k+4r+7} &= J_{2k+4r+4} J_{2k+4r+5} - J_{2k+4r+6} J_{2k+4r+7} + \sum_{i=0}^{2r+1} (-1)^i J_{2i} J_{2i+1} - \\ J_{2k} [8 \eta_n^* J_{2k-2} + \beta_n^{***}] &= \sum_{i=0}^{2(r+1)+1} (-1)^i J_{2i} J_{2i+1} - J_{2k} [8 \eta_{n+1}^* J_{2k-2} + \beta_{n+1}^{***}], \end{aligned}$$

where the last step uses Lemma 20. □

References

- [1] Z. Čerin, Properties of odd and even terms of the Fibonacci sequence, *Demonstratio Math.*, **39** (2006), 55–60.
- [2] Z. Čerin, On sums of squares of odd and even terms of the Lucas sequence, *Proceedings of the 11th Fibonacci Conference*, to appear.
- [3] Z. Čerin, Some alternating sums of Lucas numbers, *Central European J. Math.* **3** (2005), 1–13.
- [4] Z. Čerin, Alternating sums of Fibonacci products, *Atti del Seminario Matematico e Fisico dell'Università di Modena e Reggio Emilia*, (to appear).
- [5] Z. Čerin and G. M. Gianella, On sums of squares of Pell-Lucas numbers, *INTEGERS: Electronic Journal of Combinatorial Number Theory*, **6** (2006), # A15. Available at <http://www.integers-ejcnt.org/vol6.html>.
- [6] Z. Čerin and G. M. Gianella, Formulas for sums of squares and products of Pell numbers, *Acc. Sc. Torino - Atti Sci. Fis.* **140** (2006), to appear.
- [7] A. F. Horadam, Jacobsthal representation numbers, *Fib. Quart.* **34** (1996), 40–54.
- [8] V. Rajesh and G. Leversha, Some properties of odd terms of the Fibonacci sequence, *Math. Gazette*, **88** (2004), 85–86.
- [9] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, 2006.

2000 *Mathematics Subject Classification*: Primary 11B39; Secondary 11Y55, 05A19.

Keywords: Jacobsthal numbers, Jacobsthal-Lucas numbers, integer sequences, sum of squares, combinatorial identity, Maple.

(Concerned with sequences [A001045](#) and [A014551](#).)

Received May 17 2006; revised version received January 19 2007. Published in *Journal of Integer Sequences*, January 19 2007.

Return to [Journal of Integer Sequences home page](#).