Journal of Integer Sequences, Vol. 19 (2016),
Article 16.7.2

Returns, Hills, and t-ary Trees

Helmut Prodinger
Department of Mathematical Sciences
Stellenbosch University
7602 Stellenbosch
South Africa
hproding@sun.ac.za

Abstract

A recent analysis of returns and hills of generalized Dyck paths is carried over to the language of t-ary trees, from which, by explicit bivariate generating functions, all the relevant results follow quickly and smoothly. A conjecture about the (discrete) limiting distribution of hills is settled in the affirmative.

1 Introduction

In a recent paper in this journal [2], *generalized Dyck paths* where investigated: they have an up-step $\mathsf{u} = (1, 1)$ and a down-step $\mathsf{d} = (1, -t + 1)$, where $t \geq 2$, start at the origin, end on the x-axis, and never go below the x-axis. A general reference for such lattice paths is an encyclopedic paper by Banderier and Flajolet [1].

Two parameters were investigated (with the help of Riordan arrays): the number of returns to the x-axis (the origin itself does not count), and the number of (contiguous) subpaths of the form $\mathsf{u}^{t-1}\mathsf{d}$, that sit on the x-axis.

In the present note, I would like to emphasize that the language of trees, in particular t-ary trees, is favorable here, because it allows one to write the relevant generating functions with ease, without any mentioning of Riordan arrays, and also leads to settling a conjecture mentioned in the recent paper mentioned before [2].

The family of t-ary trees is recursively described: such a tree is either an external node (depicted as a square), or a root (an internal node, depicted as a circle), followed by subtrees (in this order) $T_1, \dots, T_t$. For this and many other concepts, we refer to the universal book by Flajolet and Sedgewick [3]. The generating function $T(z) = \sum_{n\geq 0} a_n z^n$, where a_n is the number of trees of size n (n internal nodes) is, following the recursive definition, given by $T(z) = 1 + zT^t(z)$. Extracting coefficients is efficiently done by setting $z = u/(1+u)^t$, thus $T = 1 + u$, and contour integration; the method is closely related to the Lagrange inversion formula. Here is an example:

$$\begin{aligned}
[z^n]T^k(z) &= \frac{1}{2\pi i} \oint \frac{dz}{z^{n+1}} T^k(z) \\
&= \frac{1}{2\pi i} \oint \frac{du(1+u-tu)(1+u)^{t(n+1)}}{(1+u)^{t+1}u^{n+1}}(1+u)^k \\
&= [u^n](1+u-tu)(1+u)^{tn+k-1} \\
&= \binom{tn+k-1}{n} - (t-1)\binom{tn+k-1}{n-1} \\
&= \frac{k}{n}\binom{tn+k-1}{n-1}.
\end{aligned}$$

This produces in particular (for $k = 1$) the numbers $a_n = \frac{1}{n}\binom{tn}{n-1}$.

There is a natural bijection between the family of generalized Dyck paths and the family of t-ary trees. It is based on the decomposition of paths according to the *first* return to the x-axis. The first part of the Dyck paths is (recursively) responsible for the first $t-1$ subtrees, and the rest of the Dyck path for the remaining t-th subtree. It is then apparent that the number of down-steps is the same as the number of internal nodes of the associated tree. Here is the situation depicted for $t = 3$.



Figure 1: The decomposition of generalized Dyck paths leading (recursively) to a ternary tree with subtrees T_1, T_2, T_3.

Now a little reflection convinces us that the number of returns is the same as the number of (internal) nodes on the path from the root to the rightmost leaf. And, further: the number of hills is the number of nodes on this rightmost path with the property that its first $t-1$ subtrees are empty (are the empty subtree, consisting only of an external node).



Figure 2: A ternary tree with 10 (internal) nodes. It has 6 returns and 3 hills.

In what follows, we will analyze these parameters in terms of t-ary trees. In particular, we will freely speak about returns and hills of t-ary trees.

Cameron and McLeod [2], defined the *negative binomial distribution* via

$$\mathbb{P}\{Y = k\} = \binom{k-1}{r-1} p^r (1-p)^{k-r}.$$

This is somewhat in contrast with the book Analytic Combinatorics [3] and *Wikipedia*, as it is a shifted version, and the roles of p and $1-p$ are interchanged from the more common definitions. Nevertheless, we will stick to this definition here, for the reason of comparisons. The numbers r and p are called the parameters of the distribution.

2 The number of returns on t-ary trees

Let $F(z, v)$ be the generating function with respect to the size and the number of returns, i. e., the coefficient of $z^n v^k$ is the number trees with n internal nodes and k returns. Then we find the equation

$$F(z, v) = 1 + zT^{t-1}(z) v F(z, v).$$

Since $zT^{t-1}(z) = \frac{T(z)-1}{T(z)}$, this leads to the explicit form

$$F(z, v) = \frac{1}{1 - v\frac{T(z)-1}{T(z)}}.$$

Therefore

$$[v^k]F(z, v) = \Big(\frac{T(z)-1}{T(z)}\Big)^k = \Big(\frac{u}{1+u}\Big)^k.$$

Furthermore

$$\begin{aligned}
[z^n][v^k]F(z,v) &= [z^n]\Big(\frac{u}{1+u}\Big)^k \\
&= \frac{1}{2\pi i}\oint \frac{dz}{z^{n+1}}\Big(\frac{u}{1+u}\Big)^k \\
&= \frac{1}{2\pi i}\oint \frac{du(1+u-tu)(1+u)^{t(n+1)}}{u^{n+1}(1+u)^{t+1}}\Big(\frac{u}{1+u}\Big)^k \\
&= [u^{n-k}](1+u-tu)(1+u)^{tn-1-k} \\
&= \binom{tn-1-k}{n-k} - (t-1)\binom{tn-1-k}{n-1-k} \\
&= \frac{k}{n}\binom{tn-1-k}{n-k}.
\end{aligned}$$

Division by a_n gives the probability that a random tree of size n has k returns:

$$p_k(n) = k\frac{\binom{tn-1-k}{n-k}}{\binom{tn}{n-1}} \to \frac{k(t-1)^2}{t^{k+1}}, \quad \text{fixed } k, \quad n\to\infty.$$

In order to compute the d-th (factorial) moment, we evaluate

$$\frac{\partial^d}{\partial v^d}F(z,v)\Big|_{v=1} = d!T(z)(T(z)-1)^d = d!(1+u)u^d.$$

Furthermore,

$$\begin{aligned}
[z^n]\frac{\partial^d}{\partial v^d}F(z,v)\Big|_{v=1} &= [u^{n-d}]d!(1+u-tu)(1+u)^{tn} \\
&= d!\binom{tn}{n-d} - d!(t-1)\binom{tn}{n-1-d} = \frac{(td+1)d!}{n-d}\binom{tn}{n-1-d}.
\end{aligned}$$

For the expected value, we consider $d=1$ and divide by a_n, with the result

$$\frac{(t+1)n}{n(t-1)+2} \sim \frac{t+1}{t-1}.$$

The second factorial moment is obtained via $d=2$, with the result

$$\frac{2(2t+1)n(n-1)}{(tn-n+3)(tn-n+2)} \sim \frac{2(2t+1)}{(t-1)^2}.$$

This leads to the variance:

$$2\frac{n(t-1)(n-1)(tn+1)}{(tn-n+3)(tn-n+2)^2} \sim \frac{2t}{(t-1)^2}.$$

This section reproved and extended the results of [2] on the number of returns. Note that the quantity $\frac{k(t-1)^2}{t^{k+1}}$ is $\mathbb{P}\{Y=k+1\}$, where Y is a random variable, distributed according to the negative binomial distribution for $r=2$ and $p=\frac{t-1}{t}$.

3 The number of hills on t-ary trees

Let $G(z,v)$ be the generating function with respect to the size (variable z) and the number of hills (variable v). Then we find the recursion

$$G(z,v) = 1 + zT^{t-1}(z)G(z,v) + z(v-1)G(z,v).$$

Since $zT^{t-1}(z) = 1 - 1/T(z)$, we find the explicit solution

$$G(z,v) = \frac{T(z)}{1-(v-1)zT(z)} = \sum_{k\ge 0}(v-1)^k z^k T^{k+1}(z).$$

By d-fold differentation, followed by setting $v=1$, we get the generating function of the d-th factorial moments (apart from normalization):

$$d!z^dT^{d+1}(z).$$

Furthermore,

$$\begin{aligned}
[z^n]d!z^dT^{d+1}(z) &= \frac{d!}{2\pi i}\oint \frac{dz}{z^{n+1-d}}T^{d+1}(z)\\
&= \frac{d!}{2\pi i}\oint \frac{du(1+u-tu)(1+u)^{t(n-d)+d}}{u^{n+1-d}}\\
&= d!\,[u^{n-d}](1+u-tu)(1+u)^{t(n-d)+d}\\
&= d!\binom{tn-(t-1)d}{n-d} - d!(t-1)\binom{tn-(t-1)d}{n-1-d}\\
&= \frac{(d+1)!}{n-d}\binom{tn-(t-1)d}{n-1-d}.
\end{aligned}$$

For $d=1$, this leads to the expected value:

$$\frac{n}{\binom{tn}{n-1}}\frac{2}{n-1}\binom{tn-t+1}{n-2} = \frac{2(tn-t+1)!(tn-n+1)!}{t(tn-1)!(tn-n-t+3)!} \to \frac{2(t-1)^{t-2}}{t^{t-1}}.$$

The variance evaluates to

$$\frac{n}{\binom{tn}{n-1}}\frac{6}{n-2}\binom{tn-2t+2}{n-3} + \frac{2(tn-t+1)!(tn-n+1)!}{t(tn-1)!(tn-n-t+3)!} - \left[\frac{2(tn-t+1)!(tn-n+1)!}{t(tn-1)!(tn-n-t+3)!}\right]^2,$$

which we do not attempt to simplify any further.

Writing

$$G(z,v) = \sum_{n,k} g_{n,k}z^nv^k,$$

it is possible to derive an explicit form for the coefficients $g_{n,k}$, but they are not as nice as the corresponding quantities in the previous section:

$$\begin{aligned}G(z,v) &= \sum_{k\ge 0}(v-1)^k z^k T^{k+1}(z)\\ &= \sum_{n\ge 0} z^n \sum_{k\ge 0}(v-1)^k \frac{k+1}{n-k}\binom{tn-(t-1)k}{n-1-k}\\ &= \sum_{n\ge 0} z^n \sum_{k\ge 0}\sum_{0\le j\le k}\binom{k}{j}v^j(-1)^{k-j}\frac{k+1}{n-k}\binom{tn-(t-1)k}{n-1-k}.\end{aligned}$$

This leads to

$$g_{n,j} = \sum_{j\le k\le n}\binom{k}{j}(-1)^{k-j}\frac{k+1}{n-k}\binom{tn-(t-1)k}{n-1-k}.$$

The limiting distribution of $g_{n,j}/a_n$, for j fixed, must thus be determined in a different way.

We need a crash course in asymptotic tree enumeration here; all this can be found in Flajolet and Sedgewick's book [3], but compare also an older paper by Meir and Moon [4], in particular the notion of *simply generated families of trees*. The procedure that we describe here is closely related to the discussion in [3, Section IX-3], where very similar parameters were analyzed.

We start from $u = z\phi(u)$, with $\phi(u) = (1+u)^t$. The quantity τ is determined via the equation $\phi(\tau) = \tau\phi'(\tau)$. In our case this leads to $\tau = \frac{1}{t-1}$. Then there is the quantity $\rho = \frac{\tau}{\phi(\tau)}$, which here evaluates to

$$\rho = \frac{(t-1)^{t-1}}{t^t}.$$

Then one knows by general principles that the function $u(z)$ has a square-root singularity around $z=\rho$, with the local expansion

$$u \sim \tau - \sqrt{\frac{2\tau}{\rho\phi''(\tau)}}\sqrt{1-z/\rho}.$$

This is here

$$T(z) = 1+u \sim \frac{t}{t-1} - \sqrt{\frac{2t}{(t-1)^3}}\sqrt{1-z/\rho}.$$

This expansion will now be used inside of $G(z,v)$, with the result (Maple):

$$G(z,v) \sim a - \frac{\sqrt{2}t^{2t-3/2}}{(t-1)^{3/2}\big(t^{t-1}+(t-1)^{t-2}-(t-1)^{t-2}v\big)^2}\sqrt{1-z/\rho},$$

with a being an unimportant constant. Note that

$$\frac{\sqrt{2}t^{2t-3/2}}{(t-1)^{3/2}\big(t^{t-1}+(t-1)^{t-2}-(t-1)^{t-2}v\big)^2}\bigg|_{v=1} = \sqrt{\frac{2t}{(t-1)^3}}.$$

Thus, the limiting distribution is given by the probability generating function

$$\frac{t^{2t-2}}{\big(t^{t-1}+(t-1)^{t-2}-(t-1)^{t-2}v\big)^2} = \frac{\frac{t^{2t-2}}{(t^{t-1}+(t-1)^{t-2})^2}}{\Big(1-\frac{(t-1)^{t-2}}{t^{t-1}+(t-1)^{t-2}}v\Big)^2}.$$

The coefficient of v^k in it given by

$$(k+1)\frac{t^{2t-2}(t-1)^{(t-2)k}}{(t^{t-1}+(t-1)^{t-2})^{k+2}}.$$

which is $\mathbb{P}\{Y = k+2\}$, for a random variable Y, which follows the negative binomial distribution with parameters $r = 2$ and $p = \frac{t^{t-1}}{t^{t-1}+(t-1)^{t-2}}$, as conjectured in [2].

4 Acknowledgment

Thanks are due to Stephan Wagner for stimulating feedback.

References

[1] C. Banderier and P. Flajolet, Basic analytic combinatorics of directed lattice paths, *Theoret. Comput. Sci.* **281** (2002), 37–80.

[2] N. T. Cameron and J. E. McLeod, Returns and hills on generalized Dyck paths, *J. Integer Sequences* **19** (2016), Article 16.6.1.

[3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.

[4] A. Meir and J. W. Moon, On the altitudes of nodes in random trees, *Canad. J. Math.* **30** (1978), 997–1015.

2010 *Mathematics Subject Classification*: Primary 05A15, 05A16; Secondary 60C05.
Keywords: t-ary trees, Dyck paths, generating functions, negative binomial distribution, asymptotic tree enumeration.

(Concerned with sequences A001764, A006013, A006629, A006630, and A006631.)

Received June 27 2016; revised versions received August 26 2016; August 27 2016. Published in *Journal of Integer Sequences*, August 29 2016.

Return to Journal of Integer Sequences home page.